

For the use of a Registered Medical Practitioner or a Hospital or a Laboratory only

Risnia 1 mg Tablets

Risperidone tablets 1 mg

COMPOSITION

Each film coated tablet contains:

Risperidone Ph .Eur..... 1 mg

Colour: Titanium Dioxide

DOSAGE FORM

Oral Tablets

DESCRIPTION

White coloured, circular, biconvex film coated tablets, plain on both the sides.

Caution: This preparation may be habit forming on prolonged use

PHARMACOLOGY

Pharmacodynamics

Risperidone is a novel antipsychotic belonging to a new class of antipsychotic agents, the benzisoxazole-derivatives.

Risperidone is a selective monoaminergic antagonist with a high affinity for both serotonergic 5-HT₂ and dopaminergic D₂ receptors. Risperidone binds also to alpha₁-adrenergic receptors and, with lower affinity, to H₁-histaminergic and alpha₂-adrenergic receptors. Risperidone has no affinity for cholinergic receptors. Although risperidone is a potent D₂ antagonist, that is considered to improve the positive symptoms of schizophrenia, it causes less depression of motor activity and induction of catalepsy than classical neuroleptics. Balanced central serotonin and dopamine antagonism may reduce the tendency to cause extrapyramidal side effects, and extend the therapeutic activity to the negative and affective symptoms of schizophrenia.

Pharmacokinetics

Risperidone is completely absorbed after oral administration, reaching peak plasma concentrations within 1 to 2 hours. Food does not affect the absorption of risperidone from the stomach. The effect of food particles in the mouth on absorption has not been studied.

The most important route of metabolism of risperidone is hydroxylation by cytochrome CYP 2D6 to 9-hydroxy-risperidone which has a similar pharmacological activity to risperidone. This hydroxylation is subject to debrisoquine-type genetic polymorphism but this does not affect the active antipsychotic fraction since this consists of risperidone and its active metabolite 9-hydroxyrisperidone. After oral administration, the elimination half-life of the active antipsychotic fraction is 24 hours.

A single-dose study showed higher active plasma concentrations and a slower elimination of risperidone in the elderly and in patients with renal insufficiency. Risperidone plasma concentrations were normal in patients with liver insufficiency.

Topiramate modestly reduces the bioavailability of risperidone, but not that of the active antipsychotic fraction. Therefore, this interaction is unlikely to be of clinical significance. The bioavailability of topiramate is slightly decreased when administered in combination with risperidone. This interaction is not likely to be clinically significant.

INDICATIONS

Treatment of a broad range of patients with schizophrenia, including first episode psychoses, acute schizophrenic exacerbations, chronic schizophrenia, and other psychotic conditions, in which positive symptoms (eg, hallucinations, delusions, thought disturbances, hostility, suspiciousness, paranoia) and/or negative symptoms (eg, blunted affect, emotional and social withdrawal, speech difficulty) are prominent. Risperidone alleviates affective symptoms (eg, depression, guilt feelings, anxiety) associated with schizophrenia.

Risperidone is also effective in maintaining the clinical improvement during continuation therapy in patients who have shown an initial treatment response.

DOSAGE AND METHOD OF ADMINISTRATION

Schizophrenia:

Adults: Risperidone given as tablets, once or twice daily.

All patients whether acute or chronic, should start with 2 mg/day risperidone. The dosage may be increased on the 2nd day to 4 mg. From then on the dosage can be maintained unchanged or further individualised, if needed. Most patients will benefit from daily doses between 4 and 6 mg. In some patients, a slower titration phase and a lower starting and maintenance dose may be appropriate.

Doses >10 mg/day have not been shown to be superior in efficacy to lower doses and may cause extrapyramidal symptoms. Since the safety of doses >16 mg/day have not been evaluated, doses above this level should not be used.

A benzodiazepine may be added to risperidone when additional sedation is required.

Renal and Liver Disease: A starting dose of 0.5 mg twice daily is recommended. This dosage can be individually adjusted with 0.5 mg twice daily increments to 1 to 2 mg twice daily.

Risperidone should be used with caution in this group of patients until further experience is gained.

Children: Experience is lacking in children <15 years.

Elderly: A starting dose of 0.5 mg twice daily is recommended. This dosage can be individually adjusted with 0.5 mg twice daily increments to 1 to 2 mg twice daily. Risperidone is well tolerated by the elderly.

Method of administration

Risperidone is for oral use. Food does not affect the absorption of risperidone.

Upon discontinuation, gradual withdrawal is advised. Acute withdrawal symptoms, including nausea, vomiting, sweating, and insomnia have very rarely been described after abrupt cessation of high doses of antipsychotic medicines (see Undesirable Effects). Recurrence of psychotic symptoms may also occur, and the emergence of involuntary movement disorders (such as akathisia, dystonia and dyskinesia) has been reported.

Switching from other antipsychotics.

When medically appropriate, gradual discontinuation of the previous treatment while risperidone therapy is initiated is recommended. Also, if medically appropriate, when switching patients from depot antipsychotics, initiate risperidone therapy in place of the next scheduled injection. The need for continuing existing anti-Parkinson medicines should be re-evaluated periodically.

CONTRAINDICATIONS

Risperidone are contraindicated in patients with a known hypersensitivity to risperidone or any other ingredients in the product.

WARNINGS AND PRECAUTIONS

General

Elderly patients with dementia

Caution should be exercised and the risks and benefits of the combination of Risperidone and Furosemide or co-medication with other potent diuretics should be considered prior to the decision to use.

Irrespective of treatment, dehydration was an overall risk factor for mortality and should therefore be carefully avoided in elderly patients with dementia.

Cerebrovascular Adverse Events (CVAE)

Risperidone are not recommended for the treatment of behavioural symptoms of dementia because of an increased risk of cerebrovascular adverse events (including cerebrovascular accidents and transient ischaemic attacks). Treatment of acute psychoses in patients with a history of dementia should be limited to short term only and should be under specialist advice.

Physicians should consider carefully the risk of cerebrovascular adverse events with Risperidone (given the observations in elderly patients with dementia detailed above) before treating any patient with a previous history of CVA/TIA. Consideration should also

be given to other risk factors for cerebrovascular disease including hypertension, diabetes, current smoking, atrial fibrillation, etc.

Alpha-blocking activity

Due to the alpha-blocking activity of Risperidone, orthostatic hypotension can occur, especially during the initial dose-titration period. A dose reduction should be considered if hypotension occurs.

Risperidone should be used with caution in patients with known cardiovascular disease including those associated with prolongation of the QT interval and the dose should be gradually titrated. In clinical trials, Risperidone was not associated with an increase in QTc intervals. As with other antipsychotics, caution is advised when prescribing with medications known to prolong the QT interval.

If further sedation is required, an additional drug (such as a benzodiazepine) should be administered rather than increasing the dose of Risperidone

Tardive Dyskinesia/Extrapyramidal Symptoms (TD/EPS)

Drugs with dopamine receptor antagonistic properties have been associated with the induction of tardive dyskinesia, characterised by rhythmical involuntary movements, predominantly of the tongue and/or face. It has been reported that the occurrence of extrapyramidal symptoms is a risk factor for the development of tardive dyskinesia. If signs and symptoms of tardive dyskinesia appear, the discontinuation of all antipsychotic drugs should be considered.

Neuroleptic Malignant Syndrome (NMS)

Neuroleptic malignant syndrome, characterised by hyperthermia, muscle rigidity, autonomic instability, altered consciousness and elevated CPK levels, has been reported to occur with neuroleptics. In this event all antipsychotic drugs including risperidone should be discontinued.

It is recommended to halve both the starting dose and the subsequent dose increments in geriatric patients and in patients with renal or liver insufficiency.

Caution should also be exercised when prescribing Risperidone to patients with Parkinson's disease since, theoretically, it may cause a deterioration of the disease.

Hyperglycemia

Hyperglycaemia or exacerbation of pre-existing diabetes has been reported in very rare cases during treatment with Risperidone. Appropriate clinical monitoring is advisable in diabetic patients and in patients with risk factors for the development of diabetes mellitus (see also Undesirable effects).

Intraoperative Floppy Iris Syndrome

Intraoperative floppy iris syndrome (IFIS) has been observed during cataract surgery in patients treated with medicines with alpha1a-adrenergic antagonist effect, including risperidone (See Undesirable Effects). IFIS may increase the risk of eye complications during and after the operation. Current or past use of medicines with alpha1a-adrenergic antagonist effect should be made known to the ophthalmic surgeon in advance of surgery. The potential benefit of stopping alpha1 blocking therapy prior to cataract surgery has not been established and must be weighed against the risk of stopping the antipsychotic therapy.

Other

Classical neuroleptics are known to lower the seizure threshold. Caution is recommended when treating patients with epilepsy.

As with other antipsychotics, patients should be advised of the potential for weight gain. Acute withdrawal symptoms, including nausea, vomiting, sweating, and insomnia have very rarely been described after abrupt cessation of high doses of antipsychotic drugs. Recurrence of psychotic symptoms may also occur, and the emergence of involuntary movement disorders (such as akathisia, dystonia and dyskinesia) has been reported. Therefore, gradual withdrawal is advisable.

Use of Risperidone for schizophrenia in children aged less than 15 years has not been formally evaluated.

Hyperglycemia and Diabetes mellitus

Hyperglycemia in some cases extreme and associated with ketoacidosis or hyperosmolar coma or death, has been reported in patients treated with atypical antipsychotics. Assessment of the relationship between atypical antipsychotics use and glucose abnormalities is complicated by the possibility of an increased background risk of diabetes mellitus in patients with schizophrenia and the increasing incidence of diabetes mellitus in the general population. Given this confounders, the relationship between atypical antipsychotic use and hyperglycemia related adverse events is not completely understood. However, epidemiological studies suggest an increased risk of treatment emergent hyperglycemia-related events in patients treated with the atypical antipsychotics. Precise risk estimates for hyperglycemia –related adverse events in patients treated with atypical antipsychotics are not available.

Patient with an established diagnosis of diabetes mellitus who are started on atypical antipsychotics should be monitored regularly for worsening of glucose control. Patients with risk factors for diabetes mellitus (e.g. obesity, family history or diabetes) who are starting treatment with atypical antipsychotic should undergo fasting blood glucose testing at the beginning of treatment and periodically during treatment. Any patient treated with atypical antipsychotics should be monitored for symptoms of hyperglycemia including polydipsia, polyuria, polyphagia, and weakness. Patients who develop symptoms of hyperglycemia during treatment with atypical antipsychotics should undergo fasting blood glucose testing. In some cases, hyperglycemia has resolved when the atypical

antipsychotic was discontinued; however some patients required continuation of anti diabetic treatment despite discontinuation of the suspect drug.

DRUG INTERACTIONS

Possible interactions of Risperidone with other drugs have not been systematically evaluated. Given the primary CNS effects of risperidone, it should be used with caution in combination with other centrally acting drugs including alcohol.

Risperidone may antagonise the effect of levodopa and other dopamine-agonists.

Carbamazepine has been shown to decrease the plasma levels of the antipsychotic fraction of Risperidone. A similar effect might be anticipated with other drugs which stimulate metabolising enzymes in the liver. On initiation of carbamazepine or other hepatic enzyme-inducing drugs, the dosage of Risperidone should be re-evaluated and increased if necessary. Conversely, on discontinuation of such drugs, the dosage of Risperidone should be re-evaluated and decreased if necessary.

Phenothiazines, tricyclic antidepressants and some beta-blockers may increase the plasma concentrations of risperidone but not those of the active antipsychotic fraction. Fluoxetine and paroxetine, CYP2D6 inhibitors, may increase the plasma concentration of risperidone but less so of the active antipsychotic fraction. When concomitant fluoxetine or paroxetine is initiated or discontinued, the physician should re-evaluate the dosing of Risperidone. Based on *in vitro* studies, the same interaction may occur with haloperidol. Amitriptyline does not affect the pharmacokinetics of risperidone or the active antipsychotic fraction. Cimetidine and ranitidine increase the bioavailability of risperidone, but only marginally that of the active antipsychotic fraction. Erythromycin, a CYP 3A4 inhibitor, does not change the pharmacokinetics of risperidone and the active antipsychotic fraction. The cholinesterase inhibitor galantamine does not show a clinically relevant effect on the pharmacokinetics of risperidone and the active antipsychotic fraction. A study of donepezil in non-elderly healthy volunteers also showed no clinically relevant effect on the pharmacokinetics of risperidone and the antipsychotic fraction.

When Risperidone is taken together with other highly protein-bound drugs, there is no clinically relevant displacement of either drug from the plasma proteins.

See warnings and precautions regarding increased mortality in elderly patients with dementia concomitantly receiving furosemide.

Risperidone does not show a clinically relevant effect on the pharmacokinetics of valproate or topiramate. The potential for reduced toleration of the combination treatment should be taken into consideration when coadministering risperidone and topiramate.

In patients on long-term lithium and older/typical neuroleptic therapy, no significant change occurred in the pharmacokinetics of lithium after substitution of the concomitant neuroleptic with risperidone.

Food does not affect the absorption of risperidone from the stomach.

Pregnancy and Lactation

Pregnancy

Although, in experimental animals, risperidone did not show direct reproductive toxicity, some indirect, prolactin- and CNS-mediated effects were observed, typically delayed oestrus and changes in mating and nursing behaviour in rats. No teratogenic effect of risperidone was noted in any study. The safety of Risperidone for use during human pregnancy has not been established.

Neonates exposed to antipsychotic drugs during the third trimester of pregnancy are at risk for extrapyramidal and/or withdrawal symptoms following delivery. There have been reports of agitation, hypertonia, hypotonia, tremor, somnolence, respiratory distress and feeding disorder in these neonates. These complications have varied in severity, while in some cases symptoms have been self-limited, in other cases neonates have required intensive care unit support and prolonged hospitalization.

Risperidone 1 mg Tablets should be used in pregnancy only if the potential benefit justifies the potential risk to the foetus.

Lactation

In animal studies, risperidone and 9-hydroxyrisperidone are excreted in the milk. It has been demonstrated that risperidone and 9-hydroxyrisperidone are also excreted in human breast milk. Therefore, women receiving Risperidone should not breast feed.

UNDESIRABLE EFFECTS

Risperidone are generally well tolerated and in many instances it has been difficult to differentiate adverse events from symptoms of the underlying disease. Adverse events observed in association with the use of Risperidone include:

Common: insomnia, agitation, anxiety, headache.

Less common: somnolence, fatigue, dizziness, impaired concentration, constipation, dyspepsia, nausea/vomiting, abdominal pain, blurred vision, priapism, erectile dysfunction, ejaculatory dysfunction, orgasmic dysfunction, urinary incontinence, rhinitis, rash and other allergic reactions.

Cerebrovascular accidents have been observed during treatment with risperidone (see warnings and precautions).

Hyperglycemia and exacerbation of pre-existing diabetes have been reported in very rare cases during risperidone treatment.

The incidence and severity of extrapyramidal symptoms are significantly less than with haloperidol. However, in some cases the following extrapyramidal symptoms may occur: tremor, rigidity, hypersalivation, bradykinesia, akathisia, acute dystonia including oculogyric crisis. If acute in nature, these symptoms are usually mild and are reversible upon dose reduction and/or administration of antiparkinson medication, if necessary. In clinical trials in patients with acute mania risperidone treatment resulted in an incidence of EPS > 10%. This is lower than the incidence observed in patients treated with classical neuroleptics.

Occasionally, orthostatic dizziness, hypotension including orthostatic, tachycardia including reflex tachycardia and hypertension have been observed following administration of Risperidone.

Risperidone can induce a dose-dependent increase in plasma prolactin concentration. Possible associated manifestations are: galactorrhoea, gynaecomastia, disturbances of the menstrual cycle and amenorrhoea.

Weight gain, oedema and increased hepatic enzyme levels have been observed during treatment with Risperidone.

A decrease in neutrophil and/or thrombocyte count has been reported.

As with classical neuroleptics, rare cases of the following have been reported in schizophrenic patients: water intoxication with hyponatraemia, either due to polydipsia or to the syndrome of inappropriate secretion of antidiuretic hormone; tardive dyskinesia, body temperature dysregulation and seizures.

Benign pituitary adenomas have been reported very rarely in risperidone users during postmarketing surveillance. No causal association has been established.

Very rare cases of angioedema have been reported in postmarketing experience. Sedation has been reported more frequently in children and adolescents than in adults. In general, sedation is mild and transient.

Withdrawal reactions have been reported in association with antipsychotic drugs (see warnings and precautions).

Post marketing data

Eye disorders

Frequency: Not Known – Floppy iris syndrome (Intraoperative)

OVERDOSE

In general, reported signs and symptoms have been those resulting from an exaggeration of the drug's known pharmacological effects. These include drowsiness and sedation, tachycardia and hypotension, and extrapyramidal symptoms. In overdose, rare cases of QT-prolongation have been reported. In case of acute over dosage, the possibility of multiple drug involvement should be considered.

Establish and maintain a clear airway, and ensure adequate oxygenation and ventilation. Gastric lavage (after intubation, if the patient is unconscious) and administration of activated charcoal together with a laxative should be considered. Cardiovascular monitoring should commence immediately and should include continuous electrocardiographic monitoring to detect possible arrhythmias.

There is no specific antidote to Risperidone. Therefore appropriate supportive measures should be instituted. Hypotension and circulatory collapse should be treated with appropriate measures such as intravenous fluids and/or sympathomimetic agents. In

case of severe extrapyramidal symptoms, anticholinergic medication should be administered. Close medical supervision and monitoring should continue until the patient recovers.

SHELF-LIFE

24 months

STORAGE AND HANDLING INSTRUCTIONS

Store below 30°C.

Protect from light and moisture.

PACK SIZE

Carton having 10 Blister of 10 tablets each.

REGISTRATION HOLDER IN MALAYSIA

CIPLA MALAYSIA SDN BHD,
Suite 1101, Amcorp Tower
Amcorp Trade Centre,
18 Persiaran Barat ,
46 050 Petaling Jaya,
Selangor , Malaysia.

MANUFACTURED BY:-

Cipla Limited,
D 7, MIDC Industrial Area,
Kurkumbh,
Dist: Pune – 413 802,
India.

Date of Revision: August 2016