



VÉTOQUINOL - Malaisie	Nom du produit : Aurizon 10/20/30 ml (notice)	
Créé par : A. Milleret	Référence article : XXXXXX	0611 B
le : 27/06/11	Dimensions : 160x120 mm	
Visa :	1 couleur	rev B
 281 U		

How to prepare AURIZON® for use

-  Remove the cap from the container.
-  Separate the flexible cannula from its cap.
-  Push and screw the cannula onto the nozzle.
-  The cap protects the product from air after use.

How to apply AURIZON®

-  Shake bottle before application.
-  Lift the ear and position the cannula.
-  Squeeze the container gently.
-  (ear seen in section)
Aurizon will be released into the external ear canal.
-  Massage base of ear after application to distribute the drops evenly.

 **Vétoquinol**



AURIZON® EAR DROP

Ear drops, oily suspension.

Homogenous, beige to yellow suspension

QUALITATIVE AND QUANTITATIVE COMPOSITION

1 ml of AURIZON contains:

Marbofloxacin	3 mg
Clotrimazole Ph.Eur	10 mg
Dexamethasone acetate	1.0 mg
(equivalent to dexamethasone)	0.9 mg

List of excipients: *Propyl gallate, sorbitan oleate, colloidal silica, medium-chain triglycerides*

The product does not contain an antimicrobial preservative.

ATC vet code: Q502CA06

Pharmacological properties

Pharmacodynamic properties :

The preparation combines three active ingredients :

- **marbofloxacin**, a synthetic bactericidal agent belonging to the fluoroquinolone family that acts by inhibiting DNA gyrase. It exhibits a broad spectrum of activity against Gram-positive bacteria (e.g. *Staphylococcus intermedius*) and against Gram-negative organisms (*Pseudomonas aeruginosa*, *Escherichia coli* and *Proteus mirabilis*).

- **clotrimazole**, an anti-fungal agent that belongs to the imidazole family and which acts by causing changes in membrane permeability, allowing intracellular compounds to leak from the cell and thus inhibiting cellular molecular synthesis. It exhibits a wide spectrum of activity and is aimed, in particular, at *Malassezia pachydermatis*.

- **dexamethasone acetate**, a synthetic glucocorticoid exhibiting anti-inflammatory and anti-pruritic activity.

Pharmacokinetic properties :

Pharmacokinetics studies in dogs at the therapeutic dosage have shown that plasma concentrations of marbofloxacin peak at 0.06 µg/ml on the 14th day of treatment.

Marbofloxacin binds weakly to plasma proteins (<10% in dogs) and is eliminated slowly, mainly in the active form, 2/3 in urine and 1/3 in faeces. Clotrimazole absorption is extremely poor (plasma concentration <0.04 µg/ml).

Plasma concentration of Dexamethasone acetate reaches 1.25 ng/ml on the 14th day of treatment. The inflammatory process induced by otitis does not increase dexamethasone resorption.

Target species : Dogs

Indications for use

In dogs :

- Treatment of otitis externa of both bacterial and fungal origin respectively due to bacteria sensitive to marbofloxacin, and fungi, especially *Malassezia pachydermatis*, sensitive to clotrimazole. Use of the product should be based on susceptibility testing.

Contra-indications

Do not administer to dogs suffering from perforation of the tympanic membrane.
Do not administer to dogs with known hypersensitivity to any of the ingredients.
Do not administer to pregnant or lactating bitches.

Adverse reactions

Usual adverse reactions associated with corticosteroid drugs may be observed (changes in biochemical and haematological parameters, such as increase of alkaline phosphatase, and of aminotransferase, some limited neutrophilia).

Prolonged and intensive use of topical corticosteroid preparations is known to trigger local and systemic effects, including suppression of adrenal function, thinning of the epidermis and delayed wound healing.

On rare occasions, the use of this product may be associated with deafness, mainly in elderly dogs and mostly of a transient nature.

Special precautions for use in animals

Heavy reliance on a single class of antibiotic may result in the induction of resistance in a bacterial population. It is prudent to reserve fluoroquinolones for the treatment of clinical conditions which have responded poorly, or are expected to respond poorly, to other classes of antibiotics.

The external ear canal should be meticulously cleaned and dried before treatment.

Before treating with the product, the integrity of the tympanic membrane must be verified.

Avoid the introduction of contamination during use.

Use during pregnancy and lactation

The safety of the product has not been demonstrated in pregnant and lactating bitches.

Interaction with other medicaments and other forms of interaction

None known.

Dosage for each species, route and method of administration

Shake well before use. Avoid the introduction of contamination during use.

Apply ten drops into the ear once daily for 7 to 14 days.

After 7 days of treatment, the veterinary surgeon should evaluate the necessity to extend the treatment a further week.

One drop of the preparation contains 71µg marbofloxacin, 237µg clotrimazole and 23.7µg dexamethasone acetate.

After application, the base of the ear may be massaged briefly and gently to allow the preparation to penetrate to the lower part of the ear canal.

When the product is intended for use in several dogs, use one cannula per dog.

Overdose (symptoms, emergency procedures, antidotes)

Changes in biochemical and haematological parameters (such as increase of alkaline phosphatase, aminotransferase, some limited neutrophilia, eosinopenia, lymphopenia) are observed with three fold the recommended dosage; such changes are not serious and will reverse once the treatment has stopped.

Special warnings for each target species : None.

Withdrawal period : Not applicable.

Special precautions to be taken by the person administering the product to animals

Wash hands carefully after applying the product.

Avoid contact with eyes. If splashed in the eye, rinse with copious amounts of water.

Persons with known hypersensitivity to compounds in the product should avoid any contact with the product.

Incompatibilities : Not applicable.

Shelf-life

2 years.

After opening: 2 months.

When the vial is opened for the first time, using the in use shelf life of 2 months, the date on which any product remaining in the container is to be discarded should be worked out. This discard date should be written in the space provided on the label.

Special storage precautions : Do not store above 30°C.

Nature and contents of container

Details of the primary packaging:

- Low-density polyethylene bottle.
- Low-density polyethylene nozzle.
- Threaded polypropylene cap.
- PVC cannula.

Presentation

- Box containing 1 x 10 ml bottle and 1 cannula
- Box containing 1 x 20 ml bottle and 2 cannulae

Date on which the package leaflet was last approved: December 2005

Registration's holder:

Gladron Chemicals Sdn. Bhd.

No.7, Jalan TP7,

UEP Industrial Park,

40400 Shah Alam,

Selangor Darul Ehsan.

Name and address of the manufacturer and site of batch release

VÉTOQUINOL

Magny-Vernois

70200 LURE

France

Special precautions for the disposal of unused product or waste material

Any unused product or waste material should be disposed of in accordance with national requirements.

UK: dispose of used packaging in the household refuse. Unused product should be returned to the veterinary surgeon.

Keep the container in the outer carton

**PRESCRIPTION ONLY MEDICINE
MARKETING AUTHORISATION NUMBERS:**

Registration number: MALXXXXX

**VETERINARY MEDICINAL PRODUCT AUTHORISED FOR USE IN UK AND IE
FOR ANIMAL TREATMENT ONLY
KEEP OUT OF REACH OF CHILDREN / JAUHI DARIPADA KANAK-KANAK
FOR EXTERNAL USE ONLY**

XXXXXX

0611 B