

Vit C 100-UD Chewable Tablet

Name of Product & Dosage Form

Vit C 100-UD Chewable Tablet

Composition

Each chewable tablet contains Ascorbic Acid 100 mg.

Description

Capsule shaped, pink tablets, bisected on one side and embossing "Atlantic" on the other side.

Size: length 11.9 – 12.7 mm, width 6.3 - 6.7 mm, thickness 4.7 – 5.4 mm.

Indication

As a dietary supplement.

Mode of administration

For oral administration.

Chew the tablet or allow to dissolve slowly in the mouth. Preferably after meals.

Recommended Dosage

Children (over 3 years): Half to one tablet per day (50 - 100 mg/day)

Adults : 1 - 5 tablets per day (100 - 500 mg/day)

Contraindications

Hypersensitivity to the ingredients.

Warning and Precautions

Patient with diabetes, prone to recurrent renal calculi (e.g. dialysis patients), undergoing stool occult blood tests and on sodium-restricted diets should not take excessive doses for extended periods of time. In large doses, ascorbic acid should be given with care to patients with hyperoxaluria.

Prolonged use of large doses of vitamin C may experience symptoms of deficiency of vitamin C when resume to normal intake. Excessive use of chewable vitamin C may lead to erosion of tooth enamel. Large dose of ascorbic acid have resulted in haemolysis in patients with G6PD deficiency.

Use in pregnancy and lactation

Pregnancy: The possibility of fetal harm appears remote. However, use during pregnancy only if clearly needed.

Lactation: Use with caution in nursing women.

Side Effects/Adverse Reactions

Ascorbic acid is usually nontoxic, however, nausea, vomiting, heartburn, abdominal cramps, fatigue, flushing, headache, insomnia, and sleepiness have been reported.

Drug Interactions

- ◆ Increased effect: Ascorbic acid enhances iron absorption in the GI tract. Concomitant ascorbic acid taken with oral contraceptives may increase contraceptive effect.
- ◆ Decreased Effect: Ascorbic acid and fluphenazine may decrease fluphenazine levels. Ascorbic acid and warfarin may decrease anticoagulant effect. Changes in dose of ascorbic acid when taken with oral contraceptives may reduce the contraceptive effect.
- ◆ Aspirin, nicotine, alcohol, phenytoin, several appetite suppressants and some anticonvulsant drugs may decrease blood levels of ascorbic acid.

Signs and Symptoms of overdose and treatment

Gastrointestinal distress may indicate toxicity. Forced fluids and diuresis may be beneficial in cases of toxicity due to massive ingestions of vitamin C.

Storage condition

Store below 25°C and protect from light.

Keep out of reach of children.

Jauhi daripada kanak-kanak.

Packaging available

1. HDPE bottle and/or amber glass bottle of 100 tablets per bottle. Nine bottles are packed in a paper box.
2. Zipped-lock aluminium sachet of 100 tablets per sachet. Nine sachets are packed in a paper box.

Manufactured for:

The United Drug (1996) Co., Ltd.
Bangkok, Thailand.

By :

Atlantic Laboratories Corporation. Ltd.
2038 Sukhumvit Road,
Bangkok, Thailand.

Revision date: April 2013