

ABOUT BLACKMORES' VITAMIN D3 1000 IU

INDICATION:

- Treatment and prevention of vitamin D deficiency
- Adjuvant treatment for osteoporosis

Pharmacodynamics and pharmacokinetics

Vitamin D3 is generally produced by the body when skin is exposed to sunlight. Individuals are at high risk of vitamin D deficiency when there is decreased natural production of vitamin D3 due to limited sun exposure, dark skin, protective clothing, traditional or religious clothing that covers most of the body, use of sunscreen and old age. Malabsorptive gastrointestinal disease and certain medications can be other factors contributing to vitamin D3 deficiency.

Pharmacodynamics

Vitamin D3, also known as cholecalciferol or calciol, is a

fat-soluble vitamin that is a positive regulator of calcium homeostasis. It is a form of vitamin D that enhances absorption of calcium from the small intestine, helps mobilise calcium from the bones and acts on the kidney to help your body retain calcium and phosphorus. Vitamin D3 has a slow onset and long duration of action.

Pharmacokinetics

- **Absorption:** Vitamin D3 is absorbed from the small intestine. The presence of bile is essential for adequate intestinal absorption.
- **Distribution:** Vitamin D3 circulates in the blood and is stored for long periods of time in adipose and muscle tissue from where it is slowly released.
- **Metabolism:** Vitamin D3 is hydroxylated in the liver by the enzyme vitamin D 25-hydroxylase to form 25-hydroxyvitamin D, which is distributed in the circulation. In the kidneys it is further hydroxylated by the

enzyme vitamin D 1-hydroxylase to form the biologically-active metabolite 1,25-dihydroxy vitamin D.

- **Excretion:** It is mainly excreted via the bile and faeces and only small amounts are eliminated in the urine.

HOW TO USE:

For the treatment and prevention of Vitamin D3 deficiency: Adults - Take 1-2 capsules a day with a meal, or as professionally prescribed. Children under 12 years - as professionally prescribed.

For adjuvant treatment for osteoporosis: Adults - Take 1-2 capsules a day with a meal, or as professionally prescribed.

CONTRAINDICATIONS, SIDE EFFECTS, PRECAUTIONS AND WARNINGS:

Contraindications:

- Hypersensitivity to vitamin D. Vitamin D3 should not be given to patients with hypercalcaemia, several renal

impairment and metastatic calcification.

Precautions & Warnings:

- Arteriosclerosis (potential exacerbation related to persistent hypercalcemic effects during therapeutic use). Cardiac conditions (potential exacerbation related to persistent hypercalcemic effects during therapeutic use). Concurrent use of calcium-containing preparations, other vitamin D-containing preparations or vitamin D analogs, or thiazide diuretics (enhanced risk of hypercalcaemia). Hyperlipidemia (potential exacerbation of LDL elevation). Hyperphosphatemia (risk of metastatic calcification; normalization of phosphate levels indicated prior to therapy). Liver disease (impaired absorption). Renal impairment (potential exacerbation related to hypercalcemic effects during therapeutic use). Renal osteodystrophy/ other conditions requiring high doses of pure vitamin D (vitamin D metabolites preferred, such as

XXXX

calciolol). Sarcoidosis/other granulomatous disease (potential for enhanced sensitivity to vitamin D).

Side Effects:

- Oral vitamin D is generally well tolerated in recommended doses, with the most common adverse effect being gastrointestinal complaints. Excessive consumption of high doses above 60,000 IU per day may cause hypercalcaemia and its associated effects including hypercalciuria, ectopic calcification, renal damage and cardiovascular damage. Hypersensitivity reactions have occurred.

Medicine Interactions:

- There is an increased risk of hypercalcaemia if vitamin D is given with thiazide diuretics, calcium or phosphate. Plasma-calcium concentrations should be monitored in such situations. Some antiepileptics may increase vitamin D requirements (e.g. carbamazepine, phenobarbital, phenytoin, and primidone). Rifampicin and isoniazid may

reduce the effectiveness of vitamin D. Corticosteroids may counteract the effect of vitamin D.

Pregnancy and Lactation:

- Use under the guidance of your doctor or pharmacist in the event you are pregnant or lactating. Due to lack of human safety information, cholecalciferol should be used in pregnant women only if the potential benefit outweighs potential risk to the foetus. Cholecalciferol, along with some of its active metabolites, passes into breast milk. Due to lack of human safety information, caution is advised when using cholecalciferol in the nursing mother. The importance of therapy to the mother should be taken into consideration.

Symptoms of overdose and treatment:

- Doses of 60,000 IU/day for long periods can cause hypercalcaemia with muscle weakness, apathy, headache, anorexia, nausea and vomiting, bone pain, ectopic calcification, proteinuria, hypertension and cardiac

arrhythmias. An overdose should be treated symptomatically. Vitamin D supplementation should be discontinued and a low-calcium diet commenced. Calcium levels, ECG and renal function should be checked. Supportive management under expert medical supervision is required.

Storage instructions:

- Keep out of reach of children. Store below 30°C in a dry place away from direct sunlight. Do not use the product beyond the expiry date printed on the pack.

ACTIVE INGREDIENT PER CAPSULE:

Vitamin D3 1000 IU equivalent to Cholecalciferol 25 mcg

Description: Oval, clear natural soft capsule filled with pale yellow oil. Unprinted.

Pack size: 60 capsules

ABOUT BLACKMORES:

Malaysia: Product registration holder: Blackmores (Malaysia) Sdn Bhd., Unit No.7-05-01, Level 5, Tower 7, UOA Business Park, No.1 Jalan Pengaturcara UI/51A Section UI, 40150 Shah Alam, Selangor, MALAYSIA. For more information ☎ 03-5022 8263

Contract manufactured for Blackmores by: Catalent Australia Pty Ltd, 217-221 Governor Road, Braeside VIC 3195 AUSTRALIA

Packed by: Blackmores Ltd., 20 Jubilee Avenue, Warriewood NSW, 2102 AUSTRALIA.

Date of amendment: June 2026

XXXX

BASE

BLACKMORES

Label Size: 160 x 40 mm

Text Height: 1.5mm

NB: TGA Require 1.5mm min. text height

CMF Job No: XXXX

New label Item Code: XXXXX

New Spec No: XXXXXX XXX

NB: PHARMA CODE are last 3 digits of New Spec No.

PRODUCT NAME: BLACKMORES® VITAMIN D3 1000 IU (60) MY DLAB

Draft No: 7

Active Proof No: -

Date: 23.06.26

EAN No: 93542524

100 ml Amber Glass Bottle

Colours used:

CYAN

MAGENTA

BLACK

PANTONE 7711

PANTONE 1665

5

Colours

LABEL SPECIFICATION:

This label is a Blackmores standard label (MCC).

PRINTER DETAILS:

PRINTER MCC

SENT TO PRINT