

Front

Back



Cloderm Shampoo 0.05% w/w

DESCRIPTION: A clear yellow viscous liquid with characteristic odour containing Clobetasol Propionate 0.05% w/w with Methyl Paraben 0.1 % w/w and Propyl Paraben 0.05% w/w as preservatives.

PHARMACODYNAMICS: Clobetasol propionate has anti-inflammatory, anti-pruritic, and vasoconstrictive actions.

PHARMACOKINETICS: When administered topically, particularly under occlusive dressings or when the skin is broken, sufficient corticosteroid may be absorbed to give systemic effects. Corticosteroids are extensively bound to plasma proteins. Only unbound corticosteroids have pharmacological effects or are metabolized. They are metabolized mainly in the liver, also in the kidneys, and are excreted in the urine.

INDICATIONS: Psoriasis and recalcitrant eczemas of the scalp. Clobetasol propionate is a highly active topical corticosteroid which is indicated for use in short courses for conditions which do not respond satisfactorily to less active steroids.

CONTRAINDICATIONS: Use of Cloderm Shampoo is contraindicated in patients who are hypersensitive to clobetasol propionate, to other corticosteroids, or to any ingredient in this preparation.

WARNINGS AND PRECAUTIONS: Discontinue use if irritation or sensitivity develops. Avoid introduction of shampoo into the eyes. Special care must be taken when giving to paediatric patients as systemic absorption can occur in topical administration causing growth retardation. Caution is advised if the medication is to be applied to extensive surface area. Long-term continuous topical therapy should be avoided. Not suitable for ophthalmic use. Avoid contact with eyes.

ROUTE OF ADMINISTRATION: Topical.

INTERACTIONS WITH OTHER MEDICAMENTS: Unknown.

PREGNANCY AND LACTATION: Safety of its use during pregnancy and lactation has not been established. Thus it should be used only if the potential benefit outweighs the potential risk to the foetus or nursing infant.

ADVERSE EFFECTS: Reported local adverse reactions are burning, itching, irritation, dryness, folliculitis, hypertrichosis, acneiform eruptions, hypopigmentation, maceration of the skin, secondary infection, skin atrophy, striae and miliaria. Large doses of corticosteroids may produce symptoms typical of hyperactivity of the adrenal cortex, with moon face, sometimes with hirsutism, buffalo hump, flushing, sometimes leading to a fully developed Cushing's syndrome.

RECOMMENDED DOSAGE: Cloderm Shampoo should be applied onto dry (not wet) scalp once a day in a thin film to the affected areas only, and left in place for 15 minutes before lathering and rinsing. Treatment should be limited to 4 weeks. The total dosage should not exceed 50g per week. Not suitable for patients under 18 years old.

SYMPTOMS AND TREATMENT FOR OVERDOSAGE: Corticosteroid applied to the skin can be absorbed in sufficient amount to produce systemic effect such as hypothalamic-pituitary-adrenal axis suppression, manifestation of Cushing's syndrome, hyperglycaemia and glucosuria. Tests which may be helpful in evaluating hypothalamic-pituitary-adrenal axis suppression include urinary free cortisol test and ACTH stimulation test. If the hypothalamic-pituitary-adrenal axis suppression is found, then the drug should be withdrawn, frequency of application reduced or a weaker steroid used. Supplemental systemic corticosteroids may be required if signs and symptoms of steroid withdrawal occur.

PACKING: Plastic bottle of 60ml, 100ml.

STORAGE: Store below 30°C. Keep out of the reach of children. Keep container well closed. Protect from light.

RECOMMENDED SHELF LIFE: 3 years.

Manufactured by:

HOE Pharmaceuticals Sdn. Bhd.
Lot 10, Jalan Sultan Mohamed 6, Bandar Sultan Suleiman,
42000 Port Klang, Selangor, MALAYSIA.

Revision date: September 2022