BAYER ARTWORK PANEL

Bayer Ref No:	**BAY526047**
Trident Reference No:	TR2536251
Brand:	Canesten
Range:	Cream
Product:	1 Percent Cream
Pack Type:	Leaflet
Pack Size:	10G/20G Pack
Action:	**A**
Date:	22/06/22
Country:	Malaysia/Brunei
Finished Goods Code:	**80435665 80435700**
Synaps Request Code:	**TL028759**
Component Code:	**CMY5/05**
CAD Ref No:	Pil 148 x 210 [Folding size 148 x 26] Encube
Printer:	AMR Print & Pack (Bhiwandi, India)
Packaging Site:	Encube Ethicals PVT Ltd [IN]
Substrate:	White Paper
Barcode 1 Type:	N/A
Barcode 1 Mag:	N/A
Barcode 2 Type:	N/A
Barcode 2 Mag:	N/A
Pharmacode No/NE:	0000110(133)

Technical & Non Printing Items

Cutter Guides

Colours

PANTONE 485 C	PANTONE Cool Gray 11 C
1 LTH	2 LTH

TRIDENT

Connaught House, Connaught Road,
Kingswood Business Park, Hull, HU7 3AP, England.
T: +44 (0) 1482 828100

Please note that any low resolution paper Canon colour copies associated with this job should be referred to for content, layout and colour separation only.

UNDER NO CIRCUMSTANCES SHOULD THIS ARTWORK BE ALTERED WITHOUT PRIOR PERMISSION FROM TRIDENT.

STUDIO USE ONLY	Divya Pandiyan	v1.0
Sm•Art check results:	G=1; O=2; R=0; - DP - 22/06/22 12:55:04	

FONT SIZE PANEL

Main Text (body copy):	8 pt
Height (based on lower case x):	1.5 mm

ARTWORK APPROVAL PANEL

Packaging Site Approval:

Name:	
Date:	
Action:	

Country Approval:

Name:	
Date:	
Action:	

148 mm

10 mm 10 mm

26 mm 26 mm

210 mm

Direction of reading

Direction of reading

Canesten®

Clotrimazole Cream 1% w/w

Broad-Spectrum Antimycotic with Fungicidal Action For local application only

Clotrimazole Cream 1% w/w

Read all of this leaflet carefully before you take this medicine.
Keep this leaflet. You may need to read it again. If you have further questions, ask your doctor or pharmacist. If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

What is in this leaflet

1. What Canesten® Clotrimazole Cream 1% w/w is used for
2. How Canesten® Clotrimazole Cream 1% w/w works
3. Before you use Canesten® Clotrimazole Cream 1% w/w
4. How to use Canesten® Clotrimazole Cream 1% w/w
5. While you are using it
6. Side effects
7. Storage and disposal of Canesten® Clotrimazole Cream 1% w/w
8. Product description

What Canesten® Clotrimazole Cream 1% w/w is used for
Canesten® Clotrimazole Cream 1% w/w is used to treat fungal skin infections such as athlete's foot, white spots, ringworm, fungal infection of the tissues adjacent to the nails, fungal infection of the skin folds & erythrasma (skin redness/irritation).

How Canesten® Clotrimazole Cream 1% w/w works
The active substance in Canesten® Clotrimazole Cream 1% w/w is clotrimazole. Clotrimazole belongs to a group of medicines called imidazoles. The primary mechanism of action of clotrimazole is against the division and growing of fungi. Clotrimazole alters the permeability of the fungal cell wall and inhibits the activity of enzymes within the cell. It specifically inhibits the biosynthesis of ergosterol required for cell membrane production.

Before you use Canesten® Clotrimazole Cream 1% w/w
When you must not use it
If you are allergic (hypersensitive) to clotrimazole or any of the other ingredients, including cetostearyl alcohol.

Before you start to use it
If you are pregnant, breast-feeding or trying for a baby, tell your doctor or midwife before using Canesten® Clotrimazole Cream 1% w/w.
If you have informed your doctor or midwife already, follow his/her instructions carefully. As a precautionary measure, it is preferable to avoid the use of clotrimazole during the first trimester of pregnancy. Breast-feeding should be discontinued during treatment with clotrimazole.

Taking other medicines
Not known.

How to use Canesten® Clotrimazole Cream 1% w/w
How much to use
Before use, pierce the tube seal by inverting the cap over the end of the tube and press. If the feet are infected, they should be washed and dried thoroughly especially between the toes before applying the cream. Canesten® Clotrimazole Cream 1% w/w should be applied thinly and evenly to the affected areas two or three times daily and rubbed in gently. A strip of cream (1/2 cm long) is enough to treat an area of about the size of the hand. The symptoms of skin infection, such as itching or soreness, should improve within a few days of treatment, although signs such as redness and scaling may take longer to disappear. If symptoms persist, consult your doctor. Patients should notify their physician if there is no improvement after 4 weeks of treatment.

5 mm 5 mm

5 mm 5 mm

148 mm

5 mm 5 mm

210 mm

5 mm 5 mm

When to use it
It can be applied anytime of the day.

How long to use it
To ensure complete healing depending on the indication the treatment should be continued according to the indications as specified below even if symptoms disappear.

Treatment duration
Athlete's foot, ringworm, fungal infection of the tissues adjacent to the nails, fungal infection of the skin folds : 3-4 weeks or as directed by physicians.
Erythrasma (skin redness/irritation) : 2-4 weeks or as directed by physicians.
White spots : 1-3 weeks or as directed by physicians.

If you forget to use it
Apply the cream as soon as possible and then continue the rest of your treatment as usual.

If you use too much (overdose)
No reports are available on cases of overdose with Canesten® Clotrimazole Cream 1% w/w. There is no specific antidote. Topical clotrimazole is intended for cutaneous use only. In the case of cutaneous overdose patients should be managed by symptomatic care; patients should wash off the product with water and cleanser, discontinue use, and see their physician so appropriate therapy can be implemented as needed. In the case of ingestion of an overdose, patients should see their physician and be managed by symptomatic and supportive care.

While you are using it
Things you must do
You can help the treatment to work if you follow these simple self-help tips:
- Keep the affected skin areas clean.
- Pay particular attention to drying the skin, but avoid excessive rubbing.
- Do not share towels, bath mats, etc. with other people as you could spread the infection to them.
- Always wash your hands after treating the infection to prevent it from spreading.

If you have athlete's foot:
- Remember to dry the skin between the toes thoroughly.
- Wash your socks, stockings and tights thoroughly in hot water to remove any shed skin or fungal spores.
- Change your footwear daily if possible.

Things you must not do
Canesten® Clotrimazole Cream 1% w/w is for external use only:
Do not put the cream in your mouth or swallow it.
Although the infected area will itch, try not to scratch. Scratching will damage the surface of the skin and cause the infection to spread further.

Things to be careful of
This product contains cetostearyl alcohol which may cause local skin irritation (e.g. rash, itching or redness).

Side effects
Like all medicines, Canesten® Clotrimazole Cream 1% w/w can cause side effects, although not everybody gets them. As with all medicines, some people may be allergic to the cream. If you are allergic, a reaction will occur soon after you start using it.

If you experience an allergic reaction, stop using Canesten® Clotrimazole Cream 1% w/w.

Signs of an allergic reaction may include:
- Rash.
- Swallowing or breathing problems.
- Swelling of your lips, face, throat or tongue.
- Weakness, feeling dizzy or faint.
- Nausea.
- Discomfort/pain.

After you apply the cream you might experience:
- Itching, rash, blisters, burning, discomfort, swelling, irritation or peeling of skin.

If you experience any of the above effects or react badly to the cream in any other way not listed in this leaflet, tell your doctor or pharmacist immediately.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my
[Consumers -> Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of Canesten® Clotrimazole Cream 1% w/w
Storage
To be stored below 30ºC.
Keep out of reach of children.

Disposal
Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product description
What it looks like
Canesten® Clotrimazole Cream 1% w/w is a white smooth cream.

Ingredients
- Active ingredient : clotrimazole 1% w/w.
- Inactive ingredients : sorbitan monostearate, polysorbate 60, cetyl palmitate, cetostearyl alcohol, octyldodecanol, benzyl alcohol & purified water.

MAL number : MAL14015020XCRZ

Manufacturer
Encube Ethicals Private Limited
C-1, Madkaim Industrial Estate,
Madkaim, Ponda, Goa, 403404 India
For Bayer Consumer Care AG Basel, Switzerland

Product Registration Holder
Bayer Co. (Malaysia) Sdn Bhd
25-03 & 25-04, Level 25, IMAZIUM,
No. 8, Jalan SS21/37, Damansara Uptown,
47400 Petaling Jaya, Selangor

Date of revision : 01 June 2022

M.L.No. 361

Bayer
CMY5/05

REBRANDING CONFORM