DEXATROL EYE DROPS

Dexamethasone 1.0mg
Neomycin (as sulphate) equivalent to Neomycin 3.5mg
Polymyxin B-sulphate 6000IU

Product Name

DEXATROL EYE DROPS

Product Description

White suspension in flexible plastic dropper bottle. This product can be used for 28 days after the first opening when conserved at room temperature below 25°C.

Composition

Each ml contains Dexamethasone 1.0mg, Neomycin sulphate equivalent to Neomycin 3.5mg and Polymycin B-sulphate 6000IU.

Pharmacodynamics

Corticoids suppress the inflammatory response to a variety of agents and they probably delay or slow healing.

Since corticosteroids may inhibit the body's defense mechanism against infection, a concomitant antimicrobial drug may be used when this inhibition is considered to be clinically significant in a particular case.

The anti-infective component in the combination is included to provide action against specific organism susceptible to it.

Neomycin sulfate is considered active against the following microorganisms:

Staphylococcus aureus, Corynebacterium diptheriae, Streptococcus viridans, Escherichia coli, Klebsiella pneumoniae, Proteus vulgaris, Aerobacter aerogenes, and Haemophilus influenza. Polymyxin B Sulfate is considered active against the following microorganisms: Pseudomonas aeruginosa, Aerobacter aerogenes, Escherichia coli, Klebsiella pneumoniae and Koch-Weeks bacillus.

Pharmacokinetics

When a decision to administer both a corticoid and an antimicrobial is made the administration of such drugs in combination has the advantage of greater patient compliance and convenience, with the added assurance that the appropriate dosage of both drugs is administered, plus assured compatibility of ingredients when types of drugs are in the same formulation and, particularly, that the correct volume of the drug is delivered and retained.

The relative potency of corticosteroids depends on the molecule structure, concentration and release from the vehicle.

Indication

Dexatrol Eye Drops indicated for local treatment for steroid-responsive inflammatory ocular conditions for which a corticosteroid is indicated and where bacterial infection or a risk of bacterial ocular infection exists.

Recommended Dosage

For topical ophthalmic use only.

One or two drops in the conjunctival sac(s).

In severe disease, drops may be used hourly, being tapered to discontinuation as the inflammation subsides.

In mild disease, drops may be used up to four to six times daily.

Contraindications

Dexatrol Eye Drops must not be used in patient with:-

- Epithelial herpes simplex keratitis (dendritic keratitis), vaccinia, varicella, and many other viral diseases of the cornea and conjunctiva.
- Mycobacterial infection of the eye.
- Fungal diseases of ocular structures.
- Hypersensitivity to a component of the medication.

Warnings and Precautions

Warnings:

Not for injection.

Do not touch dropper tip to any surface, as this may contaminate the contents. Prolonged use may result in glaucoma with damage to the optic nerve, defects in visual acuity and fields of vision, and posterior subcapsular cataract formation.

Prolonged use may suppress the host response and thus increase the hazard of formation. In those diseases causing thinning of the cornea or sclera, perforation have been known to occur with the use of topical corticosteroids.

In acute purulent conditions of the eye, corticosteroids may mask infection or enchance existing infection.

If this product is used for 10 days or longer, intraocular pressure should be routinely monitored even though it may be difficult in children and uncooperative patients.

Employment of corticosteroid medication in the treatment of herpes simplex requires great caution.

The possibility of persistent fungal infections of the cornea should be considered after prolonged corticosteroid dosing.

Interactions with Other Medicaments

No relevant information on topical use.

Pregnancy and Lactation

There are no adequate or well-controlled studies in pregnant woman.

DEXATROL Eye Drops should be used during pregnancy only if the potential benefit to the mother justifies the potential risk to the embryo or fetus.

Infants born of mothers who have received substantial doses of corticosteroids during pregnancy should be observed carefully for signs of hypoadrenalism.

It is not known whether topical administration of corticosteroids could result in sufficient systemic absorption to produce detectable quantities in human milk.

Because many drugs are excreted in human milk, caution should be exercised when **DEXATROL Eye Drops** is administered to an nursing woman.

Side Effects

Adverse reactions have occurred with corticosteroid/ anti-infective combination drugs which can be attributed to the corticosteroids component, or the combination.

Reactions occurring most ofter from the presence of the anti-infective ingredient are allergic sensitizations.

The reactions due to the corticosteroid component are elevation of intraocular pressure (IOP) with possible development of glaucoma, and infrequent optic nerve damage; posterior subcapsular cataract formation; and delayed wound healing.

Secondary Infection:

The development of secondary infection has occurred after use of combinations containing corticosteroids and antimirobials.

Fungal infections of the cornea are particularly prone to develop coincidentally with long-term applications of corticosteroids.

The possibility of fungal invasion must be considered in any persistent corneal ulceration where corticosteroid treatment has been used.

Secondary bacterial ocular infection following suppression of host responses also occurs.

Symptoms and Treatment of Overdose

Long-term intensive topical use may lead to systemic effects. Oral ingestion of the contents of up to 10 ml is unlikely to lead to any serious adverse effects.

Presentation

Dexatrol Eye drops is available in 5ml LDPE plastic dropper bottle in a folded paper box.

Storage conditions

Store at controlled room temperature below 30°C .

Store at room temperature below 25°C after opening.

Shelf Life

24 months from the date of manufacture if kept as recommended. This product can be used for 28 days after the first opening if kept as recommended.

Registration number

DEXATROL EYE DROPS – MAL14050001AZ

Name and Address of Manufacturer

Egyptian International Pharmaceutical Industries Co. (EIPICO)
Tenth of Ramadhan City, Industrial Area B1, P.O.Box : 149 Tenth, Egypt
Tel: 2-0554 499199
Fax: 2-0554 499306

Product Registration Holder

Averroes Pharmaceuticals Sdn. Bhd.
03-08-01 & 03-09-01, Block 3, Presint Alami,
Worldwide Business Center II,
Persiaran Akuatik, Seksyen 13
40100 Shah Alam, Selangor
Malaysia
Tel: +603 5511 1433
Fax: +603 5511 1431

Date of revision of package insert

5th January 2023

Black C 9.5x16.5 cm 333 8M2 PAM 002 002

EEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEEE

EGYPTIAN INT. PHARMACEUTICAL INDUSTRIES CO. E. I. P. I. CO.
10th OF RAMADAN CITY , INDUSTRIAL AREA B1, P.O. BOX: 149 TENTH, EGYPT