

Direction of reading



Direction of reading



Canesten® V Cream 1% w/w

Effective treatment & soothing relief of external thrush symptoms
Clotrimazole 1% w/w



Read all of this leaflet carefully before you take this medicine.

Keep this leaflet. You may need to read it again. If you have further questions, ask your doctor or pharmacist. If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

What is in this leaflet

1. What Canesten® V Cream 1% w/w is used for
2. How Canesten® V Cream 1% w/w works
3. Before you use Canesten® V Cream 1% w/w
4. How to use Canesten® V Cream 1% w/w
5. While you are using it
6. Side effects
7. Storage and disposal of Canesten® V Cream 1% w/w
8. Product description
9. Manufacturer & Product Registration Holder
10. Date of revision date

What Canesten® V Cream 1% w/w is used for

Canesten® V Cream 1% w/w is used to relieve irritation of the vulva (external thrush) or irritation of the end of the penis associated with thrush.

How Canesten® V Cream 1% w/w works

The active substance in Canesten® V Cream 1% w/w is clotrimazole. Clotrimazole belongs to a group of medicines called imidazoles. The primary mechanism of action of clotrimazole is against the division and growing of fungi. Clotrimazole alters the permeability of the fungal cell wall and inhibits the activity of enzymes within the cell. It specifically inhibits the biosynthesis of ergosterol required for cell membrane production.

Before you use Canesten® V Cream 1% w/w

When you must not use it

If you are allergic (hypersensitive) to clotrimazole or any of the other ingredients, including cetostearyl alcohol.

Before you start to use it

If you are pregnant, breast-feeding or trying for a baby, tell your doctor or midwife before using Canesten® V Cream 1% w/w. If you have informed your doctor or midwife already, follow his/her instructions carefully.

As a precautionary measure, it is preferable to avoid the use of clotrimazole during the first trimester of pregnancy. Breast-feeding should be discontinued during treatment with clotrimazole. As with other creams, Canesten® V Cream 1% w/w may reduce the effectiveness of rubber contraceptives, such as condoms or diaphragms. Consequently, if you are using this cream on the vulva or penis, you should use alternative precautions for at least five days after using this product.

Taking other medicines

Not known.

How to use Canesten® V Cream 1% w/w

How much to use

Before use, pierce the tube seal by inverting the cap over the end of the tube and press. Canesten® V Cream 1% w/w should be applied thinly and evenly to the affected areas two or three times daily and rubbed in gently. A strip of cream (1/2 cm long) is enough to treat an area of about the size of the hand. The symptoms of skin infection, such as itching or soreness, should improve within a few days of treatment although signs such as redness and scaling may take longer to disappear.

If symptoms persist, consult your doctor. Patients should notify their physician if there is no improvement after 4 weeks of treatment.

When to use it

It can be applied anytime of the day.

How long to use it

1-2 weeks.

If you forget to use it

Apply the cream as soon as possible and then continue the rest of your treatment as usual.

If you use too much (overdose)

No reports are available on cases of overdose with Canesten® V Cream 1% w/w. There is no specific antidote. Topical clotrimazole is intended for cutaneous use only. In the case of cutaneous overdose patients should be managed by symptomatic care; patients should wash off the product with water and cleanser, discontinue use, and see their physician so appropriate therapy can be implemented as needed. In the case of ingestion of an overdose, patients should see their physician and be managed by symptomatic and supportive care.

While you are using it

Things you must do

You can help the treatment to work if you follow these simple self-help tips:

- Keep the affected skin areas clean.
- Pay particular attention to drying the skin, but avoid excessive rubbing.
- Do not share towels, bath mats, etc. with other people as you could spread the infection to them.
- Always wash your hands after treating the infection to prevent it from spreading.

Things you must not do

Canesten® V Cream 1% w/w is for external use only:

Do not put the cream in your mouth or swallow it. Although the infected area will itch, try not to scratch. Scratching will damage the surface of the skin and cause the infection to spread further.

Things to be careful of

This product contains cetostearyl alcohol which may cause local skin irritation (e.g. rash, itching or redness).

Side effects

Like all medicines, Canesten® V Cream 1% w/w can cause side effects, although not everybody gets them. As with all medicines, some people may be allergic to the cream. If you are allergic, a reaction will occur soon after you start using it. If you experience an allergic reaction, stop using Canesten® V Cream 1% w/w.

Signs of an allergic reaction may include:

- Rash.
- Swallowing or breathing problems.
- Swelling of your lips, face, throat or tongue.
- Weakness, feeling dizzy or faint.
- Nausea.
- Discomfort/pain.

After you apply the cream you might experience:

- Itching, rash, blisters, burning, discomfort, swelling, irritation or peeling of skin.

If you experience any of the above effects or react badly to the cream in any other way not listed in this leaflet, tell your doctor or pharmacist immediately.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website nptra.gov.my [Consumers -> Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of Canesten® V Cream 1% w/w

To be stored below 30°C.

Keep out of reach of children.

Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product description

What it looks like

Canesten® V cream 1% w/w is a white smooth cream.

Ingredients

- Active ingredient : clotrimazole 1% w/w.
- Inactive ingredients : sorbitan stearate, polysorbate 60, cetyl palmitate, cetostearyl alcohol, octyldodecanol, benzyl alcohol & purified water.

MAL14085053XCRZ

Manufacturer

Encube Ethicals Private Limited
C- 1, Madkaim Industrial Estate,
Madkaim, Ponda, Goa, 403404 India
For Bayer Consumer Care AG
Basel, Switzerland

Product Registration Holder

Bayer Co. (Malaysia) Sdn Bhd,
25-03 & 25-04,
Level 25, IMAZIUM
No 8, Jalan SS21/37,
Damansara Uptown
47400 Petaling Jaya
Selangor

Date of revision: 01 June 2022

Bayer
000000