

DEXALONE SOLUTION FOR INJECTION 2mg

Composition:

Dexamethasone (as sodium phosphate) 1.52 mg
(i.e. 2 mg of dexamethasone sodium phosphate)
Phenol 5.00 mg
Excipient qs 1 ml

Product Description: A clear colourless solution, salt taste.

Pharmacodynamics: Dexamethasone is a potent glucocorticoid with a low mineralocorticoid action. Dexamethasone is ten to twenty times as active as prednisolone at equimolar doses. The corticosteroids may depress the immune response. It inhibits the capillary dilatation, leucocytes migration and phagocytosis. The glucocorticoids act on the metabolism by increasing neoglucogenesis. Dexamethasone induces parturition in ruminants if foetus is alive.

Pharmacokinetics: After administration via intramuscular administration, sodium phosphate dexamethasone is rapidly absorbed, and hydrolysed as dexamethasone (base), giving a quick action response and short in time. T_{max} in cattle, goats, horses and pigs is reached within 30 minutes of intramuscular administration. $T_{1/2}$ varies according to the species between 5 and 20 hours. The bioavailability after intramuscular administration is about 100%.

Indication:

Horse, cattle, goats and pigs: Treatment against inflammatory states, allergy or shock.

Cattle, goats: Primary ketosis treatment (acetone, pregnancy toxemia) and induction of parturition.

Horse: Treatment of osteoarticular inflammatory states.

Routes of administration: Intravenous, intramuscular, subcutaneous, intra-articular and peri-articular injection.

Dosage:

Species	Dosage (in mg of dexamethasone/kg)
Horse, cattle, goats, pigs	0.015 to 0.06 mg/kg (ie. 1 to 4 ml/100kg)

The product may also be administered via intra-articular or peri-articular injection. The recommended dosage is 0.125 to 5 ml per animal according to the weight.

The dosage may be repeated if necessary at 24-48 hours intervals.

Induce parturition:

Cow: 20 mg of dexamethasone (13 ml)

Goat: 12 to 16 mg of dexamethasone (8 to 10 ml)

Contraindications:

Do not use in animals suffering from diabetes mellitus, hyperadrenocorticism (Cushing syndrome), renal failures and cardiac failures or duodenal ulcers.

Do not use in animals with infectious disease, unless an adapted anti-infectious therapy is given at the same time.

Warnings and Precautions:

Special warnings for each target species

In horses, use dexamethasone only in beginning of laminitis.

Special precautions for use in animals

Administration of corticosteroids at end of pregnancy may induce, in ruminants, a premature calving or abortion.

Except when acetone indications and parturition induction, the corticosteroids function is to improve the state more than curing.

Therefore, it is recommended to diagnose and to treat the underlying disease. When treating a shock, intravenous liquids should be used to maintain circulation; an acid-base balance should be done.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

In case of accidental self-injection, seek for medical advice and show the package leaflet to the physician.

People with known hypersensitivity to the active substance or the excipient should avoid contact with the medicinal product.

The product should not be administered by pregnant women.

Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Interactions with other medicaments

Corticosteroids may reduce immune response to vaccines. Do not use dexamethasone during vaccination.

The use with non-steroid anti-inflammatory drugs may increase the risk of gastro-intestinal ulcers.

Statement on usage during pregnancy and lactation:

Studies in laboratory animals have demonstrated embryotoxic effects.

The use of corticosteroids in pregnant females is not recommended, except in ruminants to induce parturition during the last third of pregnancy.

Adverse Effects:

In case of long treatment, corticosteroids such as dexamethasone may induce iatrogenic hypercorticism, polyuripolydipsia (PUPD), immunodepression, bulimia and redistribution of lipidic stores in the body.

The use of corticosteroids in lactating cows and goats may induce a temporary decrease in milk production.

Induction of parturition with corticosteroids may be associated with a low viability of the progeny and an increase of the retained placenta.

Overdose: High levels of corticosteroids may lead to drowsiness and lethargy in horse.

Incompatibilities: None known

Storage Conditions: Do not store above 25°C.

Withdrawal Period:

Meat and offal: 6 days

Milk: 3 days

Packing: 50ml and 100ml vial.

FOR VETERINARY USE ONLY

Manufactured by:



DOPHARMA FRANCE, 23 rue du Prieuré, Saint Herblon, 44150 Vair sur Loire, France

Registration holder and repacker:



Rhona Ma Malaysia Sdn Bhd (200001023279)

Lot 18A & 18B, Jalan 241, Seksyen 51A, 46100 Petaling Jaya, Selangor Malaysia

Tel: 03-7873 7355 Fax: 03-2770 0119

E-mail: customercareline@rhonema.com

Date of revision of: January 2020
SLDEXAL3-0503/XXX

LABEL CODE:	VERSION	COLORS
PRODUCT: DEXALONE SOLUTION FOR INJECTION 2MG PACK SIZE: 50ML and 100 ML COUNTRY: MY DIMENSIONS: cm = 10 x 14	VERSION: 3 MMM/YY: XXX	BLACK ! Please use the official PANTONE® (P) matching system for accurate colour representation.

Marketing approval:	Regulatory approval:	GMP approval:
COMPLETE NAME: DATE: SIGNATURE:	COMPLETE NAME: DATE: SIGNATURE:	COMPLETE NAME: DATE: SIGNATURE: