

NUTRIMOM TABLET

VINAT00-0

Description:

Oblong, pink film-coated tablet, deep convex faces with 'HD' embossed and scored on the same face.

Composition:

Each tablet contains:

Vitamin A	1332.00 IU	Vitamin K1	90.00 mcg
Thiamine Mononitrate	1.50 mg	Copper	1.00 mg
(providing 1.40 mg of Vitamin B1)		Folic Acid	0.40 mg
Riboflavin	1.60 mg	Iodine	0.15 mg
Nicotinamide	18.00 mg	Iron	27.00 mg
Pyridoxine Hydrochloride	2.00 mg	Magnesium	100.00 mg
(providing 1.65 mg of Vitamin B6)		Manganese	1.80 mg
Cyanocobalamin	2.00 mcg	Pantothenic Acid	6.00 mg
Vitamin C	65.00 mg	Selenium	40.00 mcg
Vitamin D3	200.00 IU	Zinc	15.00 mg
Vitamin E	15.00 IU		

Indications:

Vitamin and mineral supplements for pregnant women.

Contraindications:

- Avoid in patients with hypervitaminosis A or D, renal insufficiency, and hypersensitivity to any of the ingredients of the preparation.
- Not recommended for patients with hyperthyroidism, hypercalcemia, hypercalcuria, calcium based kidney stones, or megaloblastic anaemia.
- Should not be taken by patients suffering from iron overload or disturbances in utilization of iron.

Precautions:

- Daily recommended dosage should not be exceeded. Certain vitamins, minerals and trace elements can cause side effects in high doses, particularly during prolonged administration.
- Overdose of vitamin D3 in pregnancy must be prevented. Sustained hypercalcaemia may lead to physical or mental retardation, congenital heart and eye diseases in the child.
- Consult a doctor / pharmacist if you are on anticoagulant/blood thinner products.

Main Side/Adverse Effects:

Diarrhoea, itching, headache, flushing or redness of skin, nausea or vomiting, gastrointestinal discomfort and yellow discoloration of urine may occur occasionally. Consult your doctor if you experience any adverse effects.

Drug Interactions:

- Effects of levodopa may be reduced by vitamin B6.
- Co-administration of probenecid may decrease gastrointestinal absorption of vitamin B2.
- Aspirin may increase urinary secretion of ascorbic acid; however, clinical significance is unclear.
- Vitamin E may facilitate absorption, hepatic storage and utilization, and reduce toxicity of vitamin A; excessive doses may deplete vitamin A stores.
- Concomitant administration of iron and tetracycline antibiotics diminishes each other's absorption. If concomitant intake should be nevertheless required, a lapse of 2 hours should be considered.
- Avoid concurrent use of medications containing cholestyramine, colestipol, digitalis, magnesium-containing antacids, mineral oil, phenytoin, high doses of phosphorus-containing preparations unless otherwise directed by physician.

Overdose:

Clinical features: Headache, nausea, vomiting, abdominal pain, anorexia, diarrhoea, drowsiness, flushing, hypertension, acute renal failure, papilloedema, generalized peeling of skin, pruritus, polyuria, polydipsia, hyperthyroidism, fatigue, muscle weakness.

Treatment for overdosage:

Emesis or gastric lavage if appropriate. Symptomatic and supportive measures.

Dosage and Administration:

Adults: 1 tablet once daily after meal.

Note: The information given here is limited. For further information, consult your doctor or pharmacist.

Storage: Store below 30°C. Protect from light and moisture.

Presentation / Packing: Bottle of 60's.

Product Registration Holder:

HOVID Bhd.,
121, Jalan Tunku Abdul Rahman, 30010 Ipoh, Perak, Malaysia

Manufactured by:

HOVID Bhd.,
Lot 56442, 7½ Miles, Jalan Ipoh / Chemor, 31200 Chemor, Perak, Malaysia.

