

**Gascovid Liquid**

VIGASxx-x

**Description**

White to off white opaque suspension with peppermint flavour.

**Composition**

Each 10ml contains Sodium Alginate 500mg, Sodium Bicarbonate 267mg and Calcium Carbonate 160mg.

**Actions and Pharmacology**

Pharmacotherapeutic classification: A02BX Other drugs for peptic ulcer and gastro-oesophageal reflux disease (GORD)

On ingestion Gascovid Liquid react rapidly with gastric acid to form a raft of alginic acid gel having a near neutral pH and which floats on the stomach contents, quickly and effectively impeding gastro-oesophageal reflux, for up to 4 hours. In severe cases the raft itself may be refluxed into the oesophagus, in preference to the stomach contents, and exert a demulcent effect.

**Pharmacokinetics**

The mode of action of Gascovid Liquid is physical and does not depend on absorption into the systemic circulation.

**Indications**

Treatment of symptoms of gastro-oesophageal reflux such as acid regurgitation, heartburn, indigestion due to reflux of stomach contents.

**Contraindications**

This medicinal product is contraindicated in patients with known or suspected hypersensitivity to the active substances or to any of the excipients.

**Pregnancy and Lactation**

Open controlled studies in 281 pregnant women did not demonstrate any significant adverse effects of Gascovid Liquid on the course of pregnancy or on the health of the foetus or new-born child. Based on this Gascovid Liquid may be used during pregnancy and lactation.

**Precautions**

This product contains methyl hydroxybenzoate and propyl hydroxybenzoate which may cause allergic reactions possibly delayed.

This product contains sodium. This should be taken into account when a highly restricted salt diet is recommended, e.g. in some cases of congestive cardiac failure and renal impairment.

This product contains calcium carbonate. Care needs to be taken in treating patients with hypercalcaemia, nephrocalcinosis and recurrent calcium containing renal calculi.

There is a possibility of reduced efficacy in patients with very low levels of gastric acid.

If symptoms do not improve after seven days, the clinical situation should be reviewed.

Treatment of children younger than 12 years of age is not generally recommended, except on medical advice.

**Interaction with Other Medicaments and Other Forms of Interaction**

Due to the presence of calcium carbonate which act as an antacid, a time-interval of 2 hours should be considered between Gascovid Liquid intake and the administration of other medicinal products, especially H<sub>2</sub>-antihistaminics, tetracyclines, digoxine, fluoroquinolone, iron salt, ketoconazole, neuroleptics, thyroxine, penicillamine, beta-blockers (atenolol, metoprolol, propranolol), glucocorticoid, chloroquine, and diphosphonates.

**Main Side/Adverse Effects**

Very rarely (<1/10,000) patients may develop allergic manifestations such as urticaria or bronchospasm, anaphylactic or anaphylactoid reactions.

**Overdose**

In the event of overdosage symptomatic treatment should be given. The patient may notice abdominal distension.

**Dosage and Administration**

For oral use.

Shake well before use.

Adult & children > 12 years old: 10 – 20ml after meals and at bed time.

Children 6 – 12 years old: 5 – 10ml after meals and at bed time.

Children under 6 years old: Should be given only on medical advice.

Note: The information given here is limited. For further information consult your doctor or pharmacists.

Storage : Store below 30°C. Do not refrigerate or freeze.  
Presentation/ packing : Bottle of 150ml, 200ml (*Not all presentations may be available locally*)

Product Registration Holder (Malaysia) : Hovid Bhd., 121, Jalan Tunku Abdul Rahman (Jalan Kuala Kangsar),  
/ Manufactured by 30010 Ipoh, Perak, Malaysia.

Information date : January 2019