

**IDAMAN PHARMA ASCORBIC ACID 100MG TABLET****DESCRIPTION**

Clean, dry, uniform, round, biconvex, even edged tablet.

Colour : Peach.

Taste : Orange Flavour.

Each tablet contains : Ascorbic Acid 100mg

**INDICATION**

Used as health supplement.

**RECOMMENDED DOSE**

**Health supplement dose:** Adult: 1 tablet once a day after meal.

**ROUTE OF ADMINISTRATION**

Oral

**CONTRAINDICATIONS**

Unknown.

**PHARMACOLOGY**

1. Absorption: Readily absorbed after oral administration.
2. Protein binding: Low
3. Storage: Present in plasma cells; higher concentration in glandular tissue
4. Metabolism: Hepatic
5. Excretion: Renal, very little as unchanged drug and metabolite except with high doses; urinary excretion increases with plasma concentration greater than 1.4mg/100ml.  
In dialysis – removal by hemodialysis.

**WARNINGS AND PRECAUTIONS**

**Pregnancy:** Ingestion of large quantities of ascorbic acid during pregnancy may result in increased requirements and scurvy in neonate.

This product contains aspartame and is unsuitable for phenylketonurics.

Importance of not taking more medication than the amount recommended; extra is excreted in urine; risk of stones in urinary tract; very high doses may interfere with glucose determinations; megadoses not recommended without physician's advice.

Missed dose: No cause for concern because of length of time necessary for depletion; remembering to take as directed.

**DRUG INTERACTIONS**

Combination containing any of the following medications may interact:

**a) Anticoagulants**

Concurrent use with ascorbic acid may interfere with the effect of oral anticoagulant.

**b) Barbiturate or Primidone**

Concurrent use may increase urinary excretion of ascorbic acid; requirement of ascorbic acid may be increased in patients receiving this medication.

**c) Cellulose sodium phosphate**

Concurrent use may result in metabolism of ascorbic acid to oxalate.

**d) Disulfiram**

Concurrent use with ascorbic, especially in chronic use or high doses, may interfere with the disulfiram-alcohol interaction.

**e) Mexiletine**

Acidification of urine resulting from use of large doses of ascorbic acid may accelerate renal excretion of mexiletine.

**f) Salicylates**

Concurrent use may increase urinary excretion of ascorbic acid; clinical significant is unclear, though some clinicians recommend vitamin supplements in patients receiving prolonged high-dose aspirin therapy.

**SIDE EFFECTS / ADVERSE REACTIONS**

Withdrawal scurvy may occur after prolonged administration of 2-3 gram/day. Occasionally, prolonged use of high doses of ascorbic acid may cause precipitation of urate, cyctine or oxalate stones in the urinary tract.

**SYMPTOMS AND TREATMENT FOR OVERDOSE**

Symptoms: Diarrhea, flushing or redness of skin, headache, nausea or vomiting, increased urination and stomach cramps

Treatment: Supportive and symptomatic.

**PACK SIZES**

Bottles of 60 Tablets and 100 Tablets

**STORAGE CONDITION**

Keep container tightly closed in a dry place, below 30°C, protect from light.

**SHELF LIFE**

Product should not be used beyond the expiry date imprinted on the product packaging.

**REGISTRATION NUMBER**

MAL16015063NZ

For further information, please consult your doctor or your pharmacist.

Revision number: 01

Revision date: 02 Apr 2026

**Lt-177.06 PRODUCT REGISTRATION HOLDER AND MANUFACTURER**

**INSPIRAPHARMA SDN. BHD.** (200401023395)

LOT 24 & 25, JALAN 8,

KAWASAN PERUSAHAAN BAKAR ARANG,

08000 SUNGAI PETANI,

KEDAH DARUL AMAN, MALAYSIA.