

RAMTACE TABLET

Ingredient(s):
Ramtace Tablet 10mg
Each tablet contains:
Ramipril 10mg

Ramtace Tablet 5mg
Each tablet contains:
Ramipril 5mg

Pharmacology (Summary of Pharmacodynamic and Pharmacokinetic):
Pharmacodynamic:
Mechanism of action

Ramiprilat, the active metabolite of the prodrug ramipril, inhibits the enzyme dipeptidylcarboxypeptidase I (synonyms: angiotensin-converting enzyme; kininase II). In plasma and tissue this enzyme catalyses the conversion of angiotensin I to the active vasoconstrictor substance angiotensin II, as well as the breakdown of the active vasodilator bradykinin. Reduced angiotensin II formation and inhibition of bradykinin breakdown lead to vasodilation. Since angiotensin II also stimulates the release of aldosterone, ramiprilat causes a reduction in aldosterone secretion. The average response to ACE inhibitor monotherapy was lower in black (Afro-Caribbean) hypertensive patients (usually a low-renin hypertensive population) than in non-black patients.

Pharmacodynamic effects:
Antihypertensive properties:

Administration of ramipril causes a marked reduction in peripheral arterial resistance. Generally, there are no renal changes in renal plasma flow and glomerular filtration rate. Administration of ramipril to patients with hypertension leads to a reduction in supine and standing blood pressure without a compensatory rise in heart rate. In most patients the onset of the antihypertensive effect of a single dose becomes apparent 1 to 2 hours after oral administration. The peak effect of a single dose is usually reached 3 to 6 hours after oral administration. The antihypertensive effect of a single dose usually lasts for 24 hours. The maximum antihypertensive effect of continued treatment with ramipril is generally apparent after 3 to 4 weeks. It has been shown that the antihypertensive effect is sustained under long term therapy lasting 2 years. Abrupt discontinuation of ramipril does not produce a rapid and excessive rebound increase in blood pressure.

Heart failure:
In addition to conventional therapy with diuretics and optional cardiac glycosides, ramipril has been shown to be effective in patients with functional classes II-IV of the New-York Heart Association. The drug had beneficial effects on cardiac haemodynamics (decreased left and right ventricular filling pressures, reduced total peripheral vascular resistance, increased cardiac output and improved cardiac index). It also reduced neuroendocrine activation.

Pharmacokinetic:
Absorption
Following oral administration, ramipril is rapidly absorbed from the gastrointestinal tract; peak plasma concentrations of ramipril are reached within 1 hour. Based on urinary recovery, the extent of absorption is at least 56% and is not significantly influenced by the presence of food in the gastrointestinal tract. The bioavailability of the active metabolite, ramiprilat, after oral administration of 2.5mg and 5mg ramipril is 45%. Peak plasma concentrations of ramiprilat, the sole active metabolite of ramipril are reached 2-4 hours after ramipril intake. Steady state plasma concentrations of ramiprilat after once daily dosing with the usual doses of ramipril are reached by about the 4th day of treatment.

Distribution:
The serum protein binding of ramipril is about 73% and that of ramiprilat about 56%.

Metabolism:
Ramipril is almost completely metabolised to ramiprilat and to the diketopiperazine ester, the diketopiperazine acid and the glucuronides of ramipril and ramiprilat.

Elimination:
Excretion of the metabolites is primarily renal. Plasma concentrations of ramiprilat decline in a polyphasic manner. Because of its potent, saturable binding to ACE and slow dissociation from the enzyme, ramiprilat shows a prolonged terminal elimination phase at very low plasma concentrations.

After multiple once-daily doses of ramipril, the effective half-life of ramiprilat concentrations was 15-17 hours for the 5-10mg doses and longer for the lower 1.25-2.5mg doses. This difference is related to the saturable capacity of the enzyme to bind ramiprilat. A single oral dose of ramipril produced an undetectable level of ramipril and its metabolite in breast milk. However, the effect of multiple doses is not known.

Patients with Renal Impairment: Renal excretion of ramiprilat is reduced in patients with impaired renal function and renal ramiprilat clearance is proportionally related to creatinine clearance. This result in elevated plasma concentrations of ramiprilat, which decrease more slowly than in subjects with normal renal function.

Patients with Hepatic Impairment: In patients with impaired liver function, the metabolism of ramipril to ramiprilat was delayed, due to diminished activity of hepatic esterases and plasma ramipril levels in these patients were increased. Peak concentrations of ramiprilat in these patients, however, are not different from those with normal hepatic function.

- Indication(s):**
1. Hypertension.
 2. Congestive heart failure.
 3. Treatment of patients who within the first few days after an acute myocardial infarction have demonstrated clinical signs of congestive heart failure.
 4. For reducing the risk of myocardial infarction, stroke, cardiovascular death or the need for revascularization procedures in patients ≥ 55 years who have clinical evidence of coronary artery disease, stroke, or peripheral vascular disease.
 5. For reducing the risk of myocardial infarction, stroke, cardiovascular death or revascularization procedures in diabetic patients ≥55 years with ≥1 of the following risk factors:
Systolic blood pressure >160 mm Hg or diastolic blood pressure > 90mm Hg (or on antihypertensive treatment); total cholesterol >5.2mmol/L; HDL cholesterol <0.9mmol/L; current smoker; known microalbuminuria; any evidence of previous vascular disease.
 6. Prevention of progressive renal failure in patients with persistent proteinuria.

Dosage and Administration(s):
The dosage is based on the desired effect and on how the patient tolerates this product. Therapy with this product is usually long term therapy; the doctor determines the duration of treatment individually for each patient.

Hypertension:
The recommended initial dose: 2.5mg once daily. Depending on the response, the dose may be increased. Any increase should be implemented by doubling the dose at intervals of 2-3 weeks.
Usual maintenance dose: 2.5-5mg daily.
Maximum Dose: 10mg daily.

In impaired renal function, i.e. a creatinine clearance between 50 and 20ml/min per 1.73m² body surface area, the initial dose is generally 1.25mg and the maximum daily dose is 5mg.
When creatinine clearance cannot be measured, it can be calculated based on the serum creatinine level using the following formulation (Cockcroft's equation):

Men:

$$\text{Creatinine Clearance (ml/min)} = \frac{\text{Body Weight (kg)} \times (140 - \text{age in years})}{72 \times \text{serum creatinine (mg/dL)}}$$

Women: Multiply the product of the above equation by 0.85

In patients with incompletely corrected fluid or salt deficiency, those with severe hypertension, as well as in those for whom a hypotensive reaction would constitute a particular risk (eg. patients with haemodynamically relevant stenoses of the coronary arteries or of the blood vessels supplying the brain) and in the elderly, a reduced initial dose of 1.25mg daily must be considered.

In patients pre-treated with diuretic, consideration must be given to discontinuing the diuretic for at least 2-3 days or depend on the duration of action of the diuretic, longer before starting treatment with this product, or at least to reducing the diuretic dose. The doctor will decide in each individual case whether such discontinuation or dose reduction is possible and how long it should last. The initial dose in such patients is generally 1.25mg.

In impaired liver function, response to treatment may be either increased or reduced. Therefore, treatment must be initiated only under close medical supervision. Maximum daily dose: 2.5mg.

Congestive Heart Failure:
Recommended initial dose: 1.25mg once daily. Depending on the response, the dose may be increased. Any increase should be implemented by doubling the dose at intervals of 1-2 weeks.
Maximum daily dose: 10mg.
The required daily dose, if equaling or exceeding 2.5mg, may be taken as a single dose or in 2 separate doses.
In impaired liver or renal function and in patients pre-treated with a diuretic, dosage recommendations for this product are identical to those mentioned previously in the treatment of hypertension. The recommendations given there in conjunction with diuretic pre-treatment also apply.

Myocardial Infarction:
Recommended initial dose: 5mg daily, divided into 2 single doses of 2.5mg each, one in the morning and one in the evening. If this dose is not well tolerated, 1.25mg should be taken twice daily over 2 days. In either event, depending on the response, the dose may then be increased. Any increase should be implemented by doubling the dose at intervals of 1-3 days. As treatment progresses, the total daily dose may be taken as a single dose.
Maximum daily dose: 10mg

Sufficient experience is still lacking in the treatment of patients with severe (NYHA IV) heart failure immediately after myocardial infarction. Treatment, if nevertheless given, should be started with 1.25mg once daily, and increased only with particular caution.

In patients with impaired liver or renal function, with incompletely corrected fluid or salt deficiency, or with severe hypertension, and in those for whom a hypotensive reaction would constitute a particular risk (eg. patients with haemodynamically relevant stenoses of the coronary arteries or of the blood vessels supplying the brain), as well as in those pre-treated with a diuretic and in the elderly, the recommendations are identical to those given previously in the treatment of hypertension.

Prevention of progressive renal failure in patients with persistent proteinuria:
The recommended initial dose is 1.25mg once daily. This should be doubled at intervals of 2-3 weeks, depending on how the drug is tolerated. There are no efficacy data regarding doses above 5mg/day in patients with nephropathy. In hypertensive patients, a target diastolic blood pressure of <90mmHg should be pursued. In patients pre-treated with a diuretic, consideration must be given to discontinuing the diuretic for at least 2-3 days or longer (depending on duration of action) or at least consideration should be given to reducing the dose, before initiating this product.

Patients at increased cardiovascular risk:
Recommended initial dose: 2.5mg once daily. Depending on the tolerability, the dose should be doubled after 1 week of treatment and after 3 weeks, should be increased to 10mg.
Usual maintenance dose: 10mg daily.

Patients with impaired renal function:
In hypertensive patients with creatinine clearance levels of ≥50ml/min (serum creatinine < 1.5mg/dL) a dosage adjustment is not required.

For patients with creatinine clearance levels between 20 and 50ml/min (serum creatinine between 1.5 and 3mg/dL), the recommended initial dose is 1.25mg once daily. This should be doubled at intervals of 2-3 weeks, depending on how the drug is tolerated.

Particular care should be exercise in patients with impaired renal function who are to be treated for heart failure post-myocardial infarction as such patients may be susceptible to hypotension. Patients with impaired renal function treated for heart failure post MI have not been studied systemically.

Patients with impaired liver function:
In patients with impaired liver function, the metabolism of ramipril and therefore, the formation of the bioactive metabolite ramiprilat, is delayed due to diminished activity of the esterases in the liver, resulting in elevated plasma ramipril levels. Treatment with this product should therefore be initiated under close medical supervision and should not exceed 2.5mg daily.

Elderly:
Recommended starting dose: 1.25mg once daily, which can then be increased according to the individual patient's blood pressure response.

Paediatric population:
This product is not recommended for use in children and adolescents below 18 years of age due to insufficient data on safety and efficacy.

Mode of Administration:
To be taken orally. It is recommended that this product is taken each day at the same time of the day.
In case of angioedema, this product must be discontinued. Emergency therapy should be instituted promptly. Patient should be kept under observation for at least 12 to 24 hours and discharged after complete resolution of the symptoms. Intestinal angioedema has been reported in patients treated with ACE inhibitors. These patients presented with abdominal pain (with or without nausea or vomiting).

- Contraindications(s):**
1. Hypersensitivity to ramipril and any other ACE (Angiotensin converting enzyme) inhibitors.
 2. History of angioedema (hereditary, idiopathic or due to previous angioedema with ACE inhibitors or AIIRAs)
 3. Extracorporeal treatments leading to contact of blood with negatively charged surfaces.
 4. Significant bilateral renal artery stenosis or renal artery stenosis in a single functioning kidney
 5. 2nd and 3rd trimester of pregnancy.
 6. Ramipril must not be used in patients with hypotensive or haemodynamically unstable states.

Precaution(s) / Warning(s):

ACE INHIBITORS

WARNING:
INCREASED RISK OF BIRTH DEFECTS, FOETAL AND NEONATAL MORBIDITY
AND DEATH WHEN USED THROUGHOUT PREGNANCY

Special Populations
Pregnancy: ACE inhibitors such as ramipril, or Angiotensin II Receptor Antagonists (AIIRAs) should not be initiated during pregnancy. Unless continued ACE inhibitor/ AIIRAs therapy is considered essential, patients planning pregnancy should be changed to alternative anti-hypertensive treatments which have an established safety profile for use in pregnancy. When pregnancy is diagnosed with ACE inhibitor/ AIIRAs, treatment should be stopped immediately, and, if appropriate, alternative therapy should be started.

- **Patients at particular risk of hypotension**
 - **Patients with strongly activated renin-angiotensin-aldosterone system**
- Patients with the strongly activated renin-angiotensin-aldosterone system are at risk of an acute pronounced fall in blood pressure and deterioration of renal function due to ACE inhibition, especially when an ACE inhibitor or a concomitant diuretic is given for the first time or at first dose increase. Significant activation of renin-angiotensin-aldosterone system is to be anticipated and medical supervision including blood pressure monitoring is necessary, for example in:
- patients with severe hypertension
 - patients with decompensated congestive heart failure
 - patients with haemodynamically relevant left ventricular inflow or outflow impediment (e.g. stenosis of the aortic or mitral valve)
 - patients with unilateral renal artery stenosis with a second functional kidney
 - patients in whom fluid or salt depletion exists or may develop (including patients with diuretics)
 - patients with liver cirrhosis and/or ascites
 - patients undergoing major surgery or during anaesthesia with agents that produce hypotension

Cough
Cough has been reported with the use of ACE inhibitors. Characteristically, the cough is non-productive, persistent and resolves after discontinuation of therapy. ACE inhibitor-induced cough should be considered as part of the differential diagnosis of cough.

Effects on ability to drive and use machines
Some adverse effects (e.g. symptoms of a reduction in blood pressure such as dizziness) may impair the patient's ability to concentrate and react and, therefore, constitute a risk in situations where these abilities are of particular importance (e.g. operating a vehicle or machinery). This can happen especially at the start of treatment, or when changing over from other preparations. At the first dose or subsequent increases in dose it is not advisable to drive or operate machinery for several hours.

Interaction with Other Medicaments:
Contra-indicated combinations
Extracorporeal treatments leading to contact of blood with negatively charged surfaces such as dialysis or haemofiltration with certain high-flux membranes (e.g. polyacrylonitril membranes) and low density lipoprotein apheresis with dextran sulphate due to increased risk of severe anaphylactoid reactions. If such treatment is required, consideration should be given to using a different type of dialysis membrane or a different class of antihypertensive agent.

Precautions for use
Potassium salts, heparin, potassium-retaining diuretics and other plasma potassium increasing active substances (including Angiotensin II antagonists, trimethoprim, tacrolimus, ciclosporin): Hyperkalaemia may occur, therefore close monitoring of serum potassium is required.

Antihypertensive agents (e.g. diuretics) and other substances that may decrease blood pressure (e.g. nitrates, tricyclic antidepressants, anaesthetics, acute alcohol intake, baclofen, alfuzosin, doxazosin, prazosin, tamsulosin, terazosin): Potentiation of the risk of hypotension is to be anticipated.

Angioedema
Angioedema has been reported in patients treated with ACE inhibitors including ramipril. In case of angioedema, this product must be discontinued. Emergency therapy should be instituted promptly. Patient should be kept under observation for at least 12 to 24 hours and discharged after complete resolution of the symptoms. Intestinal angioedema has been reported in patients treated with ACE inhibitors. These patients presented with abdominal pain (with or without nausea or vomiting).

Anaphylactic reactions during desensitization
The likelihood and severity of anaphylactic and anaphylactoid reactions to insect venom and other allergens are increased under ACE inhibition. A temporary discontinuation of this product should be considered prior to desensitization.

Hyperkalaemia
Hyperkalaemia has been observed in some patients treated with ACE inhibitors. Patients at risk for development of hyperkalaemia include those with renal insufficiency, age (>70 years), uncontrolled diabetes mellitus, or those using potassium salts, potassium retaining diuretics and other plasma potassium increasing active substances, or conditions such as dehydration, acute cardiac decompensation, metabolic acidosis. If concomitant use of the above mentioned agents is deemed appropriate, regular monitoring of serum potassium is recommended.

Neutropenia/agranulocytosis
Neutropenia/agranulocytosis, as well as thrombocytopenia and anaemia, have been rarely seen and bone marrow depression has also been reported. It is recommended to monitor the white blood cell count before detection of a possible leucopenia. More frequent monitoring is advised in the initial phase of treatment and in patients with impaired renal function, those with concomitant collagen disease (e.g. lupus erythematosus or scleroderma), and all those treated with other medicinal products that can cause changes in the blood picture.

Ethnic differences
ACE inhibitors cause higher rate of angioedema in black patients than in non black patients. As with other ACE inhibitors, ramipril may be less effective in lowering blood pressure in black people than in non black patients, possibly because of a higher prevalence of hypertension with low renin level in the black hypertensive population.

Cough
Cough has been reported with the use of ACE inhibitors. Characteristically, the cough is non-productive, persistent and resolves after discontinuation of therapy. ACE inhibitor-induced cough should be considered as part of the differential diagnosis of cough.

Effects on ability to drive and use machines
Some adverse effects (e.g. symptoms of a reduction in blood pressure such as dizziness) may impair the patient's ability to concentrate and react and, therefore, constitute a risk in situations where these abilities are of particular importance (e.g. operating a vehicle or machinery). This can happen especially at the start of treatment, or when changing over from other preparations. At the first dose or subsequent increases in dose it is not advisable to drive or operate machinery for several hours.

Interaction with Other Medicaments:
Contra-indicated combinations
Extracorporeal treatments leading to contact of blood with negatively charged surfaces such as dialysis or haemofiltration with certain high-flux membranes (e.g. polyacrylonitril membranes) and low density lipoprotein apheresis with dextran sulphate due to increased risk of severe anaphylactoid reactions. If such treatment is required, consideration should be given to using a different type of dialysis membrane or a different class of antihypertensive agent.

Precautions for use
Potassium salts, heparin, potassium-retaining diuretics and other plasma potassium increasing active substances (including Angiotensin II antagonists, trimethoprim, tacrolimus, ciclosporin): Hyperkalaemia may occur, therefore close monitoring of serum potassium is required.

Antihypertensive agents (e.g. diuretics) and other substances that may decrease blood pressure (e.g. nitrates, tricyclic antidepressants, anaesthetics, acute alcohol intake, baclofen, alfuzosin, doxazosin, prazosin, tamsulosin, terazosin): Potentiation of the risk of hypotension is to be anticipated.

Vasopressor sympathomimetics and other substances (e.g. isoproterenol, dobutamine, dopamine, epinephrine) that may reduce the antihypertensive effect of this product: Blood pressure monitoring is recommended.

Allopurinol, immunosuppressants, corticosteroids, procainamide, cytostatics and other substances that may change the blood cell count: Increased likelihood of haematological reactions.

Lithium salts: Excretion of lithium may be reduced by ACE inhibitors and therefore lithium toxicity may be increased. Lithium level must be monitored.

Antidiabetic agents including insulin: Hypoglycaemic reactions may occur. Blood glucose monitoring is recommended.

Non-steroidal anti-inflammatory drugs and acetylsalicylic acid: Reduction of the antihypertensive effect of this product is to be anticipated. Furthermore, concomitant treatment of ACE inhibitors and NSAIDs may lead to an increased risk of worsening of renal function and to an increase in kalaemia.

Pregnancy and Lactation:

ACE INHIBITORS

USE IN PREGNANCY:
INCREASED RISK OF BIRTH DEFECTS, FOETAL AND NEONATAL MORBIDITY
AND DEATH WHEN USED THROUGHOUT PREGNANCY

This product is not recommended during the first trimester of pregnancy and contraindicated during the second and third trimesters of pregnancy.

Epidemiological evidence regarding the risk of teratogenicity following exposure to ACE inhibitors during the first trimester of pregnancy has not been conclusive; however a small increase in risk cannot be excluded. Unless continued ACE inhibitor therapy is considered essential, patients planning pregnancy should be changed to alternative anti-hypertensive treatments which have an established safety profile for use in pregnancy. When pregnancy is diagnosed, treatment with ACE inhibitors should be stopped immediately, and, if appropriate, alternative therapy should be started.

ACE inhibitor/ Angiotensin II Receptor Antagonist (AIIRA) therapy exposure during the second and third trimesters is known to induce human fetotoxicity (decreased renal function, oligohydramnios, skull ossification retardation) and neonatal toxicity (renal failure, hypotension, hyperkalaemia). Should exposure to ACE inhibitor have occurred from the second trimester of pregnancy, ultrasound check of renal function and skull is recommended. Newborns whose mothers have taken ACE inhibitors should be closely observed for hypotension, oliguria and hyperkalaemia.

Because insufficient information is available regarding the use of ramipril during breastfeeding, ramipril is not recommended and alternative treatments with better established safety profiles during breast-feeding are preferable, especially while nursing a newborn or preterm infant.

Side Effect(s):
The safety profile of ramipril includes persistent dry cough and reactions due to hypotension. Serious adverse reactions include angioedema, hyperkalaemia, renal or hepatic impairment, pancreatitis, severe skin reactions and neutropenia/agranulocytosis.

- **Cardiac disorders:** Myocardial ischaemia including angina pectoris or myocardial infarction, tachycardia, arrhythmia, palpitations, oedema peripheral.
- **Blood and lymphatic system disorders:** Eosinophilia, white blood cell count decreased (including neutropenia or agranulocytosis), red blood cell count decreased, haemoglobin decreased, platelet count decreased, bone marrow failure, pancytopenia, haemolytic anaemia.
- **Nervous system disorders:** Headache, dizziness, vertigo, paraesthesia, ageusia, dysgeusia, tremor, balance disorder, cerebral ischaemia including ischaemic stroke and transient ischaemic attack, psychomotor skills impaired, burning sensation, parosmia.
- **Eye disorders:** Visual disturbance including blurred vision, conjunctivitis.
- **Ear and labyrinth disorders:** Hearing impaired, tinnitus.
- **Respiratory, thoracic and mediastinal disorders:** Non-productive tickling cough, bronchitis, sinusitis, dyspnoea, bronchospasm including asthma aggravated, nasal congestion.

Gastrointestinal disorders: Gastrointestinal inflammation, digestive disturbances, abdominal discomfort, dyspepsia, diarrhoea, nausea, vomiting, pancreatitis (cases of fatal outcome have been very exceptionally reported with ACE inhibitors), pancreatic enzymes increased, small bowel angioedema, abdominal pain upper including gastritis, constipation, dry mouth, glossitis, aphthous stomatitis

Renal and urinary disorders: Renal impairment including renal failure acute, urine output increased, worsening of a pre-existing proteinuria, blood urea increased, blood creatinine increased.

Skin and subcutaneous tissue disorders: Rash in particular maculopapular, angioedema; very exceptionally, the airway obstruction resulting from angioedema may have a fatal outcome; pruritus, hyperhidrosis, exfoliative dermatitis, urticaria, onycholysis, photosensitivity reaction, toxic epidermal necrolysis, Stevens-Johnson syndrome, erythema multiforme, pemphigus, psoriasis aggravated, dermatitis psoriasiform, pemphigoid or lichenoid exanthema or enanthema, alopecia.

Musculoskeletal and connective tissue disorders: Muscle spasms, myalgia, arthralgia.

Metabolism and nutrition disorders: Blood potassium increased, anaemia, decreased appetite, blood sodium decreased.

Vascular disorders: Hypotension, orthostatic blood pressure decreased, syncope, flushing, vascular stenosis, hypoperfusion, vasculitis, Raynaud's phenomenon.

General disorders and administration site conditions: Chest pain, fatigue, pyrexia, asthenia.

Immune system disorders: Anaphylactic or anaphylactoid reactions, antinuclear antibody increased.

Hepatobiliary disorders: Hepatic enzymes and/or bilirubin conjugated increased, jaundice cholestatic, hepatocellular damage, acute hepatic failure, cholestatic or cytolytic hepatitis (fatal outcome has been very exceptional).

Reproductive system and breast disorders: Transient erectile impotence, libido decreased, gynaecomastia.

Psychiatric disorders: Depressed moods, anxiety, nervousness, withdrawal syndrome, sleep disorder including somnolence, confusional state, disturbance in attention.


Symptoms and Treatment for Overdosage, and Antidote(s):
Symptoms associated with overdosage of ACE inhibitors may include excessive peripheral vasodilation (with marked hypotension, shock), bradycardia, electrolyte disturbances and renal failure. The patient should be closely monitored and the treatment should be symptomatic and supportive. Suggested measures include primary detoxification (gastric lavage, administration of adsorbents) and measures to restore haemodynamic stability, including administration of α1-adrenergic agonists or angiotensin II (angiotensinamide) administration. Ramiprilat, the active metabolite of ramipril is poorly removed from the general circulation by haemodialysis.

Shelf-Life:
2 years from the date of manufacture.

Storage Condition(s):
Store at temperature below 30°C. Protect from light and moisture.

Product Description & Packing(s):
Ramtace Tablet 10mg
A brownish-red color tablet, oblong with notched sides at double bisect, one side impressed with a score in the middle, 'R' on the left and '10' on the right. The other side impressed with a score. 

Blister packing of 10's x 3, 10's x 10, 10's x 50, 10's x 100

Ramtace Tablet 5mg:
A brownish-red color oblong tablet, one side impressed with a score in the middle, 'R' on the left and '5' on the right. The other side impressed with a score. 

Blister packing of 10's x 3, 10's x 10, 10's x 50, 10's x 100


Manufacturer and Product Registration Holder :
Y.S.P. INDUSTRIES (M) SDN. BHD. (192891-U)
Lot 5, 5 & 7, Jalan P17, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor Darul Ehsan, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3678