

GDMY Anti-Fungal Cream_Insert

Size: 100 (W) x 160 (H) mm

GD961LR2W

guardian

ANTI-FUNGAL CREAM 1% W/W

Composition: Clotrimazole 1% w/w, Chlorocresol 0,2% w/w.

Description: White colour, smooth application, sticky and phenolic odour cream.

Indications: For treatment of skin fungal infections including ringworm (tinea corporis), athlete's foot (tinea pedis), jock itch (tinea cruris), white spot (tinea versicolor) and candidiasis (topikal).

Recommended dose: General: The affected area should be washed and thoroughly dried. Thin layer of clotrimazole cream is evenly rubbed into the affected area twice daily. Treatment should continue for 2-4 weeks after the disappearance of all signs of infection to ensure mycological cure and prevent relapse.

Adult & paediatric dosing based on type of infection:

- Candidiasis (topikal): Twice daily for up to 4 weeks
- White spot (tinea versicolor): Twice daily for up to 4 weeks
- Tinea corporis: Twice daily for up to 4 weeks
- Tinea cruris: Twice daily for up to 4 weeks
- Tinea pedis: Twice daily for up to 4 weeks

Symptoms & treatment of overdose: Acute overdosage with topical application of clotrimazole is unlikely and would not be expected to lead to a life-threatening situation.

Side effects: Local reactions include irritation, burning sensation, contact dermatitis, blister, edema, erythema, peeling of skin, pruritus, urticaria and stinging of skin.

Warning & precautions: For irritation or sensitivity caused with the use of clotrimazole; treatment should be discontinued and appropriate therapy instituted. The patient should be advised to use the medication for the full treatment time even though the symptoms may have improved.

This medicine should be used in the first trimester of pregnancy after consideration of the benefit / risk relation and by physician decision. It is not known whether clotrimazole is excreted in human being milk. As many drugs are excreted in human being milk, caution should be exercised with clotrimazole used by nursing mothers.

Effects on ability to drive and use machines: It is unlikely to impair a patient's ability to drive or use machinery.

Contraindication: Hypersensitivity to clotrimazole.

Interactions with other medicaments: Not known.

Pharmacodynamic: Clotrimazole is an imidazole derivative with a broad spectrum antimycotic activity.

Mechanism of action:

Clotrimazole acts against fungi by inhibiting ergosterol synthesis. Inhibition of ergosterol synthesis leads to structural and functional impairment of the cytoplasmic membrane.

Pharmacodynamic effects:

Clotrimazole has a broad antimycotic spectrum of action in-vitro and in-vivo, which includes dermatophytes, yeasts, moulds, etc.

Under appropriate test conditions, the minimum inhibitory concentration (MIC) values for these types of fungi are in the region of less than 0,062-4(-8) mg/ml substrate. The mode of action of clotrimazole is primarily fungistatic. In-vitro activity is limited to proliferating fungal elements; fungal spores are slightly sensitive.

In addition to its antimycotic action, clotrimazole also acts on trichomonas vaginalis, gram-positive microorganisms (streptococci/staphylococci), and gram-negative microorganisms (bacteroides/gardnerella vaginalis).

In vitro clotrimazole inhibits the multiplication of corneobacteria and gram-positive cocci-with the exception of enterococci in concentrations of 0,5-10 mg/ml substrate and exerts a trichomonacidal action at 100 mg/ml.

Primary resistant variants of sensitive fungal species are very rare; the development of secondary resistance by sensitive fungi has so far only been observed in very isolated cases under therapeutic conditions.

Pharmacokinetic: Absorption of clotrimazole is less than 0,5% after application to skin/ topical application.

Storage condition: Store below 30°C. Keep out of reach of children.

Shelf life: Please refer to the packaging. Do not use beyond the expiry date. **Packing:** 15g in aluminum tube.

The information contained in this leaflet is limited. For further information, please consult your doctor or pharmacist.

Manufactured under the commission of DFI Brands Limited of 5/F Devon House, Talook Place, Quarry Bay, Hong Kong Distributed By Guardian Health & Beauty: 21 Tampines North Drive 2, #03-01, Giant, Singapore 528765 Tel: (65) 6891 8100	Malaysia Product Registration Holder Guardian Health and Beauty Sdn. Bhd. (201401024993) Level 11, 8 First Avenue, Bandar Utama, 47800, Petaling Jaya, Selangor Darul Ehsan, Malaysia. Tel: (60) 3 2777 7100 Guardian Health and Beauty (B) Sdn. Bhd. (PC0000294) Lot 58865 Rimba Point, Kampong Rimba, Mukim Gadong BE3119, Brunei Darussalam	Manufactured By: Syarikat Wen Ken Drug Sdn Bhd 24, Jalan Lambak, Taman Johor, 81200 Johor Bahru, Johor, Malaysia. MAL17035018XCZ BRU24083458P Date of revision: Mar 2025 GD961LR2W
---	---	--

guardian

KRIM ANTI KULAT 1% W/W

Komposisi: Krim ini mengandungi Clotrimazole 1% w/w, Chlorocresol 0,2% w/w.

Deskripsi: Krim berwarna putih, mudah disapu, melekit dan berbau fenolik.

Indikasi: Untuk rawatan jangkitan kulat pada kulit termasuk kurap (tinea corporis), kaki atlet (tinea pedis), gatal pangkal paha (tinea cruris), panau (tinea versicolor) serta kandidiasis (topikal).

Dos saranan: Umum: Basuh dan keringkan dengan menyeluruh bahagian yang dijangkiti. Sapu lapisan nipis krim clotrimazole dengan sekata kepada bahagian yang dijangkiti dua kali sehari. Rawatan perlu diteruskan 2-4 minggu selepas kehilangan semua tanda-tanda jangkitan untuk memastikan penyembuhan yang sempurna dan mengelakkan jangkitan berulang.

Dos dewasa dan pediatrik bergantung kepada jenis jangkitan:

- Kandidiasis (topikal): Dua kali sehari untuk jangkamasa sehingga 4 minggu
- Panau (tinea versicolor): Dua kali sehari untuk jangkamasa sehingga 4 minggu
- Tinea corporis: Dua kali sehari untuk jangkamasa sehingga 4 minggu
- Tinea cruris: Dua kali sehari untuk jangkamasa sehingga 4 minggu
- Tinea pedis: Dua kali sehari untuk jangkamasa sehingga 4 minggu

Symptom & rawatan overdos: Overdos akut dengan aplikasi topikal clotrimazole jarang berlaku dan tidak akan menyebabkan keadaan yang mengancam nyawa.

Kesan sampingan: Tindak balas setempat termasuk iritasi, sensasi panas, dermatitis kontak, lepuh, edema, eritema, pengelupasan kulit, gatal-gatal, urtikaria dan sengatan kulit.

Amaran & perhatian: Untuk iritasi atau sensitiviti yang diakibatkan oleh penggunaan clotrimazole; rawatan perlu dihentikan dan terapi yang sesuai perlu dimulakan. Pesakit adalah dinasihat untuk menghabiskan ubat untuk seluruh jangkamasa rawatan walaupun simptom mungkin telah bertambah baik. Ubat ini boleh digunakan dalam trimester pertama kehamilan selepas menimbang hubungan manfaat/risiko dan dengan nasihat doktor, Ia tidak diketahui samada clotrimazole dikeluarkan dalam susu ibu. Oleh kerana banyak ubat dikeluarkan dalam susu ibu, perhatian perlu diberikan kepada penggunaan clotrimazole oleh ibu yang menyusukan anak.

Kesan terhadap kebolehan memandu dan mengendalikan mesin: Tiada kemungkinan kesan terhadap kebolehan memandu atau mengendalikan mesin.

Kontraindikasi: Hipersensitiviti kepada clotrimazole.

Kehamilan & laktasi: Timbangkan potensi manfaat rawatan ubat terhadap potensi risiko sebelum menetapkan preskripsi ubat ini semasa kehamilan dan laktasi.

Interaksi dengan ubat lain: Tidak ada interaksi ubat yang diketahui.

Farmakodinamik: Clotrimazole adalah derivatif imidazole dengan aktiviti antimikotik spektrum luas.

Tindakan mekanisme:

Clotrimazole bertindak terhadap fungus dengan merencatkan sintesis ergosterol. Perencatan sintesis ergosterol membawa kepada kemerosotan struktur dan fungsi membrane sitoplasmik.

Kesan farmakodinamik:

Clotrimazole memiliki tindakan antimikotik berspektrum luas in-vitro dan in-vivo yang berkesan terhadap dermatofit, yis, kulat dan sebagainya.

Dalam keadaan ujian yang sesuai, nilai kepekatan perencatan minimum (MPC) untuk jenis-jenis fungus ini adalah dalam lingkungan kurang daripada 0,064-4 (-8) mg/ml substrat. Cara tindakan utama clotrimazole ialah fungistatik. Aktiviti in-vitro adalah terhad kepada elemen pembiakan fungus; spora fungus kurang sensitif kepadanya.

Tambahan kepada tindakan antimikotik, clotrimazole juga bertindak terhadap trichomonas vaginalis, mikroorganisma gram-positif (streptococci/staphylococci), mikroorganisma gram-negatif (bacteroides/gardnerel) avaginalis).

Clotrimazole merencatkan pembiakan corneobacteria dan koki gram-positif kecuali enterococci dalam kepekatan 0,5-1,0 mg/ml substrat dan memberi tindakan trichomonasida pada 100 mg/ml.

Jarang sekali berlaku variasi resisten primer spesies fungus yang sensitif; kemunculan resisten sekunder oleh spesies fungus yang sensitif hanya diperhatikan dalam kes asing dalam keadaan terapeutik.

Farmakokinetik: Penyerapan clotrimazole adalah kurang daripada 0,5% selepas sapu pada kulit/ aplikasi topikal.

Kadaan Penyimpanan: Simpan di bawah 30°C. Jauhkan daripada capaian kanak-kanak.

Tempoh hayat: Sila rujuk kepada bungkusan. Jangan guna selepas tamat tarikh luput. **Saiz bungkusan:** 15g dalam tiub aluminium.

Maklumat dalam risalah ini adalah terhad. Untuk informasi selanjut, sila jumpa doktor atau ahli farmasi.

Manufactured under the commission of DFI Brands Limited of 5/F Devon House, Talook Place, Quarry Bay, Hong Kong Distributed By Guardian Health & Beauty: 21 Tampines North Drive 2, #03-01, Giant, Singapore 528765 Tel: (65) 6891 8100	Malaysia Product Registration Holder Guardian Health and Beauty Sdn. Bhd. (201401024993) Level 11, 8 First Avenue, Bandar Utama, 47800, Petaling Jaya, Selangor Darul Ehsan, Malaysia. Tel: (60) 3 2777 7100 Guardian Health and Beauty (B) Sdn. Bhd. (PC0000294) Lot 58865 Rimba Point, Kampong Rimba, Mukim Gadong BE3119, Brunei Darussalam	Manufactured By: Syarikat Wen Ken Drug Sdn Bhd 24, Jalan Lambak, Taman Johor, 81200 Johor Bahru, Johor, Malaysia. MAL17035018XCZ BRU24083458P Date of revision: Mar 2025 GD961LR2W
---	---	--