

OLANZ Tablets 5mg

OLANZ Tablets 10mg

NAME AND STRENGTH OF ACTIVE SUBSTANCE:

OLANZ Tablets 5mg

Each tablet contains : Olanzapine 5mg

OLANZ Tablets 10mg

Each tablet contains : Olanzapine 10mg

PRODUCT DESCRIPTION:

OLANZ Tablets 5mg

A yellow, scored, convex of diameter 8.5mm round tablet with “MPI” marking.

OLANZ Tablets 10mg

A yellow, scored, convex of diameter 11mm round tablet.

PHARMACODYNAMICS:

Olanzapine is an antipsychotic agent that demonstrates a broad pharmacologic profile across a number of receptor systems. Olanzapine exhibited a range of receptor affinities (K_i ; < 100 nM) for serotonin 5 HT_{2A/2C}, 5 HT₃, 5 HT₆; dopamine D₁, D₂, D₃, D₄, D₅; cholinergic muscarinic receptors m1-m5; α_1 adrenergic; and histamine H₁ receptors. Olanzapine reduced a conditioned avoidance response, a test indicative of antipsychotic activity, at doses below those producing catalepsy, an effect indicative of motor side-effects. Unlike some other antipsychotic agents, olanzapine increases responding in an “anxiolytic” test. Olanzapine produced a higher 5 HT_{2A} than dopamine D₂ receptor occupancy. In addition, olanzapine-responsive patients had lower striatal D₂ occupancy than some other antipsychotic- and risperidone-responsive patients, while being comparable to clozapine-responsive patients. Olanzapine was associated with statistically significantly greater improvements in negative as well as positive symptoms. Olanzapine demonstrated statistically significant superiority over placebo on the primary endpoint of bipolar recurrence. Olanzapine also showed a statistically significant advantage over placebo in terms of preventing either recurrence into mania or recurrence into depression.

PHARMACOKINETICS:

Olanzapine is well absorbed after oral administration, reaching peak plasma concentrations within 5 to 8 hours. The absorption is not affected by food. Absolute oral bioavailability relative to intravenous administration has not been determined.

Olanzapine is metabolized in the liver by conjugative and oxidative pathways. The major circulating metabolite is the 10-N-glucuronide, which does not pass the blood brain barrier. Cytochromes P450-CYP1A2 and P450-CYP2D6 contribute to the formation of the N-desmethyl and 2-hydroxymethyl metabolites, both exhibited significantly less *in vivo* pharmacological activity than olanzapine in animal studies. The predominant pharmacologic activity is from the parent olanzapine.

After oral administration, the mean terminal elimination half-life of olanzapine in healthy subjects varied on the basis of age and gender. The plasma clearance of olanzapine is lower in elderly versus young subjects, in females versus males, and in non-smokers versus smokers. However, the magnitude of the impact of age, gender, or smoking on olanzapine clearance and half-life is small in comparison to the overall variability between individuals.

Adolescents (ages 13 to 17 years)

The pharmacokinetics of olanzapine are similar between adolescents and adults.

The plasma protein binding of olanzapine was about 93% over the concentration range of about 7 to about 1000 ng/mL.

Olanzapine is bound predominantly to albumin and α_1 -acid-glycoprotein.

INDICATION:

Olanzapine is indicated for the treatment of schizophrenia.

Olanzapine is effective in maintaining the clinical improvement during continuing therapy in patients who have shown initial treatment response.

Olanzapine is indicated for short-term treatment of acute manic episode associated with Bipolar I Disorder.

Olanzapine is indicated for preventing recurrence of manic, mixed or depressive episodes in Bipolar I Disorder.

RECOMMENDED DOSAGE:

Schizophrenia

The recommended starting dose for olanzapine is 10 mg/day.

Manic episode

The starting dose is 15 mg as a single daily dose in monotherapy or 10 mg daily in combination therapy.

Preventing recurrence in bipolar disorder

The recommended starting dose is 10mg/day. For patients who have been receiving olanzapine for treatment of manic episode, continue therapy for preventing recurrence at the same dose. If a new manic, mixed, or depressive episode occurs, olanzapine treatment should be continued (with dose optimisation as needed), with supplementary therapy to treat mood symptoms, as clinically indicated.

During treatment for both schizophrenia, manic episode and recurrence prevention in bipolar disorder, daily dosage may subsequently be adjusted on the basis of individual clinical status within the range 5 – 20 mg/day. An increase to a dose greater than the recommended starting dose is advised only after appropriate clinical reassessment and should generally occur at intervals of not less than 24 hours. Olanzapine can be given without regards for meals as absorption is not affected by food. Gradual tapering of the dose should be considered when discontinuing olanzapine.

Children

Olanzapine is not recommended for use in children and adolescents below 18 years of age due to a lack of data on safety and efficacy. A greater magnitude of weight gain, lipid and prolactin alterations has been reported in short term studies of adolescent patients than in studies of adults patients.

Elderly patients

A lower starting dose (5 mg/day) is not routinely indicated but should be considered for those 65 and over when clinical factors warrant.

Patients with renal and/or hepatic impairment

A lower starting dose (5 mg) should be considered for such patients. In cases of moderate hepatic insufficiency (cirrhosis, Child-Pugh Class A or B), the starting dose should be 5 mg and only increased with caution.

Gender

The starting dose and dose range need not be routinely altered for female patients relative to male patients.

Smokers

The starting dose and dose range need not be routinely altered for non-smoking patients relative to smoking patients. When more than one factor is present which might result in slower metabolism (female gender, geriatric age, non-smoking status), consideration should be given to decreasing the starting dose. Dose escalation, when indicated, should be conservative in such patients.

MODE OF ADMINISTRATION:

Oral

CONTRAINDICATIONS:

Hypersensitivity to olanzapine or any of the excipients. Patients with known risk for narrow-angle glaucoma.

WARNINGS & PRECAUTIONS:

Hyperglycemia and Diabetes Mellitus

Hyperglycemia in some extreme cases and associated with ketoacidosis or hyperosmolar coma or death, has been reported in patients treated with atypical antipsychotics. Assessment of the relationship between atypical antipsychotics use and glucose abnormalities is complicated by the possibility of an increased background risk of diabetes mellitus in patients with schizophrenia and the increasing incidence of diabetes mellitus in the general population. Given this confounders, the relationship between atypical antipsychotic use and hyperglycemia-related adverse events is not completely understood. However, epidemiological studies suggest an increased risk of treatment-emergent hyperglycemia-related events in patients treated with the atypical antipsychotics. Precise risk estimates for hyperglycemia-related adverse events in patients treated with atypical antipsychotics are not available.

Patients with an established diagnosis of diabetes mellitus who are started on atypical antipsychotics should be monitored regularly for worsening of glucose control. Patients with risk factors for diabetes mellitus (e.g. obesity, family history of diabetes) who are starting treatment with atypical antipsychotics should undergo fasting blood glucose testing at the beginning of treatment and periodically during treatment. Any patient treated with atypical antipsychotics should be monitored for symptoms of hyperglycemia including polydipsia, polyuria, polyphagia and weakness. Patients who develop symptoms of hyperglycemia during treatment with atypical antipsychotics should undergo fasting blood glucose testing. In some case, hyperglycemia has resolved when the atypical antipsychotics was discontinued; however, some patients required continuation of anti-diabetic treatment despite discontinuation of the suspect drug.

During antipsychotics treatment, improvement in the patient's clinical condition may take several days to some weeks. Patients should be closely monitored during this period.

Dementia-related psychosis and/or behavioural disturbances

Olanzapine is not approved for the treatment of dementia-related psychosis and/or behavioural disturbances and is not recommended for use in this particular group of patients because of an increase in mortality and the risk of cerebrovascular accident. The higher incidence of death was not associated with olanzapine dose (mean daily dose 4.4 mg) or duration of

treatment. Risk factors that may predispose this patient population to increased mortality include age >65 years, dysphagia, sedation, malnutrition and dehydration, pulmonary conditions (e.g. pneumonia, with or without aspiration), or concomitant use of benzodiazepines.

Cerebrovascular adverse events (CVAE e.g. stroke, transient ischemic attack), including fatalities, were reported.

Parkinson's disease

The use of olanzapine in the treatment of dopamine agonist associated psychosis in patients with Parkinson's disease is not recommended.

Neuroleptic Malignant Syndrome (NMS)

NMS is a potentially life-threatening condition associated with antipsychotic medicinal product. Rare cases reported as NMS have also been received in association with olanzapine. Clinical manifestations of NMS are hyperpyrexia, muscle rigidity, altered mental status, and evidence of autonomic instability (irregular pulse or blood pressure, tachycardia, diaphoresis, and cardiac dysrhythmia). Additional signs may include elevated creatine phosphokinase, myoglobinuria (rhabdomyolysis), and acute renal failure. If a patient develops signs and symptoms indicative of NMS, or presents with unexplained high fever without additional clinical manifestations of NMS, all antipsychotic medicines, including olanzapine must be discontinued.

Lipid alterations

Lipid alterations should be managed as clinically appropriate, particularly in dyslipidemic patients and in patients with risk factors for the development of lipids disorders. Patients treated with any antipsychotic agents, including olanzapine, should be monitored regularly for lipids in accordance with utilised antipsychotic guidelines.

Anticholinergic activity

Caution is advised when prescribing for patients with prostatic hypertrophy, or paralytic ileus and related conditions.

Hepatic function

Transient, asymptomatic elevations of hepatic transaminases, ALT, AST have been seen commonly, especially in early treatment. Caution should be exercised in patients with elevated ALT and/or AST, in patients with signs and symptoms of hepatic impairment, in patients with pre-existing conditions associated with limited hepatic functional reserve, and in patients who are being treated with potentially hepatotoxic medicines. In the event of elevated ALT and/or AST during treatment, follow-up should be organised and dose reduction should be considered. In cases where hepatitis (including hepatocellular, cholestatic or mixed liver injury) has been diagnosed, olanzapine treatment should be discontinued.

Neutropenia

Caution should be exercised in patients with low leukocyte and/or neutrophil counts for any reason, in patients receiving medicines known to cause neutropenia, in patients with a history of drug-induced bone marrow depression/toxicity, in patients with bone marrow depression caused by concomitant illness, radiation therapy or chemotherapy and in patients with hypereosinophilic conditions or with myeloproliferative disease. Neutropenia has been reported commonly when olanzapine and valproate are used concomitantly.

Discontinuation of treatment

Acute symptoms such as sweating, insomnia, tremor, anxiety, nausea, or vomiting have been reported very rarely when olanzapine is stopped abruptly.

QT interval

As with other antipsychotics, caution should be exercised when olanzapine is prescribed with medicines known to increase QTc interval, especially in the elderly, in patients with congenital long QT syndrome, congestive heart failure, heart hypertrophy, hypokalaemia or hypomagnesaemia.

Thromboembolism

Temporal association of olanzapine treatment and venous thromboembolism has very rarely been reported. A causal relationship between the occurrence of venous thromboembolism and treatment with olanzapine has not been established. However, since patients with schizophrenia often present with acquired risk factors for venous thromboembolism all possible risk factors of VTE e.g. immobilisation of patients, should be identified and preventive measures undertaken.

General CNS activity

Given the primary CNS effects of olanzapine, caution should be used when it is taken in combination with other centrally acting medicines and alcohol. As it exhibits in vitro dopamine antagonism, olanzapine may antagonize the effects of direct and indirect dopamine agonists.

Seizures

Olanzapine should be used cautiously in patients who have a history of seizures or are subject to factors which may lower the seizure threshold. Seizures have been reported to occur rarely in patients when treated with olanzapine. In most of these cases, a history of seizures or risk factors for seizures were reported.

Tardive Dyskinesia

Olanzapine was associated with a statistically significant lower incidence of treatment emergent dyskinesia. However the risk of tardive dyskinesia increases with long term exposure, and therefore if signs or symptoms of tardive dyskinesia appear in a patient on olanzapine, a dose reduction or discontinuation should be considered. These symptoms can temporarily deteriorate or even arise after discontinuation of treatment.

Postural hypotension

As with other antipsychotics, it is recommended that blood pressure is measured periodically in patients over 65 years to avoid postural hypotension.

Use in children and adolescents under 18 years of age

Olanzapine is not indicated for use in the treatment of children and adolescents.

Effects on ability to drive and use machines

Because olanzapine may cause somnolence and dizziness, patients should be cautioned about operating hazardous machinery, including motor vehicles.

Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)

Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS) has been reported with olanzapine exposure. DRESS consists of a combination of three or more of the following: cutaneous reaction (such as rash or exfoliative dermatitis), eosinophilia, fever lymphadenopathy and one or more systemic complications such as hepatitis, nephritis, pneumonitis, myocarditis, and pericarditis. Discontinue olanzapine if DRESS is suspected.

INTERACTIONS WITH OTHER MEDICAMENTS:

Caution should be exercised in patients who receive medicinal products that can cause central nervous system depression.

Potential interactions affecting olanzapine

Since olanzapine is metabolised by CYP1A2, substances that can specifically induce or inhibit this isoenzyme may affect the pharmacokinetics of olanzapine.

Induction of CYP1A2

The metabolism of olanzapine may be induced by smoking and carbamazepine, which may lead to reduced olanzapine concentrations. Only slight to moderate increase in olanzapine clearance has been observed. The clinical consequences are likely to be limited, but clinical monitoring is recommended and an increase of olanzapine dose may be considered if necessary.

Inhibition of CYP1A2

Fluvoxamine, a specific CYP1A2 inhibitor, has been shown to significantly inhibit the metabolism of olanzapine. A lower starting dose of olanzapine should be considered in patients who are using fluvoxamine or any other CYP1A2 inhibitors, such as ciprofloxacin or ketoconazole. A decrease in the dose of olanzapine should be considered if treatment with an inhibitor of CYP1A2 is initiated.

Decreased bioavailability

Activated charcoal reduces the bioavailability of oral olanzapine by 50 to 60% and should be taken at least 2 hours before or after olanzapine. Fluoxetine (a CYP2D6 inhibitor), single doses of antacid (aluminium, magnesium) or cimetidine have not been found to significantly affect the pharmacokinetics of olanzapine.

Potential for olanzapine to affect other medicinal products

Olanzapine may antagonise the effects of direct and indirect dopamine agonists. No inhibition of metabolism of the following active substances was found: tricyclic antidepressant (representing mostly CYP2D6 pathway), warfarin (CYP2C9), theophylline (CYP1A2) or diazepam (CYP3A4 and 2C19). Olanzapine showed no interaction when co-administered with lithium or biperiden. Therapeutic monitoring of valproate plasma levels did not indicate that valproate dosage adjustment is required after the introduction of concomitant olanzapine.

STATEMENT ON USAGE DURING PREGNANCY AND LACTATION:

Neonates exposed to antipsychotic drugs during the third trimester of pregnancy are at risk for extrapyramidal and/or withdrawal symptoms following delivery. There have been reports of agitation, hypertonia, hypotonia, tremor, somnolence, respiratory distress and feeding disorder in these neonates. These complications have varied in severity; while in some cases symptoms have been self-limited, in other cases neonates have required intensive care unit support and prolonged hospitalisation.

Olanzapine tablet should be used during pregnancy only if the potential benefit justifies the potential risk to the foetus.

There are no adequate and well-controlled studies in pregnant women. Patients should be advised to notify their physician if they become pregnant or intend to become pregnant during treatment with olanzapine.

Patients should be advised not to breast feed an infant if they are taking olanzapine.

ADVERSE EFFECTS/ UNDESIRABLE EFFECTS:

Adults

The most frequently reported adverse reactions associated with the use of olanzapine are somnolence, weight gain, eosinophilia, elevated prolactin, cholesterol, glucose and triglyceride levels, glucosuria, increased appetite, dizziness, akathisia, parkinsonism, dyskinesia, orthostatic hypotension, anticholinergic effects, transient asymptomatic elevations of hepatic transaminases, rash, asthenia, fatigue and oedema.

The following adverse reactions and laboratory investigations has been reported with olanzapine.

Blood and the lymphatic system disorders

Common : Eosinophilia

Uncommon: Leucopenia, neutropenia

Not known : Thrombocytopenia

Immune system disorders

Not known :Allergic reaction

Metabolism and nutrition disorders

Very common: Weight gain

Common: Elevated cholesterol levels, elevated glucose levels, elevated triglyceride levels, glucosuria, increased appetite

Not known: Development or exacerbation of diabetes occasionally associated with ketoacidosis or coma, including some fatal cases, hypothermia

Nervous system disorders

Very common: Somnolence

Common: Dizziness, akathisia, parkinsonism, dyskinesia

Not known: Seizures where in most cases a history of seizures or risk factors for seizures were reported, neuroleptic malignant syndrome, dystonia (including oculogyration), tardive dyskinesia, discontinuation symptoms

Cardiac disorders

Uncommon: Bradycardia, QTc prolongation

Not known: Ventricular tachycardia / fibrillation, sudden death

Vascular disorders

Common: Orthostatic hypotension

Not known: Thromboembolism (including pulmonary embolism and deep vein thrombosis)

Gastrointestinal disorders

Common: Mild, transient anticholinergic effects including constipation and dry mouth

Not known: Pancreatitis

Hepato-biliary disorders

Common: Transient, asymptomatic elevations of hepatic transaminases (ALT, AST), especially in early treatment.

Not known: Hepatitis (including hepatocellular, cholestatic or mixed liver injury)

Skin and subcutaneous tissue disorders

Common: Rash

Uncommon: Photosensitivity reaction, alopecia

Very rare: Drug reaction with Eosinophilia and Systemic Symptoms (DRESS)

Musculoskeletal and connective tissue disorders

Not known: Rhabdomyolysis

Renal and urinary disorders

Uncommon: Urinary incontinence

Not known: Urinary hesitation

Reproductive system and breast disorders

Not known: Priapism

General disorders and administration site conditions

Common: Asthenia, fatigue, oedema

Investigations

Very common: Elevated plasma prolactin levels

Uncommon: High creatinine phosphokinase, increased total bilirubin

Not known: Increased alkaline phosphatase

Long-term exposure (at least 48 weeks)

Patients who had adverse, clinically significant changes in weight gain, glucose, total/LDL/HDL cholesterol or triglycerides increased over time. In adult patients on long-term therapy for 9-12 months, the rate of increase in mean blood glucose slowed after approximately 6 months.

Other reported adverse events are: Personality disorder, dyspepsia, accidental injury, back pain, chest pain, hypertension, ecchymosis, extremity pain, joint pain, articulation impairment, rhinitis, cough increased, pharyngitis, amblyopia, increased salivation, amnesia, paresthesia, thirst, apathy, confusion, euphoria, in coordination, dyspnea, acne, dry skin, abnormal vision, dysmenorrhea, vaginitis, chills, suicide attempt, hangover effect, vasodilation, tongue edema, ileus, intestinal obstruction, liver fatty deposit, hypoproteinemia, osteoporosis, ataxia, stupor, epistaxis, lung edema, abnormality of accommodation, dry eyes, mydriasis, amenorrhea, breast pain, decreased menstruation, impotence, menorrhagia, metrorrhagia, polyuria, urinary frequency, urinary retention, urination impaired, injection site pain, syncope, creatinine phosphokinase increased, sinusitis, arthralgia.

Additional information on special populations

In elderly patients with dementia, olanzapine treatment is associated with a higher incidence of death and cerebrovascular adverse reactions. Very common adverse reactions associated with the use of olanzapine in this patient group were abnormal gait and falls. Pneumonia, increased body temperature, lethargy, erythema, visual hallucinations and urinary incontinence were observed commonly.

In patients with drug-induced (dopamine agonist) psychosis associated with Parkinson's disease, worsening of Parkinsonian symptomatology and hallucinations were reported.

In patients with bipolar mania, valproate combination therapy with olanzapine neutropenia has been reported; a potential contributing factor could be high plasma valproate levels. Olanzapine administered with lithium or valproate resulted in increased levels of tremor, dry mouth, increased appetite, and weight gain. Speech disorder was also reported commonly. During treatment with olanzapine in combination with lithium or divalproex up to 6 weeks, an increase of baseline body weight has been reported. Also, long-term olanzapine treatment (up to 12 months) was associated with an increase from baseline body weight.

Paediatrics

Olanzapine is not indicated for the treatment of children and adolescent patients below 18 years.

The adverse reactions have been reported with a greater frequency in adolescent patients (aged 13-17 years). Clinically significant weight gain occurs more frequently in the adolescent population. The magnitude of weight gain is greater with long-term exposure (at least 24 weeks) than with short-term exposure.

Metabolism and nutrition disorders

Very common: Weight gain, elevated triglyceride levels, increased appetite.

Common: Elevated cholesterol levels.

Nervous system disorders

Very common: Sedation (including : hypersomnia, lethargy, somnolence).

Gastrointestinal disorders

Common: Dry mouth

Hepato-biliary disorders

Very common: Elevations of hepatic transaminases (ALT/AST)

Investigations

Very common: Decreased total bilirubin, increased GGT, elevated plasma prolactin levels.

ECG Changes

In adults as well as adolescents, no significant differences are reported in ECG parameters, including QT, QTc (Fridericia corrected), and PR intervals. Olanzapine use is associated with a mean increase in heart rate.

OVERDOSE AND TREATMENT:*Signs and Symptoms*

Very common symptoms in overdose include tachycardia, agitation/ aggressiveness, dysarthria, various extrapyramidal symptoms, and reduced level of consciousness ranging from sedation to coma.

Other medically significant sequelae of overdose include delirium, convulsion, coma, possible neuroleptic malignant syndrome, respiratory depression, aspiration, hypertension or hypotension, cardiac arrhythmias and cardiopulmonary arrest.

Management of Overdose

There is no specific antidote for olanzapine. Induction of emesis is not recommended. Standard procedures for management of overdose may be indicated (i.e. gastric lavage, administration of activated charcoal). The concomitant administration of activated charcoal was shown to reduce the oral bioavailability of olanzapine by 50 to 60%.

Symptomatic treatment and monitoring of vital organ function should be instituted according to clinical presentation, including treatment of hypotension and circulatory collapse and support of respiratory function. Do not use epinephrine, dopamine, or other sympathomimetic agents with beta-agonist activity since beta stimulation may worsen hypotension. Cardiovascular monitoring is necessary to detect possible arrhythmias. Close medical supervision and monitoring should continue until the patient recovers.

STORAGE CONDITIONS:

Store in the original package and in a dry place. Do not store above 30°C.

Keep out of reach of children.

DOSAGE FORMS AND PACKAGING AVAILABLE:

Packing Size: **OLANZ Tablets 5mg** 50 x 10's (blister pack)

OLANZ Tablets 10mg 50 x 10's (blister pack)

NAME AND ADDRESS OF MANUFACTURER / PRODUCT REGISTRATION HOLDER:

MALAYSIAN PHARMACEUTICAL INDUSTRIES SDN BHD (101323-U)

Plot 14, Lebuhraya Kampung Jawa, 11900 Bayan Lepas,
Pulau Pinang, Malaysia.

DATE OF REVISION:

19/06/2017