

TINACORT CREAM

Description

White, odourless, opaque cream.

Composition

Miconazole Nitrate 2% w/w
Hydrocortisone 1% w/w

Pharmacodynamics

Miconazole acts as an antifungal by altering cell membrane permeability, resulting in the cell membrane being unable to function as a selective barrier. There is little systemic absorption when miconazole is applied topically.

The anti-inflammatory effect of topical application of hydrocortisone, a less potent corticosteroid, is due to its ability to diffuse across cell membranes and subsequently enters the cell nucleus. It then binds with DNA, and stimulates transcription of messenger RNA resulting in protein synthesis of various enzymes. It is absorbed percutaneously, particularly in denuded areas. Occlusion or use over extensive areas increases topical absorption.

Pharmacokinetics

Tinacort combines an antifungal activity against the common dermatophytes, yeasts and various other fungi with an antibacterial activity against certain gram positive bacilli and cocci. Tinacort inhibits the biosynthesis of ergosterol in fungi and changes the composition of other lipid components in the membrane, resulting in fungi cell necrosis.

Miconazole in combination with hydrocortisone acts very rapidly on pruritus, which frequently accompanies dermatophyte and yeast infections. This symptomatic improvement is seen before the first signs of healing are observed. However, treatment with hydrocortisone is

symptomatic and lesions may flare up again after a discontinuation of the treatment.

Miconazole does not produce detectable blood levels after topical application. Depending on the location and type of infection, hydrocortisone penetrates well into the skin and is partially absorbed. More than 90% of the absorbed fraction is bound to plasma proteins. The metabolism takes place in the liver and tissues and the metabolites are excreted with the urine, mostly as glucuronides, together with a very small fraction of unchanged hydrocortisone. The biological half life is about 100 minutes.

Indications

Infection of the skin by dermatophytes of *Candida* spp. in which inflammatory symptoms are prominent.

Thus, Tinacort Cream is particularly indicated for the initial stages of treatment. Once the inflammatory symptoms have disappeared, treatment may be continued with miconazole nitrate 20mg/g topical cream if preferred. In view of Tinacort Cream's antibacterial effect on gram-positive bacteria, the product may also be used for myotic affections with bacterial superinfection.

Contraindications

True hypersensitivity to any of the ingredients. Tubercular or viral infections of the skin or those caused by Gram-negative bacteria.

Warning and Precautions

If a reaction suggesting sensitivity or irritation should occur, the treatment should be discontinued. Avoid the introduction of Tinacort cream into eyes. As with any topical corticosteroid, care is advised with infants and children when Tinacort is to be applied to extensive surface areas or under occlusive dressings including baby napkins.

Similarly application to the face should be avoided. In infants, long term continuous topical corticosteroid

therapy should be avoided. Adrenal suppression can occur even without occlusion.

Pregnancy and Lactation

Caution is recommended during pregnancy and lactation period. Treatment of large surfaces and the application under occlusive dressing should be avoided during that time.

Interaction with Other Medicaments and Other Forms of Interaction

None known

Main Side/ Adverse Effects

Tinacort cream is usually well tolerated.

There have been isolated reports of irritation or burning sensation associated with the application of Tinacort cream. As with all substances applied to the skin, an allergic reaction to any of the ingredient of the cream may occur.

Overdose

Prolonged and excessive use can result in skin irritation which usually disappears after discontinuation of therapy. Topical applied, corticosteroids can be absorbed in sufficient amounts to produce systemic effects.

If accidental ingestion of large quantities of the product occurs, an appropriate method of gastric emptying may be used if considered to be necessary.

Dosage and Administration

Tinacort cream should be applied topically to the lesion once to twice daily. It should be rubbed in gently with finger until it has been completely penetrated into the skin. The treatment with Tinacort cream (or subsequently with miconazole nitrate 20mg/g topical cream) should be continued without interruption until the lesion has completely disappeared (usually after 2 to 5 weeks). This dosage applies to both adults and children.

Special populations

Pediatrics

In infants and children, caution is advised when Tinacort cream is applied to extensive surface areas or under occlusive dressings including baby napkins (diapers). In infants, long term continuous topical corticosteroid therapy should be avoided (see Warnings and Precautions).

Elderly

Natural thinning of the skin occurs in the elderly, hence corticosteroids should be used sparingly and for short periods of time.

Note: The information given here is limited. For further information, consult your doctor or pharmacist.

Storage : Store below 25 degree Celsius

Presentation/
packing : 15 gram in Aluminium Tube

Product
Registration
Holder : Hovid Bhd., 121, Jalan Tunku
Abdul Rahman, 30010 Ipoh,
Malaysia

Manufactured
by : Hovid Bhd., 121, Jalan Tunku
Abdul Rahman, 30010 Ipoh,
Malaysia

Revision date : August 2017