



# DYNA FUSIDIC CREAM 2%

## MAL18026194AZ

### CONTENT & DESCRIPTION

Dyna Fusidic Cream contains fusidic acid 2% w/w.  
White cream with natural fragrance.  
FOR EXTERNAL (TOPICAL) USE ONLY

### PHARMACOLOGICAL PROPERTIES

#### Pharmacodynamic properties

Fusidic acid is a potent antibacterial agent. Fusidic acid and its salts show fat and water solubility and strong surface activity and exhibit unusual ability to penetrate intact skin. Concentrations of 0.03 - 0.12 mcg fusidic acid per ml inhibit nearly all strains of *Staphylococcus aureus*. Topical application of fusidic acid is also effective against streptococci, corynebacteria, neisseria and certain clostridia.

#### Pharmacokinetic properties

Fusidic acid is excreted mainly in the bile with little excreted in the urine.

### CLINICAL PARTICULARS

#### Therapeutic indications

DYNA FUSIDIC CREAM is indicated for treatment of skin infections caused by staphylococci, streptococci, *Propionibacterium acnes*, *Corynebacterium minutissimum*, and other organisms sensitive to fusidic acid. The most important indications being: impetigo, infected wounds, folliculitis, boils, sycosis barbae, carbuncles, hidradenitis, paronychia, erythrasma.

DYNA FUSIDIC CREAM is a non-staining preparation, it is cosmetically acceptable for treatment of face and scalp infections.

#### Direction for Use

Apply DYNA FUSIDIC CREAM to the affected area 2-3 times daily, generally for a period of 7 days, except in treatment of acne where therapy should be extended for a longer period according to the nature of this disease.

#### Contraindications

Known hypersensitivity to fusidic acid/sodium fusidate or to any of the excipients (refer section on 'List of Excipients').

Infection caused by non-susceptible organisms.

#### Special warnings and precautions for use

Bacterial resistance has been reported to occur with the use of fusidic acid. As with all antibiotics, extended or recurrent use may increase the risk of developing antibiotic resistance.

Extended or recurrent use may increase the risk of developing contact sensitisation.

Fusidic acid does not appear to cause conjunctival irritation in experimental animals. Caution should still be exercised, however, when this cream is used near the eyes.

#### Interactions with Other Medicaments

None known

#### Pregnancy and lactation

##### Pregnancy

No effects during pregnancy are anticipated, since systemic exposure to topically-applied fusidic acid/sodium fusidate is negligible. Topical Fusidic Acid Cream can be used during pregnancy.

##### Breast-feeding

No effects on the breast-fed new-born/infant are anticipated since the systemic exposure of topically-applied fusidic acid/sodium fusidate to the breast-feeding woman is negligible. Topical Fusidic Acid Cream can be used during breast-feeding but it is recommended to avoid applying topical Fusidic Acid Cream on the breast.

#### Effects on ability to drive and use machines

Fusidic Acid Cream administered topically has no or negligible influence on the ability to drive and to use machines.

### Side effects

The most frequently reported adverse reactions during treatment are various skin reactions such as pruritus and rash, followed by application site conditions such as pain and irritation. Hypersensitivity and angioedema have been reported.

|                                                      |                                                                                                                                                                                                                                                      |
|------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Immune system disorders                              |                                                                                                                                                                                                                                                      |
| Rare                                                 | Hypersensitivity                                                                                                                                                                                                                                     |
| Eye disorders                                        |                                                                                                                                                                                                                                                      |
| Rare                                                 | Conjunctivitis                                                                                                                                                                                                                                       |
| Skin and subcutaneous tissue disorders               |                                                                                                                                                                                                                                                      |
| Uncommon                                             | Dermatitis (including dermatitis contact, eczema)<br>Rash*<br>Pruritus<br>Erythema<br>*Various types of rash reactions such as erythematous, pustular, vesicular, maculo-papular and papular have been reported. Rash generalised has also occurred. |
| Rare                                                 | Angioedema<br>Urticaria<br>Blister                                                                                                                                                                                                                   |
| General disorders and administration site conditions |                                                                                                                                                                                                                                                      |
| Uncommon                                             | Application site pain (including skin burning sensation)<br>Application site irritation                                                                                                                                                              |

### Paediatric population

Frequency, type and severity of adverse reactions in children are expected to be the same as in adults.

### Overdose

Overdose is unlikely to occur.

Unless hypersensitivity to Fusidic acid or any of the excipients exists, accidental ingestion of Fusidic Acid Cream is unlikely to cause any harm. The total quantity of fusidic acid (30g Fusidic Acid Cream contains 600mg fusidic acid) will usually not exceed the approved total daily oral dose of fusidic acid containing products except in children aged less than 1 year and weighing  $\leq 10$ kg. Although in this instance a child of this particular age group is unlikely to ingest a whole tube of Fusidic Acid Cream. The concentration of the excipients is too low to constitute a safety risk.

### PHARMACEUTICAL PARTICULARS

#### List of excipients

Glycerine, Chlorocresol, Cetomacrogol  
Emulsifying Wax, Liquid Paraffin, Propylene  
Glycol, White Soft Paraffin and Purified Water

#### Shelf life

3 years from date of manufacture.

#### Packing / Pack size(s)

Aluminium tubes of 15 gram.

#### Storage

JAUHI UBAT DARIPADA KANAK-KANAK  
KEEP OUT OF REACH OF CHILDREN  
Keep Container Tightly Closed  
Store in A Dry Place below 30°C  
Protect From Light

### MANUFACTURER & PRODUCT REGISTRATION HOLDER:

**DYNAPHARM (M) SDN. BHD.** 198001011897 (65683-V)

2497, Mukim 1, Lorong Perusahaan Baru 5, Kawasan Perusahaan Perai 3,  
13600 Perai, Pulau Pinang, MALAYSIA

PI1PE D011-01

**Latest Revision:** Feb 2022