

BAROSCON® DOUBLE ACTION Suspension

Ingredient(s):

Each sachet (10ml) contains:

Sodium Alginate	500mg
Calcium Carbonate	325mg
Sodium Bicarbonate	213mg
Methyl Parahydroxybenzoate (preservative)	28mg
Propyl Parahydroxybenzoate (preservative)	4mg

Product Description:

Peppermint-flavoured, pale yellow or cream-coloured suspensions contained in aluminium foil pouch.

Indication:

Treatment of symptoms of gastro-oesophageal reflux such as acid regurgitation, heartburn and indigestion, and for symptoms of excess stomach acid (hyperacidity).

Recommended Dose:

Adults and children aged 12 and above: 1 to 2 sachets after meals and before bedtime.

Pharmacokinetics:

The mode of action of the medicinal product is physical and does not depend on absorption into the systemic circulation.

Pharmacodynamics:

Pharmacotherapeutic classification: other drugs for peptic ulcer and gastro-oesophageal reflux disease.

The medicinal product is a combination of two antacids (calcium carbonate and sodium bicarbonate) and an alginate.

On ingestion, the medicinal product reacts rapidly with gastric acid to form a raft of alginic acid gel having a near neutral pH and which floats on the stomach contents effectively impeding gastro-oesophageal reflux, for up to 4 hours.

In severe cases the raft itself may be refluxed into the oesophagus, in preference to the stomach contents, and exert a demulcent effect. Calcium carbonate neutralizes gastric acid to provide fast relief from indigestion and heartburn. This effect is increased by the addition of sodium bicarbonate which also has a neutralising action.

Mode of Administration:

For oral administration.

Interaction with Other Medicaments:

Due to the presence of calcium carbonate which act as an antacid, a time-interval of 2 hours should be considered between the medicinal product intake and the administration of other medicinal products, especially H₂-antihistaminics, tetracyclines, digoxine, fluoroquinolone, iron salt, ketoconazole, neuroleptics, thyroxine, penicillamine, beta-blockers (atenolol, metoprolol, propranolol), glucocorticoid, chloroquine, and diphosphonates.

Pregnancy & Lactation:

There is no significant adverse effects of the product known on the course of pregnancy or on the health of the foetus/new-born child. The medicinal product may be used during pregnancy and lactation.

Contraindication:

Hypersensitivity, including the esters of hydroxybenzoates (parabens).

Item Code	: BRSs - 03 draft 27 Jun 2024
PM Code	: Ins.P - BRSs
Measurement(mm) ± 2mm	: 270(L) x 110(W)
Colour of Artwork	: K
Source of Artwork	: MAC 2 / CC

H
270mm
(±2mm)

Item Code	: BRSs - 03 draft 27 Jun 2024
PM Code	: Ins.P - BRSs
Measurement(mm) ± 2mm	: 270(L) x 110(W)
Colour of Artwork	: K
Source of Artwork	: MAC 2 / CC

Warning and Precautions:

Each 10ml (single sachet) dose has a sodium content of 127.9mg (5.56mmol). This should be taken into account when a highly restricted salt diet is recommended, e.g. in some cases of congestive cardiac failure and renal impairment.

Each 10ml (single sachet) dose contains 130mg (3.25mmol) of calcium. Care needs to be taken in treating patients with hypercalcaemia, nephrocalcinosis and recurrent calcium containing renal calculi.

Treatment of children younger than 12 years of age is not generally recommended, except on medical advice.

If symptoms do not improve after seven days, the clinical situation should be reviewed.

Contains methyl parahydroxybenzoate (E218) and propyl parahydroxybenzoate (E216) which may cause allergic reactions (possibly delayed).

Side Effects:

Very rarely (<1/10,000) patients sensitive to the ingredients may develop allergic manifestations such as urticaria or bronchospasm, anaphylactic or anaphylactoid reactions.

Ingestion of large quantities of calcium carbonate may cause alkalosis, hypercalcaemia, acid rebound, milk alkali syndrome or constipation. These usually occur following larger than recommended dosages.

Symptoms and Treatment of Overdose:

In the event of overdose symptomatic treatment should be given. The patient may notice abdominal distension.

Storage Condition:

Store at temperature below 30°C. Protect from direct sunlight and moisture.

Shelf Life:

2 years from the date of manufacture.

Packing(s):

Sachet packing of 10ml x 4's, 10ml x 5's, 10ml x 16's and 10ml x 24's.

H
270mm
(±2mm)



Manufacturer:
Yungjin Pharm. Co. Ltd.,
66, Muha-ro, Namyang-eup, Hwaseong-si,
Gyeonggi-do, Republic of Korea.

Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7 & 12, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679

xxxxxx printing code xxxxxx

Date of revision: 28 Feb 2020

W
110mm (±2mm)