
UNIDINE

Povidone Iodine Solution 10% w/v

Name and Strength of Active Ingredient:

Povidone Iodine 10% w/v (Available Iodine 1% w/v)

Product Description:

A deep brown liquid.

Pharmacodynamics:

Povidone iodine is a complex of iodine, which shows all the broad spectrum germicidal activity of elemental iodine. The germicidal activity is maintained in the presence of blood, pus, serum and necrotic tissue.

Pharmacokinetics:

Povidone Iodine is slightly absorbed when applied to the skin. Iodides are excreted mainly in the urine, with smaller amounts appearing in the faeces, saliva and sweat.

Indications:

Disinfectant and antiseptic for the treatment of contaminated wounds, cuts and burns and also for the pre-operative preparation of the skin and mucous membrane.

Recommended Dose:

To be used undiluted. Apply directly to skin. Allow to dry before applying dress, drape or cast.

Route of administration:

For topical administration.

Contraindications:

Hypersensitivity to iodine, polyvinylpyrrolidone or to any excipient. History of abnormal thyroid function or goitre (in particular nodular colloid goitre, endemic goitre and Hashimoto's thyroiditis). Use in children under two years of age. Regular use should be avoided in patients on concurrent lithium therapy.

Warning and Precautions:

Use of this preparation may interfere with tests of thyroid function. Iodine is absorbed through burns and broken skin and to a lesser extent through intact skin and may lead to toxic levels of iodine in the blood, particularly in patients with renal insufficiency. If symptoms occur suggesting changes in thyroid function, these should be investigated. In patients with impaired renal function, blood levels of iodine should be monitored.

Interactions with Other Medicaments:

Use with concurrent lithium therapy has been shown to exhibit additive hypothyroidic effects. Absorption of iodine from povidone iodine through either intact skin or broken skin may interfere with thyroid function tests. Contamination with povidone iodine of several types of tests for the detection of occult blood in faeces or blood in urine may produce false- positive results.

Pregnancy and Lactation:

Iodine freely crosses the placenta and is secreted in breast milk. Therefore it is recommended that breastfeeding should stop during treatment, unless no alternative is possible. Thyroid function disorders have been reported in the offspring of mothers exposed to pharmacological doses of iodine. Povidone iodine should not be used regularly during pregnancy unless there is no alternative treatment available.

Side Effects:

Local irritation, skin burns and sensitivity reactions have been reported rarely. Anaphylactic reactions, anaphylactoid reactions and anaphylactic shock have been reported uncommonly with products containing povidone iodine or povidone. Excess iodine can produce goitre and hypothyroidism or hyperthyroidism. Such effects have

occasionally been seen with extensive or prolonged use of povidone iodine. Other effects that have been reported are metabolic acidosis and acute renal failure.

Symptoms and Treatment of Overdose:

Excess iodine can produce goitre and hypothyroidism or hyperthyroidism. Systemic absorption of iodine after repeated application of povidone iodine to large areas of wounds or burns may lead to a number of adverse effects: Metallic taste in mouth, increased salivation, burning or pain in the throat or mouth, irritation and swelling of the eyes, pulmonary oedema, skin reactions, gastrointestinal upset and diarrhea, metabolic acidosis, hypernatraemia and renal impairment.

Deliberate or accidental ingestion of large quantities of povidone iodine will result in high blood levels of iodine and gastrointestinal toxicity including vomiting and diarrhoea. Symptomatic and supportive treatment should be started with special attention to monitoring electrolyte balance, renal function, thyroid function and liver function. Iodine can be removed by dialysis.

Effects on Ability to Drive and Use Machine:

None known.

Instructions for Use:

For topical use on the skin only.

Storage Condition:

Store below 30°C & Protect from light.

Pack Size:

100ml, 200ml, 250ml, 500ml and 1000ml HDPE bottle.

Registration Number:

MAL18056086XZ

Product Registration Holder:

KCK Pharmaceutical Industries Sdn. Bhd.

Plot 61, Bayan Lepas Industrial Park Phase 4,
Lintang Bayan Lepas 1, 11900 Bayan Lepas,
Pulau Pinang, Malaysia.

Manufactured by:

KCK Pharmaceutical Industries Sdn. Bhd.

No. 5 & 7 Hala Beringin,
Diamond Valley Industrial Park, 11960 Bayan Lepas,
Pulau Pinang, Malaysia.

Date of Revision:

02 January 2024