

System Organ class	Adverse drug reaction
Infections and infestations	
Common	Nasopharyngitis
Blood and lymphatic system disorders	
Uncommon	Neutropaenia
Immune system disorders	
Uncommon	Hypersensitivity
Rare	Angioedema, allergic reaction
Metabolism and nutrition disorders	
Common	Appetite increased
Un common	Anorexia, hypoglycaemia
Psychiatric disorders	
Common	Euphoric mood, confusion, irritability, disorientation, insomnia, libido decreased
Uncommon	Hallucination, panic attack, restlessness, agitation, depression, depressed mood, elevated mood, <i>aggression</i> , mood swings, depersonalization, word finding difficulty, abnormal dreams, libido increased, anorgasmia, apathy
Rare	Disinhibition
Nervous system disorders	
Very Common	Dizziness, somnolence, headache
Common	Ataxia, coordination abnormal, tremor, dysarthria, amnesia, memory impairment, disturbance in attention, paraesthesia, hypoaesthesia, sedation, balance disorder, lethargy
Uncommon	Syncope, stupor, myoclonus, <i>loss of consciousness</i> , psychomotor hyperactivity, dyskinesia, dizziness postural, intention tremor, nystagmus, cognitive disorder, <i>mental impairment</i> , speech disorder, hyporeflexia, hyperaesthesia, burning sensation, ageusia, <i>malaise</i>
Rare	<i>Convulsions</i> , parosmia, hypokinesia, dysgraphia
Eye disorders	
Common	Vision blurred, diplopia
Uncommon	Peripheral vision loss, visual disturbance, eye swelling, visual field defect, visual acuity reduced, eye pain, asthenopia, photopsia, dry eye, lacrimation increased, eye irritation
Rare	<i>Vision loss</i> , <i>keratitis</i> , oscillopsia, altered visual depth perception, mydriasis, strabismus, visual brightness
Ear and labyrinth disorders	
Common	Vertigo
Un Common	Hyperacusis
Cardiac disorders	
Uncommon	Tachycardia, atrioventricular block first degree, sinus bradycardia, <i>congestive heart failure</i>
Rare	<i>QT prolongation</i> , sinus tachycardia, sinus arrhythmia
Vascular disorders	
Uncommon	Hypotension, hypertension, hot flushes, flushing, peripheral coldness
Respiratory, thoracic and mediastinal disorders	
Uncommon	Dyspnoea, epistaxis, cough, nasal congestion, rhinitis, snoring, nasal dryness
Rare	<i>Pulmonary oedema</i> , throat tightness
Gastrointestinal disorders	
Common	Vomiting, <i>nausea</i> , constipation, <i>diarrhoea</i> , flatulence, abdominal distension, dry mouth
Uncommon	Gastroesophageal reflux disease, salivary hypersecretion, hypoaesthesia oral
Rare	Ascites, pancreatitis, <i>swollen tongue</i> , dysphagia
Skin and subcutaneous tissue disorders	
Uncommon	Rash papular, urticaria, hyperhidrosis, <i>pruritus</i>
Rare	<i>Stevens Johnson syndrome</i> , cold sweat
Musculoskeletal and connective tissue disorders	
Common	Muscle cramp, arthralgia, back pain, pain in limb, cervical spasm
Uncommon	Joint swelling, myalgia, muscle twitching, neck pain, muscle stiffness
Rare	Rhabdomyolysis
Renal and urinary disorders	
Uncommon	Urinary incontinence, dysuria
Rare	Renal failure, oliguria, <i>urinary retention</i>
Reproductive system and breast disorders	
Common	Erectile dysfunction
Uncommon	Sexual dysfunction, ejaculation delayed, dysmenorrhoea, breast pain
Rare	Amenorrhoea, breast discharge, breast enlargement, <i>gynaecomastia</i>
General disorders and administration site conditions	
Common	Oedema peripheral, oedema, gait abnormal, fall, feeling drunk, feeling abnormal, fatigue
Uncommon	Generalised oedema, <i>face oedema</i> , chest tightness, pain, pyrexia, thirst, chills, asthenia
Investigations	
Common	Weight increased
Uncommon	Blood creatine phosphokinase increased, alanine aminotransferase increased, aspartate aminotransferase increased, blood glucose increased, platelet count decreased, blood creatinine increased, blood potassium decreased, weight decreased
Rare	White blood cell count decreased

After discontinuation of short-term and long-term treatment with Pregabalin withdrawal symptoms have been observed in some patients. The following reactions have been mentioned: insomnia, headache, nausea, anxiety, diarrhoea, flu syndrome, convulsions, nervousness, depression, pain, hyperhidrosis and dizziness, suggestive of physical dependence. The patient should be informed about this at the start of the treatment.

Concerning discontinuation of long-term treatment of Pregabalin, data suggest that the incidence and severity of withdrawal symptoms may be dose-related.

Overdose

In the postmarketing experience, the most commonly reported adverse reactions observed when Pregabalin was taken in overdose included somnolence, confusional state, agitation, and restlessness.

In rare occasions, cases of coma have been reported. Treatment of Pregabalin overdose should include general supportive measures and may include haemodialysis if necessary.

PHARMACOLOGICAL PROPERTIES

Pharmacodynamic properties

Pharmacotherapeutic group: Anti-epileptics, other anti-epileptics ATC code: N03AX16

The active substance, Pregabalin, is a gamma-aminobutyric acid analogue [(S)-3-(aminomethyl)-5-methylhexanoic acid].

Mechanism of action:

Pregabalin is an analog of the neurotransmitter GABA. It binds potently to the α2-δ subunit resulting in modulation of Ca channels and reduction in the release of several neurotransmitters, including glutamate, norepinephrine, serotonin, dopamine, calcitonin gene-related peptide and substance P. Absorption: Rapidly absorbed. Bioavailability: Approx 90%. Time to peak plasma concentration: with in 1.5 hr.

Distribution: Volume of distribution: 0.5 L/kg. Not bound to plasma protein.

Metabolism: Negligible metabolism.

Excretion: Via urine (approx 98%) as unchanged drug). Elimination half-life: 6.3 hr.

Pharmacokinetic properties

Absorption

Pregabalin is rapidly absorbed, with peak plasma concentration within 1 hour. Pregabalin oral bioavailability is ≥ 90%. The rate of Pregabalin absorption is decreased with food resulting in a decrease in Cmax by approximately 25-30% and a delay in tmax to approximately 2.5 hours. However, administration of Pregabalin with food has no significant effect on the extent of Pregabalin absorption.

Distribution

Pregabalin crosses the blood brain barrier. Pregabalin crosses the, the apparent volume of distribution of Pregabalin is approximately 0.56 l/kg. Pregabalin is not bound to plasma proteins.

Biotransformation

Pregabalin undergoes negligible metabolism.

Elimination

Pregabalin eliminates from the systemic circulation primarily by renal excretion as unchanged drug.

Pregabalin mean elimination half-life is 6.3 hours. Pregabalin plasma clearance and renal clearance are directly proportional to creatinine clearance.

Dose adjustment in patients with reduced renal function or undergoing haemodialysis is necessary.

Linearity / non-linearity

Pregabalin pharmacokinetics are linear over the recommended daily dose range. Inter-subject pharmacokinetic variability for Pregabalin is low (< 20%).

Gender

Gender does not influence on the plasma concentrations of Pregabalin.

Renal impairment

Pregabalin clearance is directly proportional to creatinine clearance.

Hepatic impairment

Since Pregabalin does not undergo significant metabolism and is excreted predominantly as unchanged drug in the urine, impaired liver function would not be expected to significantly alter Pregabalin plasma concentrations.

Elderly (over 65 years of age)

Pregabalin clearance tends to decrease with increasing age. This decrease in Pregabalin oral clearance is consistent with decreases in creatinine clearance associated with increasing age. Reduction of Pregabalin dose may be required in patients who have age related compromised renal function.

Pharmaceutical particulars

List of excipients

Capsules content

Maize starch
Talc

Capsules Shell
Iron Oxide Red (E172)
Titanium Dioxide (E171)
Gelatin
Sodium Lauryl Sulfate

Capsule printing ink:
Shellac
Black Iron Oxide (E172)

Incompatibilities
None Known.

Shelf life
Please refer outer package for expiry date.

Special precautions for storage
Store in a dry place below 30° C.

Nature and contents of container
PEGASET 50 mg/75 mg/150 mg: Blister of 28 Capsules

Manufactured by:


AUROBINDO

Aurobindo Pharma Limited,
Unit XV, Plot No 17a, E Bonangi Village,
Parawada Mandal, Anakapalli District,
Anakapalli, 531021, India.

Product Registration Holder
Healol Pharmaceuticals Sdn.Bhd.
74-3, Wangsa Delima 6, KLSC Wanga Maju,
53300 Kuala Lumpur.

DATE OF REVISION OF PACKAGE INSERT
07/2025