

pharmaniaga®

i-CALAMINE LOTION

DESCRIPTION

Pinkish red colored, smooth suspension with a strong menthol smell.

COMPOSITION

Calamine 8% w/v
Zinc Oxide 8% w/v

Preservative:

Phenol 0.1% w/v

Alcohol content:

Ethyl alcohol 0.1% v/v

PHARMACODYNAMICS

Calamine has mild astringent and antipruritic actions on the skin and is used to relieve the discomfort of dermatitis as it produces a cooling effect and helps to divert attention away from the itching and thus giving relief from skin irritations. Zinc oxide is applied externally as a mild astringent for the skin. Zinc oxide has soothing and protective properties and is used in topical preparations for eczema, slight excoriations, wounds and haemorrhoids. It also reflects ultraviolet radiation and can be used as a physical sunscreen.

PHARMACOKINETICS

Absorption

Calamine and Zinc Oxide are not absorbed through the skin.

INDICATIONS

Calamine Lotion is used for the relief of skin discomfort due to pruritis, mild sunburn, nappy rashes, and insect stings.

RECOMMENDED DOSAGE

Apply to the affected area twice or three times daily and allowed to dry. Repeated applications may be administered as needed.

Shake well before use. Wash and dry affected area before each application.

**Extremes of temperature, wearing tight fitting nappies or non-breathable clothing such as polyester or nylon can make itching worse.

ROUTE OF ADMINISTRATION

Topical

CONTRAINDICATIONS

Hypersensitivity to any of the ingredients listed.

WARNINGS AND PRECAUTIONS

- For external use only.
- Avoid contacts with the eyes and other mucous membranes such as mouth, nose and genital region.
- Not to be applied to abraded, blistered, inflamed, raw or oozing areas of the skin.
- Do not use on open wounds or burns.
- Discontinue use if rash, irritation, burning sensation or sensitivity develops/occurs.
- If symptoms persist, consult your pharmacist or doctor.

PREGNANCY AND LACTATION

The safety of calamine and zinc oxide during pregnancy and lactation has not been established but use during these periods is not considered to constitute a hazard.

SIDE EFFECTS

Calamine and zinc oxide is well-tolerated except for occasional dryness following and excessive use. Occasional hypersensitivity reactions such as itchiness in sensitive individuals.

SYMPTOMS AND TREATMENT OF OVERDOSE

Overdose arising from the topical application is extremely remote. As the ingredients are inert and very poorly absorbed, effects if any arising from accidental overdose (orally) is unlikely to be life threatening. If ingested, symptoms of overdose include gastritis, nausea, vomiting, epigastric discomfort and diarrhoea. Treatment consists of symptomatic measures only. Maintain adequate hydration if vomiting or diarrhea becomes extensive. Milk or antacids may be used to allay gastritis in the event of oral ingestion.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINE

Unlikely to impair a patient's ability to drive or use machinery.

STORAGE CONDITIONS

Store below 30°C, in airtight containers, away from direct light.

SHELF LIFE

Product should not be used beyond the expiry date imprinted on the product packaging.

PACKING

Plastic Container of 120mL.

REGISTRATION NO

MAL18086094XZ

KEEP MEDICINES OUT OF REACH OF CHILDREN
JAUHI UBAT-UBATAN DARI KANAK-KANAK

For further information, please consult your doctor or your pharmacist.

Revision Number : 02

Revision Date : 02 Apr 2026

LT-190N.03 PRODUCT REGISTRATION AND MANUFACTURER

INSPIRAPHARMA SDN. BHD. (200401023395)

LOT 24 & 25, JALAN 8,
KAWASAN PERUSAHAAN BAKAR ARANG,
08000 SUNGAI PETANI,
KEDAH DARUL AMAN, MALAYSIA.