

Duofilm

Solution

1. Product Name

Duofilm Solution

2. Qualitative And Quantitative Composition

Salicylic acid 16.7% w/w

Lactic acid 15.0% w/w

Excipient:

Flexible collodion (contains pyroxylin, colophony, virgin castor oil, ethanol and ether).

This product contains Ethanol 90%.

3. Product Description

A yellowish to amber coloured clear viscous liquid with a strong odour ether.

4. Pharmacodynamics

ATC Code

Pharmacotherapeutic group: Wart and anti-corn preparations.

ATC code: D11AF

Mechanism of Action

Topically applied salicylic acid is keratolytic. The keratolytic activity produces desquamation by solubilising the intercellular cement in the stratum corneum resulting in the shedding of skin scales.

Lactic acid affects the keratinisation process, reducing the hyperkeratosis which is characteristic of warts, corns and calluses. At high concentrations it can cause epidermolysis, leading to the destruction of the keratotic tissue, and in the case of warts, of the causative virus. It also has antiseptic properties.

Flexible collodion provides a viscous vehicle that allows accurate application of the active ingredients to the wart, corn or callus. It also forms a film that helps to hydrate and promote the destruction of hyperkeratotic tissue.

5. Pharmacokinetics

Absorption

Salicylic acid is absorbed through the skin; where detectable, maximum plasma levels are found 6 to 12 hours after application. Systemic absorption of salicylic acid has been reported to range from 9% to 25% after topical application of other salicylic acid-containing preparations. The extent of absorption is variable depending on the duration of contact and the vehicle. Despite percutaneous absorption, the systemic exposure is low given the low dose topically administered to small, localised areas of hyperkeratotic tissue.

Human abdominal skin in a flow-through diffusion system was used to assess the in vitro percutaneous absorption of lactic acid. At a pH of 3, the amount of radioactivity detected in the receptor fluid, stratum corneum, epidermis, and dermis was 3.6, 6.3, 6.6, and 13.9%, respectively.

Distribution

Following percutaneous absorption, salicylic acid is distributed in the extracellular space; approximately half of which is protein bound to albumin.

Metabolism

Salicylates are metabolised in the liver by microsomal enzymes to salicyluric acid and phenolic glucuronides of salicylic acid. That which is not metabolised is excreted in the urine as unchanged salicylic acid.

Elimination

Within 24 hours of salicylic acid being absorbed and distributed in the intercellular space, approximately 95% of the absorbed dose can be recovered in the urine.

6. Mode Of Administration

For topical application only.

7. Indication

Duofilm Solution is indicated in the treatment of warts, corns and calluses.

8. Recommended Dose

For topical use only.

- **Adults, the elderly and children aged 2 years and over**

Salicylic acid and lactic acid solution should be applied to the affected areas only. Use in children should be under the supervision of an adult. Salicylic acid and lactic acid solution should be applied to the wart, corn or callus once daily, preferably at bedtime.

Procedure for application:

1. The wart, corn or callus should be soaked in warm water for 5 minutes and dried thoroughly with a clean towel.
2. The surface of the wart, corn or callus should be rubbed with a nail file, pumice stone, emery board or coarse washcloth, with care taken not to cause bleeding.
3. A thin layer of salicylic acid and lactic acid solution should be applied directly to the wart, corn or callus. Care should be taken to avoid the healthy surrounding skin.
4. The solution should be allowed to dry thoroughly. The wart, corn or callus should be covered with a plaster (dressing) if it is large or if it is on the foot to help penetration of ingredients.

It is recommended that treatment continues until whichever of the following occurs first:

- corns and calluses have been treated for 2 weeks
- warts have been treated for 12 weeks
- or until the wart, corn or callus is completely cleared and the normal ridgelines of the skin have been restored.

For warts, clinically visible improvement should occur in 1-2 weeks, but the maximum effect may be expected after 4-8 weeks.

If warts persist beyond 12 weeks of treatment, the patient should be advised to consult their pharmacist or doctor. Consider alternative treatments if warts cover a large area of the body (more than 5 cm²) (see Warnings and Precautions). Patients should be advised to consult a pharmacist or doctor if skin irritation develops.

Due to the flammable nature of salicylic acid and lactic acid solution, patients should avoid smoking or being near an open flame during application and immediately after use.

- **Infants under 2 years of age**

Treatment of infants under the age of 2 years is not recommended.

- **Elderly**

No dosage adjustment is required as clinically significant systemic exposure is not expected.

- **Hepatic Impairment**

No dosage adjustment is required as clinically significant systemic exposure is not expected.

• **Renal Impairment**

No dosage adjustment is required as clinically significant systemic exposure is not expected.

9. Contraindication

Duofilm Solution is contraindicated in patients with a prior hypersensitivity reaction to salicylic acid, lactic acid or any other ingredient of the preparation.

Do not use on open wounds, irritated or reddened skin, or any area that is infected.

Do not use on moles, birthmarks, genital warts, warts on the face or mucous membranes, or warts with hair growing from them, red edges, or an unusual colour.

10. Warning and Precautions

Duofilm Solution may cause eye irritation. Avoid contact with eyes and other mucous membranes. In case of accidental contact with the eyes or other mucous membranes, flush with water for 15 minutes.

Avoid exposure to healthy skin (see Adverse Reactions). Duofilm Solution may cause skin irritation. If undue skin irritation develops treatment should be discontinued. Consider alternative treatments if warts cover a large area of the body (more than 5 cm²) due to the potential risk of salicylate toxicity.

Duofilm Solution is not recommended in patients with diabetes, circulatory problems or peripheral neuropathy except under the supervision of a doctor.

Oral salicylates taken during or immediately after a viral illness have been associated with Reye's syndrome and hence there is a theoretical risk with topical salicylates. Therefore, do not use in children or teenagers during or immediately after chickenpox, influenza, or other viral infections.

It has been reported that salicylates are excreted via breastmilk (see Pregnancy and Lactation).

Patients should be advised not to inhale the vapour. Keep out of the sight and reach of children.

11. Interactions With Other Medicaments

Topical salicylic acid and lactic acid solution may increase the absorption of other topically applied medicines. Therefore, concomitant use of Duofilm Solution and other topical medicines on the treated area should be avoided. As systemic exposure of topically applied salicylic acid and lactic acid solution is low, interaction with systemically administered medicines is not anticipated.

12. Pregnancy and Lactation

Pregnancy

The safety of salicylic acid and lactic acid solution during human pregnancy has not been established. Studies in animals given salicylic acid orally demonstrated embryo toxicity at high doses. Duofilm Solution is not recommended during pregnancy.

Lactation

Salicylates are excreted in human milk. Duofilm Solution is not recommended during lactation. If used or administered during lactation, care should be taken to avoid contact with the breast area in order to avoid accidental ingestion by the infant.

13. Side Effects

The following convention is used for the classification of the frequency of an adverse reaction and is based on the CIOMS guidelines:

- Very common: ≥1/10
- Common: ≥1/100 to <1/10
- Uncommon: ≥1/1000 to <1/100
- Rare: ≥1/10000 to <1/1000
- Very rare: <1/10000
- Not known: (Cannot be estimated from the available data)

Clinical Trial Data

Immune system disorders

Common: Rash.
Skin and subcutaneous tissue disorders
Very common: Application site reaction, pruritus, burning sensation, erythema, scaling, dryness.
Common: Skin hypertrophy.

Post Marketing Data

Immune system disorders

Rare: Application site hypersensitivity including inflammation.
Skin and subcutaneous tissue disorders
Rare: Application site pain and irritation, application site discoloration/skin discoloration, exposure to healthy skin can lead to application site blistering and skin exfoliation (see Warnings and Precautions), allergic dermatitis.

14. Symptoms And Treatment of Overdose

Symptoms and Signs

In the event of accidental oral ingestion symptoms of salicylate toxicity may occur. The risk of developing symptoms of salicylate poisoning or salicylism is increased if topical salicylic acid and lactic acid solution is used excessively or if it is used for prolonged periods of time. Therefore, duration of use and recommended frequency compliance is very important.

Treatment

Management should be as clinically indicated or as recommended by the national poisons centre, where available. There is no specific treatment for accidental oral ingestion of salicylic acid and lactic acid solution. If accidental oral ingestion occurs, the patient should be treated according to local guidelines with appropriate monitoring as necessary.

15. Storage Condition

Contents are flammable.
Keep away from fire or flame.
Keep bottle tightly closed when not in use and store at room temperature away from heat.
Do not store above 30°C.

16. Dosage Form and Packaging Available

Solution
15ml amber glass bottle

17. Manufacturer

Delpharm Bladel B. V.
Industrieweg 1,
BLADEL, 5531AD,
Netherlands

18. Product Registration Holder

DKSH Malaysia Sdn Bhd
B-11-01, The Ascent, Paradigm, No.1, Jalan SS7/26A, Kelana Jaya, 47301, Petaling Jaya, Selangor, Malaysia

Date of Revision: 31 May 2021



Produkt/Product: DUOFILM SOLUTION MY:PIL
STADA Mat.-Nr./Mat.-No.: Mockup
STADA Art.-Nr./Art.-No.: 172943
Hersteller/Manufacturer: Delpharm Bladel BV
Hersteller Art.-Nr./ Manufacturer Art.-No.: -
Format/Size: -
Schriftgröße/Font-size: 7.5 pt - 14 pt
PZN/EAN: -
Pharma-Code: -
Flattermarke/Collating mark: -
1. Farbe/Colour: Black
2. Farbe/Colour: -
3. Farbe/Colour: -
4. Farbe/Colour: -
5. Farbe/Colour: -
6. Farbe/Colour: -
7. Farbe/Colour: -
8. Farbe/Colour: -
Version: 1. Loop Datum/Date: 03.09.2021
Agentur/Agency: STADA Agency
Grafik erstellt von/Graphic made by: N.V.
Aktueller Umlauf/Current loop: N.V.
Genehmigt/Approved