
DULOX 30 (Duloxetine Hydrochloride Delayed Release Capsules 30mg)
DULOX 60 (Duloxetine Hydrochloride Delayed Release Capsules 60mg)

NAME OF THE PRODUCT

DULOX 30 (Duloxetine Hydrochloride Delayed Release Capsules 30mg)

DULOX 60 (Duloxetine Hydrochloride Delayed Release Capsules 60mg)

NAME AND STRENGTH OF ACTIVE INGREDIENT

Name: Duloxetine Hydrochloride Capsules

Strength: 30 mg

Name: Duloxetine Hydrochloride Capsules

Strength: 60 mg

PRODUCT DESCRIPTION

DULOX 30 (Duloxetine Hydrochloride Delayed Release Capsules 30mg)

Opaque blue cap / Opaque white body size '3' hard gelatin capsules imprinted with 'H' on cap and '191' on body with golden yellow ink, filled with white to off white colored pellets.

DULOX 60 (Duloxetine Hydrochloride Delayed Release Capsules 60mg)

Opaque blue cap / Opaque green body size '1' hard gelatin capsules imprinted with 'H' on cap and '192' on body with white ink, filled with white to off white colored pellets.

PHARMACODYNAMICS

Pharmacotherapeutic group: Other antidepressants.

ATC code: N06AX21.

Mechanism of action

Duloxetine is a combined serotonin (5-HT) and noradrenaline (NA) reuptake inhibitor. It weakly inhibits dopamine reuptake, with no significant affinity for histaminergic, dopaminergic, cholinergic, and adrenergic receptors. It was reported that duloxetine dose-dependently increases extracellular levels of serotonin and noradrenaline in various brain areas.

Pharmacodynamic effects

Duloxetine normalised pain thresholds in several models of neuropathic and inflammatory pain and attenuated pain behaviour in a model of persistent pain. The pain inhibitory action of duloxetine is believed to be a result of potentiation of descending inhibitory pain pathways within the central nervous system.

PHARMACOKINETICS

Duloxetine is administered as a single enantiomer. Duloxetine is extensively metabolised by oxidative enzymes (CYP1A2 and the polymorphic CYP2D6), followed by conjugation. The pharmacokinetics of duloxetine demonstrate large intersubject variability (generally 50-60%), partly due to gender, age, smoking status, and CYP2D6 metaboliser status.

Absorption: Duloxetine is well absorbed after oral administration, with a C_{max} occurring 6 hours post-dose. The absolute oral bioavailability of duloxetine ranged from 32% to 80% (mean of 50%). Food delays the time to reach the peak concentration from 6 to 10 hours and it marginally decreases the extent of absorption (approximately 11%). These changes do not have any clinical significance.

Distribution: Duloxetine is approximately 96% bound to human plasma proteins. Duloxetine binds to both albumin and alpha1-acid glycoprotein. Protein binding is not affected by renal or hepatic impairment.

Biotransformation: Duloxetine is extensively metabolised and the metabolites are excreted principally in urine. Both cytochromes P450-2D6 and 1A2 catalyse the formation of the two major metabolites, glucuronide conjugate of 4-hydroxy duloxetine and sulfate conjugate of 5-hydroxy, 6-methoxy duloxetine. The pharmacokinetics of duloxetine in patients who are poor metabolisers with respect to CYP2D6 has not been specifically investigated. Limited data suggest that the plasma levels of duloxetine

DULOX 30 (Duloxetine Hydrochloride Delayed Release Capsules 30mg)
DULOX 60 (Duloxetine Hydrochloride Delayed Release Capsules 60mg)

are higher in these patients.

Elimination: The elimination half-life of duloxetine ranges from 8 to 17 hours (mean of 12 hours). After an intravenous dose the plasma clearance of duloxetine ranges from 22 l/hr to 46 l/hr (mean of 36 l/hr). After an oral dose the apparent plasma clearance of duloxetine ranges from 33 to 261 l/hr (mean 101 l/hr).

Special Populations

Gender: Pharmacokinetic differences have been identified between males and females (apparent plasma clearance is approximately 50% lower in females). Based upon the overlap in the range of clearance, gender-based pharmacokinetic differences do not justify the recommendation for using a lower dose for female patients.

Age: Pharmacokinetic differences have been identified between younger and elderly females (≥ 65 years) (AUC increases by about 25% and half-life is about 25% longer in the elderly), although the magnitude of these changes is not sufficient to justify adjustments to the dose. As a general recommendation, caution should be exercised when treating the elderly.

Renal impairment: End stage renal disease (ESRD) patients receiving dialysis had 2-fold higher duloxetine C_{max} and AUC values. Pharmacokinetic data on duloxetine is limited in patients with mild or moderate renal impairment.

Hepatic impairment: Moderate liver disease (Child-Pugh Class B) affected the pharmacokinetics of duloxetine. The apparent plasma clearance of duloxetine was 79% lower, the apparent terminal half-life was 2.3-times longer, and the AUC was 3.7-times higher in patients with moderate liver disease. The pharmacokinetics of duloxetine and its metabolites have not been studied in patients with mild or severe hepatic insufficiency.

Breast-feeding mothers: Duloxetine is detected in breast milk, and steady-state concentrations in breast milk are about one-fourth those in plasma. The amount of duloxetine in breast milk is approximately 7 µg/day while on 40 mg twice-daily dosing. Lactation did not influence duloxetine pharmacokinetics.

Paediatric population: It is reported that duloxetine steady-state plasma concentrations in paediatric patients were mostly within the concentration range observed in adult patients.

INDICATIONS

Treatment of major depressive disorder. Management of neuropathic pain associated with diabetic peripheral neuropathy in adults. Treatment of generalized anxiety disorder.

RECOMMENDED DOSE

For oral use.

Major Depressive Disorder

The starting and recommended maintenance dose is 60 mg once daily with or without food. Dosages above 60 mg once daily, up to a maximum dose of 120 mg per day administered in evenly divided doses, have been evaluated from a safety perspective. However, there is no clinical evidence suggesting that patients not responding to the initial recommended dose may benefit from dose up-titrations. Therapeutic response is usually seen after 2-4 weeks of treatment. After consolidation of the antidepressive response, it is recommended to continue treatment for several months, in order to avoid relapse.

Diabetic Peripheral Neuropathic Pain

Duloxetine should be administered at a dose of 60mg once daily with or without food. Dosages above 60mg once daily, up to a maximum dose of 120mg per day administered in evenly divided doses, have been evaluated from a safety perspective. The plasma concentration of duloxetine displays large inter-individual variability. Hence, some patients that respond insufficiently to 60mg may benefit from a higher dose. Response to treatment should be evaluated after 2 months. In patients with inadequate initial

DULOX 30 (Duloxetine Hydrochloride Delayed Release Capsules 30mg)
DULOX 60 (Duloxetine Hydrochloride Delayed Release Capsules 60mg)

response, additional response after this time is unlikely. The therapeutic benefit should be reassessed regularly (at least every three months.)

Generalized Anxiety Disorder

Duloxetine should be administered at a dose of 60 mg once daily without regard to meals. For some patients, it may be desirable to start at 30mg once daily for one week to allow patients to adjust to medication before increasing to 60mg once daily. While a 120mg once daily was shown to be effective, there is no evidence that doses greater than 60mg/day confer additional benefit. Nevertheless, if a decision is made to increase the dose beyond 60mg once daily, dose increases should be in increments of 30mg once daily. The safety of doses above 120 mg once daily has not been adequately evaluated.

Special Populations and General Information:

Hepatic impairment

Duloxetine should not be used in patients with liver disease resulting in hepatic impairment.

Renal insufficiency

No dosage adjustment is necessary for patients with mild or moderate renal dysfunction (creatinine clearance 30 to 80 ml/min).

Paediatric population

Duloxetine should not be used in children and adolescents under the age of 18 years for the treatment of major depressive disorder because of safety and efficacy concerns. The safety and efficacy of duloxetine for the treatment of diabetic peripheral neuropathic pain or generalized anxiety disorder have not been studied. No data are available.

Elderly Major Depressive Disorder:

Depression may have different features in the elderly which make it difficult to extrapolate efficacy and safety data from a younger population. Only limited information on the use of Duloxetine in elderly patients with major depressive disorders is available. Therefore, caution should be exercised when treating the elderly. Until more efficacy data are available the use of Duloxetine in the very elderly population (>75 years) is not recommended.

Diabetic Peripheral Neuropathic Pain:

No dosage adjustment is recommended for elderly patients solely on the basis of age. However, caution should be exercised when treating the elderly.

Discontinuation of treatment

Abrupt discontinuation should be avoided. When stopping treatment with Duloxetine the dose should be gradually reduced over a period of at least one to two weeks in order to reduce the risk of withdrawal reactions. If intolerable symptoms occur following a decrease in the dose or upon discontinuation of treatment, then resuming the previously prescribed dose may be considered.

Subsequently, the physician may continue decreasing the dose, but at a more gradual rate.

MODE OF ADMINISTRATION

Oral

CONTRAINDICATIONS

Hypersensitivity to the active substance or to any of the excipients listed.

Concomitant use of Duloxetine with non-selective, irreversible monoamine oxidase inhibitors (MAOIs) is contraindicated. Liver disease resulting in hepatic impairment.

Duloxetine should not be used in combination with fluvoxamine, ciprofloxacin or enoxacin (i.e., potent CYP1A2 inhibitors), since the combination results in elevated plasma concentrations of duloxetine.

Severe renal impairment (creatinine clearance <30 ml/min).

The initiation of treatment with Duloxetine is contraindicated in patients with uncontrolled hypertension

DULOX 30 (Duloxetine Hydrochloride Delayed Release Capsules 30mg)
DULOX 60 (Duloxetine Hydrochloride Delayed Release Capsules 60mg)

that could expose patients to a potential risk of hypertensive crisis.

WARNINGS AND PRECAUTIONS

Mania and Seizures

Duloxetine should be used with caution in patients with a history of mania or a diagnosis of bipolar disorder, and/or seizures.

Mydriasis

Mydriasis has been reported in association with duloxetine, therefore, caution should be used when prescribing duloxetine to patients with increased intraocular pressure or those at risk of acute narrow-angle glaucoma.

Blood Pressure and Heart Rate

Duloxetine has been associated with an increase in blood pressure, and clinically significant hypertension in some patients. This may be due to the noradrenergic effect of duloxetine. Cases of hypertensive crisis have been reported with duloxetine, especially in patients with pre-existing hypertension. Therefore, in patients with known hypertension and/or other cardiac disease, blood pressure monitoring is recommended, especially during the first month of treatment. Duloxetine should be used with caution in patients whose conditions could be compromised by an increased heart rate or by an increase in blood pressure. Caution should also be exercised when duloxetine is used with medicinal products that may impair its metabolism. For patients who experience a sustained increase in blood pressure while receiving duloxetine, either dose reduction or gradual discontinuation should be considered. In patients with uncontrolled hypertension, duloxetine should not be initiated.

Renal Impairment

Increased plasma concentrations of duloxetine occur in patients with severe renal impairment on haemodialysis (creatinine clearance <30 ml/min).

Serotonin syndrome

As with other serotonergic agents, serotonin syndrome, a potentially life-threatening condition, may occur with duloxetine treatment, particularly with concomitant use of other serotonergic agents (including SSRIs, SNRIs, tricyclic antidepressants or triptans), with agents that impair metabolism of serotonin such as MAOIs, or with antipsychotics or other dopamine antagonists that may affect the serotonergic neurotransmitter systems.

Serotonin syndrome symptoms may include mental status changes (e.g., agitation, hallucinations, coma), autonomic instability (e.g., tachycardia, labile blood pressure, hyperthermia), neuromuscular aberrations (e.g., hyperreflexia, incoordination) and/or gastrointestinal symptoms (e.g., nausea, vomiting, diarrhoea). If concomitant treatment with duloxetine and other serotonergic agents that may affect the serotonergic and/or dopaminergic neurotransmitter systems is clinically warranted, careful observation of the patient is advised, particularly during treatment initiation and dose increases.

St John's Wort
Adverse reactions may be more common during concomitant use of duloxetine and herbal preparations containing St John's Wort (*Hypericum perforatum*).

Suicide

Major Depressive Disorder and Generalised Anxiety Disorder: Depression is associated with an increased risk of suicidal thoughts, self-harm, and suicide (suicide-related events). This risk persists until significant remission occurs. As improvement may not occur during the first few weeks or more of treatment, patients should be closely monitored until such improvement occurs. It is general clinical experience that the risk of suicide may increase in the early stages of recovery.

Other psychiatric conditions for which duloxetine is prescribed can also be associated with an increased risk of suicide-related events. In addition, these conditions may be co-morbid with major depressive disorder. The same precautions observed when treating patients with major depressive disorder should therefore be observed when treating patients with other psychiatric disorders.

DULOX 30 (Duloxetine Hydrochloride Delayed Release Capsules 30mg)
DULOX 60 (Duloxetine Hydrochloride Delayed Release Capsules 60mg)

Patients with a history of suicide-related events or those exhibiting a significant degree of suicidal thoughts prior to commencement of treatment, are known to be at greater risk of suicidal thoughts or suicidal behaviour, and should receive careful monitoring during treatment.

Cases of suicidal thoughts and suicidal behaviours have been reported during duloxetine therapy or early after treatment discontinuation.

Close supervision of patients, and in particular those at high risk, should accompany medicinal product therapy, especially in early treatment and following dose changes. Patients (and caregivers of patients) should be alerted about the need to monitor for any clinical worsening, suicidal behaviour or thoughts, and unusual changes in behaviour, and to seek medical advice immediately if these symptoms present.

Diabetic Peripheral Neuropathic Pain

As with other medicinal products with similar pharmacological action (antidepressants), isolated cases of suicidal ideation and suicidal behaviours have been reported during duloxetine therapy or early after treatment discontinuation. Physicians should encourage patients to report any distressing thoughts or feelings at any time.

Use in Children and Adolescents Under 18 Years of Age

Duloxetine should not be used in the treatment of children and adolescents under the age of 18 years. Suicide-related behaviours (suicide attempts and suicidal thoughts) and hostility (predominantly aggression, oppositional behaviour, and anger) were more frequently observed among children and adolescents treated with antidepressants. If, based on clinical need, a decision to treat is nevertheless taken, the patient should be carefully monitored for the appearance of suicidal symptoms. In addition, long-term safety data in children and adolescents concerning growth, maturation, and cognitive and behavioural development are lacking.

Haemorrhage

There have been reports of bleeding abnormalities, such as ecchymoses, purpura, and gastrointestinal haemorrhage, with selective serotonin reuptake inhibitors (SSRIs) and serotonin/noradrenaline reuptake inhibitors (SNRIs), including duloxetine. Caution is advised in patients taking anticoagulants and/or medicinal products known to affect platelet function (e.g., NSAIDs or acetylsalicylic acid (ASA)), and in patients with known bleeding tendencies.

Hyponatraemia

Hyponatraemia has been reported when administering duloxetine, including cases with serum sodium lower than 110 mmol/l. Hyponatraemia may be due to a syndrome of inappropriate anti-diuretic hormone secretion (SIADH). The majority of cases of hyponatraemia were reported in the elderly, especially when coupled with a recent history of, or condition pre-disposing to, altered fluid balance. Caution is required in patients at increased risk for hyponatraemia, such as elderly, cirrhotic, or dehydrated patients, or patients treated with diuretics.

Discontinuation of Treatment

Withdrawal symptoms when treatment is discontinued are common, particularly if discontinuation is abrupt. The risk of withdrawal symptoms seen with SSRIs and SNRIs may be dependent on several factors, including the duration and dose of therapy and the rate of dose reduction. Generally, these symptoms are mild to moderate; however, in some patients they may be severe in intensity. They usually occur within the first few days of discontinuing treatment, but there have been very rare reports of such symptoms in patients who have inadvertently missed a dose. Generally, these symptoms are self-limiting and usually resolve within 2 weeks, though in some individuals they may be prolonged (2-3 months or more). It is therefore advised that duloxetine should be gradually tapered when discontinuing treatment over a period of no less than 2 weeks, according to the patient's needs.

Elderly

Caution should be exercised when treating the elderly with the maximum dosage. Akathisia/Psychomotor

DULOX 30 (Duloxetine Hydrochloride Delayed Release Capsules 30mg)
DULOX 60 (Duloxetine Hydrochloride Delayed Release Capsules 60mg)

Restlessness

The use of duloxetine has been associated with the development of akathisia, characterised by a subjectively unpleasant or distressing restlessness and need to move, often accompanied by an inability to sit or stand still. This is most likely to occur within the first few weeks of treatment. In patients who develop these symptoms, increasing the dose may be detrimental.

Medicinal Products Containing Duloxetine

Duloxetine is used under different trademarks in several indications (treatment of diabetic neuropathic pain, major depressive disorder, generalised anxiety disorder and stress urinary incontinence). The use of more than one of these products concomitantly should be avoided.

Hepatitis/Increased Liver Enzymes

Cases of liver injury, including severe elevations of liver enzymes (>10-times upper limit of normal), hepatitis, and jaundice have been reported with duloxetine. Most of them occurred during the first months of treatment. The pattern of liver damage was predominantly hepatocellular. Duloxetine should be used with caution in patients treated with other medicinal products associated with hepatic injury.

Sucrose

Duloxetine hard gastro-resistant capsules contain sucrose. Patients with rare hereditary problems of fructose intolerance, glucose-galactose malabsorption or sucrose-isomaltase insufficiency should not take this medicine.

Effects on ability to drive and use machines

No studies on the effects on the ability to drive and use machines have been performed. Duloxetine may be associated with sedation and dizziness. Patients should be instructed that if they experience sedation or dizziness they should avoid potentially hazardous tasks such as driving or operating machinery.

Suicidality in Children and Adolescents

Antidepressants increase the risk of suicidal thinking and behavior (suicidality) in children and adolescents with major depressive disorder (MDD) and other psychiatric disorders.

Anyone considering the use of an antidepressant in a child or adolescent for any clinical use must balance the risk of increased suicidality with the clinical need.

Patients who are started on therapy should be observed closely for clinical worsening, suicidality, or unusual changes in behavior. Families and caregivers should be advised to closely observe the patient and to communicate with the prescriber. The indication(s) approved in paediatric for the particular drug should be clearly stated / included.

INTERACTION WITH OTHER MEDICAMENTS

Monoamine Oxidase Inhibitors (MAOIs)

Due to the risk of serotonin syndrome, duloxetine should not be used in combination with non- selective, irreversible monoamine oxidase inhibitors (MAOIs) or within at least 14 days of discontinuing treatment with an MAOI. Based on the half-life of duloxetine, at least 5 days should be allowed after stopping duloxetine before starting an MAOI.

The concomitant use of duloxetine with selective, reversible MAOIs, like moclobemide, is not recommended. The antibiotic linezolid is a reversible non- selective MAOI and should not be given to patients treated with duloxetine.

Inhibitors of CYP1A2

Because CYP1A2 is involved in duloxetine metabolism, concomitant use of duloxetine with potent inhibitors of CYP1A2 is likely to result in higher concentrations of duloxetine. Therefore, duloxetine should not be administered in combination with potent inhibitors of CYP1A2 like fluvoxamine.

CNS Medicinal Products

Caution is advised when duloxetine is taken in combination with other centrally-acting medicinal products or substances, including alcohol and sedative medicinal products (e.g., benzodiazepines,

DULOX 30 (Duloxetine Hydrochloride Delayed Release Capsules 30mg)
DULOX 60 (Duloxetine Hydrochloride Delayed Release Capsules 60mg)

morphinomimetics, antipsychotics, phenobarbital, sedative antihistamines).

Serotonergic agents: In rare cases, serotonin syndrome has been reported in patients using SSRIs/SNRIs concomitantly with serotonergic agents. Caution is advisable if duloxetine is used concomitantly with serotonergic agents like SSRIs, SNRIs, tricyclic antidepressants like clomipramine or amitriptyline, MAOIs like moclobemide or linezolid, St John's Wort (*Hypericum perforatum*) or triptans, tramadol, pethidine, and tryptophan.

Effect of Duloxetine on Other Medicinal Products

Medicinal products metabolised by CYP1A2: The pharmacokinetics of theophylline, a CYP1A2 substrate, were not significantly affected by co-administration with duloxetine (60 mg twice daily).

Medicinal products metabolised by CYP2D6: Duloxetine is a moderate inhibitor of CYP2D6. Caution is advised if duloxetine is co-administered with medicinal products that are predominantly metabolised by CYP2D6 (risperidone, tricyclic antidepressants [TCAs], such as nortriptyline, amitriptyline, and imipramine), particularly if they have a narrow therapeutic index (such as flecainide, propafenone, and metoprolol).

Oral contraceptives and other steroidal agents: Reported studies demonstrate that duloxetine does not induce the catalytic activity of CYP3A. Specific drug interaction studies have not been performed.

Anticoagulants and antiplatelet agents: Caution should be exercised when duloxetine is combined with oral anticoagulants or antiplatelet agents due to a potential increased risk of bleeding attributable to a pharmacodynamic interaction. Furthermore, increases in INR values have been reported when duloxetine was co-administered to patients treated with warfarin.

Effects of Other Medicinal Products on Duloxetine

Antacids and H₂ antagonists: Co-administration of duloxetine with aluminium- and magnesium-containing antacids, or duloxetine with famotidine, had no significant effect on the rate or extent of duloxetine absorption after administration of a 40 mg oral dose.

Inducers of CYP1A2: Population pharmacokinetic analyses have shown that smokers have almost 50% lower plasma concentrations of duloxetine compared with non-smokers.

PREGNANCY AND LACTATION

Pregnancy

There are no adequate data on the use of duloxetine in pregnant women. The potential risk for humans is unknown. Epidemiological data have suggested that the use of SSRIs in pregnancy, particularly in late pregnancy, may increase the risk of persistent pulmonary hypertension in the newborn (PPHN). Although no studies have investigated the association of PPHN to SNRI treatment, this potential risk cannot be ruled out with duloxetine, taking into account the related mechanism of action (inhibition of the re-uptake of serotonin).

As with other serotonergic medicinal products, discontinuation symptoms may occur in the neonate after maternal duloxetine use near term. Discontinuation symptoms seen with duloxetine may include hypotonia, tremor, jitteriness, feeding difficulty, respiratory distress and seizures. The majority of cases have occurred either at birth or within a few days of birth.

Duloxetine should be used in pregnancy only if the potential benefit justifies the potential risk to the foetus. Women should be advised to notify their physician if they become pregnant, or intend to become pregnant, during therapy.

Breast-Feeding

Duloxetine is very weakly excreted into human milk. The estimated daily infant dose on a mg/kg basis is approximately 0.14% of the maternal dose. As the safety of duloxetine in infants is not known, the use of duloxetine while breast-feeding is not recommended.

DULOX 30 (Duloxetine Hydrochloride Delayed Release Capsules 30mg)
DULOX 60 (Duloxetine Hydrochloride Delayed Release Capsules 60mg)

SIDE EFFECTS

A. Summary of the safety profile

The most commonly reported adverse reactions in patients treated with duloxetine were nausea, headache, dry mouth, somnolence and dizziness. However, the majority of common adverse reactions were mild to moderate; they usually started early in therapy, and most tended to subside even as therapy was continued.

B. Tabulated summary of adverse reactions

Table 1 gives the adverse reactions observed.

Table 1: Adverse reactions

Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

Very common	Common	Uncommon	Rare	Very rare
<i>Infections and infestations</i>				
		Laryngitis		
<i>Immune system disorders</i>				
			Anaphylactic reaction; Hypersensitivity disorder	
<i>Endocrine disorders</i>				
			Hyperthyroidism	
<i>Metabolism and nutrition disorders</i>				
	Decreased appetite	Hyperglycemia (reported especially in diabetic patients)	Dehydration; Hyponatremia; SIADH ⁶	
<i>Psychiatric disorders</i>				
	Insomnia; Agitation; Libido decreased; Anxiety; Orgasm abnormal; Abnormal dreams	Suicidal ideation; ^{5,7} Sleep disorders; Bruxism; Disorientation Apathy	Suicidal behaviour; ^{5,7} Mania; Hallucinations; Aggression and anger ⁴	
<i>Nervous system disorders</i>				
Headache; Somnolence	Dizziness; Lethargy; Tremor; Paraesthesia	Myoclonus; Akathisia ⁷ Nervousness; Disturbance in attention; Dysgeusia; Dyskinesia; Restless legs syndrome; Poor quality sleep	Serotonin syndrome; ⁶ Convulsion; ¹ Psychomotor restlessness; ⁶ Extrapyramidal symptoms ⁶	
<i>Eye disorders</i>				
	Blurred vision	Mydriasis; Visual impairment	Glaucoma	
<i>Ear and labyrinth disorders</i>				
	Tinnitus ¹	Vertigo; Ear pain		
<i>Cardiac disorders</i>				
	Palpitations	Tachycardia; Supraventricular arrhythmia, mainly atrial fibrillation		

DULOX 30 (Duloxetine Hydrochloride Delayed Release Capsules 30mg)
DULOX 60 (Duloxetine Hydrochloride Delayed Release Capsules 60mg)

Very common	Common	Uncommon	Rare	Very rare
<i>Vascular disorders</i>				
	Blood pressure increase; ³ Flushing	Syncope; ² Hypertension; ^{3,7} Orthostatic hypotension; ² Peripheral coldness	Hypertensive crisis ^{3, 6}	
<i>Respiratory, thoracic and mediastinal disorders</i>				
	Yawning	Throat tightness; Epistaxis		
<i>Gastrointestinal disorders</i>				
Nausea; Dry mouth	Constipation; Diarrhea; Abdominal pain; Vomiting; Dyspepsia; Flatulence	Gastrointestinal haemorrhage; ⁷ Gastroenteritis; Eructation; Gastritis; Dyspepsia	Gastrointestinal haemorrhage; ⁷ Gastroenteritis; Eructation; Gastritis; Dysphagia	
<i>Hepato-biliary disorders</i>				
		Hepatitis; ³ Elevated liver enzymes (ALT, AST, alkaline phosphatase); Acute liver injury	Hepatic failure; ⁶ Jaundice ⁶	
<i>Skin and subcutaneous tissue disorders</i>				
	Sweating increased; Rash	Night sweats; Urticaria; Dermatitis contact; Cold sweat; Photosensitivity reactions; Increased tendency to bruise	Steven-Johnson Syndrome; ⁶ Angioneurotic oedema ⁶	Cutaneous vasculitis
<i>Musculoskeletal and connective tissue disorders</i>				
	Musculoskeletal pain; Muscle spasm	Muscle tightness; Muscle twitching	Trismus	
<i>Renal and urinary disorders</i>				
	Dysuria; Pollakiuria	Urinary retention; Urinary hesitation; Nocturia; Polyuria; Urine flow decreased	Urine color abnormal	
<i>Reproductive system and breast disorders</i>				
	Erectile dysfunction; Ejaculation disorder; Ejaculation delayed	Gynecological hemorrhage; Menstrual disorder; Sexual dysfunction; Testicular pain	Menopausal symptoms; Galactorrhoea; Hyperprolactinaemia	
<i>General disorders and administration site conditions</i>				
	Falls; ⁸ Fatigue	Chest pain; ⁷ Feeling abnormal; Feeling cold; Thirst; Chills; Malaise;		

DULOX 30 (Duloxetine Hydrochloride Delayed Release Capsules 30mg)
DULOX 60 (Duloxetine Hydrochloride Delayed Release Capsules 60mg)

Very common	Common	Uncommon	Rare	Very rare
		Feeling hot; Gait disturbance		
<i>Investigations</i>				
	Weight decrease	Weight increase; Blood creatine phosphokinase increased; Blood potassium increased	Blood cholesterol increased	

- 1 Cases of convulsion and cases of tinnitus have also been reported after treatment discontinuation.
- 2 Cases of orthostatic hypotension and syncope have been reported especially at the initiation of treatment.
- 3 See Warnings and Precautions.
- 4 Cases of aggression and anger have been reported particularly early in treatment or after treatment discontinuation.
- 5 Cases of suicidal ideation and suicidal behaviours have been reported during duloxetine therapy or early after treatment discontinuation.
- 6 Estimated frequency of post-marketing surveillance reported adverse reactions; not observed in placebo-controlled clinical trials.
- 7 Not statistically significantly different from placebo.
- 8 Falls were more common in the elderly (≥ 65 years old).
- 9 Estimated frequency based on all clinical trial data.

C. Description of selected adverse reactions

Discontinuation of duloxetine (particularly when abrupt) commonly leads to withdrawal symptoms. Dizziness, sensory disturbances (including paraesthesia or electric shock-like sensations, particularly in the head), sleep disturbances (including insomnia and intense dreams), fatigue, somnolence, agitation or anxiety, nausea and/or vomiting, tremor, headache, myalgia, irritability, diarrhoea, hyperhidrosis and vertigo are the most commonly reported reactions. Generally, for SSRIs and SNRIs, these events are mild to moderate and self-limiting; however, in some patients they may be severe and/or prolonged. It is therefore advised that when duloxetine treatment is no longer required, gradual discontinuation by dose tapering should be carried out.

D. Paediatric population

In general, the adverse reaction profile of duloxetine in children and adolescents was similar to that seen for adults.

SYMPTOMS AND TREATMENT OF OVERDOSE

Cases of overdoses, alone or in combination with other medicinal products, with duloxetine doses of 5400 mg were reported. Some fatalities have occurred, primarily with mixed overdoses, but also with duloxetine alone at a dose of approximately 1000 mg.

Signs and symptoms of overdose (duloxetine alone or in combination with other medicinal products) included somnolence, coma, serotonin syndrome, seizures, vomiting and tachycardia. No specific antidote is known for duloxetine but if serotonin syndrome ensues, specific treatment (eg, with cyproheptadine and/or temperature control) may be considered.

Monitoring of cardiac and vital signs is recommended, along with appropriate symptomatic and supportive measures. Gastric lavage may be indicated if performed soon after ingestion or in symptomatic patients. Activated charcoal may be useful in limiting absorption. Duloxetine has a large volume of distribution and forced diuresis, haemoperfusion and exchange perfusion are unlikely to be beneficial.

STORAGE CONDITION

Store below 30°C and protect from moisture

DULOX 30 (Duloxetine Hydrochloride Delayed Release Capsules 30mg)
DULOX 60 (Duloxetine Hydrochloride Delayed Release Capsules 60mg)

DOSAGE AND PACKAGING AVAILABLE

DULOX 30: 3 x 10's, 4 x 7's and 10 x 7's Alu-PVC/Alu Blister pack

DULOX 60: 3 x 10's and 10 x 10's Alu-PVC/Alu Blister pack

PRODUCT REGISTRATION HOLDER

Camber Laboratories Sdn Bhd
Unit E-13A-02, Menara SUEZCAP 2,
No.2, KL Gateway, Jalan Kerinchi,
Gerbang Kerinchi Lestari,
59200 Kuala Lumpur, Malaysia

MANUFACTURED BY

HETERO LABS LIMITED

Unit-III, 22-110, I.D.A., Jeedimetla, Hyderabad - 500 055, Telangana, INDIA.

DATE OF REVISION

October 2024