

Veterinary Package

Insert

TP-BROMPIRIN 40MG/ML SOLUTION

PRODUCT DESCRIPTION

Slightly yellow to brown liquid, which contains bromhexine hydrochloride 40mg (equivalent to 36.5mg of bromhexine) and propylene glycol 312mg (as preservative) in 1ml of solution.

PHARMACODYNAMICS & PHARMACOKINETICS

Bromhexine decreases viscosity and causes liquefaction of respiratory epithelial secretions by producing hydrolysis and dissolution of acidic mucopolysaccharide (MPSA) fibers. In this way, secretion is easily eliminated. It also has a direct expectorant action.

Following oral administration plasma concentrations are obtained at about 30 minutes in all species. It is metabolized rapidly by giving an active metabolite ambroxol. The main route of excretion of bromhexine and its metabolites is the urinary route.

INDICATION

Diseases of the respiratory tract that occur with an increase in the production of mucus and / or its viscosity.

TARGET SPECIES

Swine

RECOMMENDED DOSAGE AND ADMINISTRATION

Swine:

Administration in drinking water

From 5 - 25 kg: 0.014 - 0.027 ml of product/kg/day (equivalent to 0.5 - 1 mg of bromhexine/kg/day) for 5 consecutive days.

From 25 - 100 kg: 0.014 ml of product/kg/day (equivalent to 0.5 mg of bromhexine/kg/day) for 5 consecutive days.

Due to the form of administration and the water consumption depends on the clinical situation of the animal and the time of year, to ensure a correct dosage, the concentration of the drug will be adjusted taking into account the daily water consumption.

According to the recommended dose, the number of animals that should receive the treatment, the daily dose should be calculated by applying the following formula:

$$\begin{array}{lcl} \text{ml of product/} & = & \frac{\text{dose of active substance (mg/kg /day)} \\ \text{liter of drinking} & & \times \\ \text{water} & & \text{average weight of animals (kg)} \\ & & \text{mg of active substance in ml of} \\ & & \text{product (mg/ml)} \\ & & \times \\ & & \text{average water consumption} \\ & & \text{(liters/day)} \end{array}$$

CONTRAINDICATIONS

Do not use in animals with pulmonary edema.

Do not use in case of known hypersensitivity to the active substance and / or to any excipient.

WARNINGS & PRECAUTIONS.

Special precautions for use in animals

In respiratory diseases of bacterial origin, administer in combination with the appropriate anti-infective treatment.

In case of verminous bronchitis, use it 3 days after the use of an anthelmintic.

Specific precautions to be taken by the person administering the veterinary medicine to the animals

People with known hypersensitivity to bromhexin or to any excipient should avoid all contact with the veterinary medication.

Administer the veterinary medicinal product with caution and use gloves during their incorporation into drinking water to avoid contact with the skin.

In case of accidental contact with the eyes, wash with plenty of water.

Wash hands and exposed skin after administration of the veterinary medicinal product.

INTERACTIONS WITH OTHER MEDICATIONS

None known.

PREGNANCY AND LACTATION

It can be used during pregnancy.

ADVERSE EFFECTS

Data not available.

OVERDOSE & TREATMENT

Bromhexine has a wide margin of safety.

WITHDRAWAL PERIOD

Swine: Meat: 2 days

MAXIMUM RESIDUAL LIMIT (MRL)

Data not available.

STORAGE CONDITION

Store below 30°C. Protect from direct sunlight.

SHELF LIFE

2 years.

After 1st opening, reconstitution or dilution, use within 24 hours.

PACKING

1 L

THYE PHARMA SDN. BHD.

(manufacturer & product registration holder)

No.2 Lorong Industri Ringan Permatang

Tinggi 8,

Kawasan Industri Ringan Permatang Tinggi,

14000 Bukit Mertajam,

Pulau Pinang.

Revised September 2018