

ATC VITAMIN B₁₂



ATC VITAMIN B₁₂ INJECTION 1000 mcg/ml (10 ml)

Name and Strength of Active Substance(s)

Cyanocobalamin 1000 mcg/ml

Product Description

ATC Vitamin B₁₂ Injection 1000 mcg/ml is a red clear solution contained in a clear glass vial.

Pharmacodynamics

Vitamin B₁₂ is water-soluble. Vitamin B₁₂ acts as a coenzyme to various metabolic functions, including fat and carbohydrate metabolism and protein synthesis. It is necessary for growth, cell replication, hematopoiesis, and nucleoprotein and myelin synthesis, largely due to its effects on the metabolism of methionine, folic acid and malonic acid.

Pharmacokinetics

Absorption: Vitamin B₁₂ is readily absorbed from intramuscular and subcutaneous site of injection. Plasma level compound reaches within 1 hour after intramuscular injection. Absorbed cyanocobalamin is transported via specific B₁₂ binding proteins, transcobalamin I and transcobalamin II to the various tissue.

Distribution: Vitamin B₁₂ is distributed into the liver.

Metabolism: Vitamin B₁₂ is metabolised in the liver.

Excretion: Vitamin B₁₂ is excreted in the bile. Excess beyond daily needs is excreted, largely unchanged in the urine.

Indication

ATC Vitamin B₁₂ Injection 1000 mcg/ml is indicated for prophylaxis of anaemia associated with Vitamin B₁₂ deficiency and for uncomplicated pernicious anaemia or Vitamin B₁₂ malabsorption.

Recommended Dosage

i) Prophylaxis of anaemia:

250-1000 mcg IM every month

ii) Uncomplicated pernicious anaemia or Vitamin B₁₂ malabsorption:

Initial 100 mcg daily for 5-10 days followed by 100-200 mcg monthly until complete remission is achieved.

Maintenance: up to 1000 mcg monthly.

Children: 30-50 mcg daily for 2 or more weeks (to a total dose of 1-5mg) or as prescribed.

Route of Administration

ATC Vitamin B₁₂ Injection Injection 1000 mcg/ml is administered by intramuscular only.

Contraindications

Any allergies to cyanocobalamin or any ATC Vitamin B₁₂ Injection 1000 mcg/ml ingredients. Vitamin B₁₂ Injection 1000 mcg/ml should not be used for Leber's disease or tobacco amblyopia since optic neuropathies may degenerate further.

Warnings and Precautions

ATC Vitamin B₁₂ Injection Injection 1000 mcg/ml is not a suitable form of vitamin B₁₂ for the treatment of optic neuropathies associated with raised plasma concentrations of cyanocobalamin.

As this preparation contains benzyl alcohol, its use should be avoided in children under two years of age. Not to be used in neonates.

Interaction with other medicament

Drugs that can decrease the absorption of ATC Vitamin B₁₂ Injection 1000 mcg/ml:

- Colchicine
- Antibiotics: neomycin.
- Omeprazole
- H₂ antagonist
- Aminosalicic acid

Large and continuous dose of folic acid may reduce vitamin B₁₂ concentration in blood.

Statement on usage during pregnancy and lactation

Pregnancy: Vitamin B₁₂ has been assigned to pregnancy category C by the FDA. Studies have revealed a progressive decline in the maternal serum level and an increase in the urinary excretion of vitamin B₁₂ during pregnancy. The fetus has been shown to concentrate vitamin B₁₂ since fetal cord serum levels are approximately three times maternal values. In view of a demonstrated increase in urinary excretion of vitamin B₁₂ during pregnancy, the recommended daily allowance (RDA) during pregnancy includes the addition of 1 mcg/day above the basal requirement of 3 mcg/day

Lactation: Vitamin B₁₂ appears in breastmilk. It is also excreted into human milk. American Academy of Pediatrics considers Vitamin B₁₂ is compatible with breastfeeding.

Adverse Effects/ Undesirable Effects

ATC Vitamin B₁₂ Injection 1000 mcg/ml is usually well tolerated. Rarely allergic reactions have been reported. Symptoms of serious allergic reaction including rash, itching/swelling (especially face/tongue/throat), severe dizziness and trouble breathing.

Overdose and Treatment

No overdosage has been reported with this drug.

Incompatibilities

Refer to 'Interactions with Other Medicaments' section.

Storage Conditions

Store below 30°C and protect from light.

Once opened, store in temperature below 30°C and can be used within 14 days.

Dosage forms and packaging available

Available in injection form, in multi-doses vial 10 mL.

Manufacturer

ATLANTIC LABORATORIES CORPORATION LTD.
111 Moo 7, Bang Phli Noi, Bang Bo, Samut Prakan
10560, Thailand.

Product Registration Holder

ATLANTIC LABORATORIES (M) SDN. BHD.
S-G-11, Gaya Resort Homes, Persiaran Kemuning Damai,
Seksyen 32, 40460 Shah Alam, Selangor, Malaysia.

Date Revision

03/12/2025

Atlantic Laboratories Corporation Ltd.
Samut Prakan, Thailand.

031225 MAL(R)