

ANTI HIST JUNIOR SYRUP 2mg/5ml

COMPOSITION

Each 5ml syrup contains: Chlorpheniramine Maleate 2.0mg,

Preservatives:

Methylparaben 9.0mg

Propylparaben 0.1mg

PRODUCT DESCRIPTION

A clear, orange liquid with orange flavour

PHARMACODYNAMICS

Chlorpheniramine is a potent antihistamine (H₁- antagonist).

Antihistamine diminish or abolish the actions of histamine in the body by competitive reversible blockade of histamine H₁-receptor sites on tissues. Chlorpheniramine also has anticholinergic activity.

Antihistamines act to prevent the release of histamine, prostaglandins and leukotrienes and have been shown to prevent the migration of inflammatory mediators. The actions of chlorpheniramine include inhibition of histamine on smooth muscle, capillary permeability and hence reduction of oedema and wheal in hypersensitivity reactions such as allergy and anaphylaxis.

PHARMACOKINETICS

Chlorpheniramine maleate is well absorbed from the gastro-intestinal tract, following oral administration. The effects develop within 30 minutes, are maximal within 1 to 2 hours and last 4 to 6 hours. The plasma half-life has been estimated to be 12 to 15 hours.

Chlorpheniramine is metabolised to the monodesmethyl and didesmethyl derivatives. About 22% of an oral dose excreted unchanged in the urine.

INDICATION

For the symptomatic relief of hayfever, vasomotor rhinitis, urticaria, angioneurotic oedema, food allergy, drug and serum reactions, insect bites.

RECOMMENDED DOSAGE

Children 6 to 12 years of age	: 5ml (2mg) every 4 to 6 hours. Maximum daily dose: 30ml (12mg) in 24 hours as required.
Children 2 to 5 years of age	: 2.5ml (1mg) every 4 to 6 hours. Maximum daily dose: 15ml (6mg) in 24 hours as required

ROUTE OF ADMINISTRATION

Oral administration

CONTRAINDICATIONS

Antihist Junior syrup is contra-indicated in patients who are hypersensitive to antihistamines or to any of the syrup ingredients.

The anticholinergic properties of chlorpheniramine are intensified by monoamine oxidase inhibitors (MAOIs). Contra-indicated in patients who have been treated with MAOIs within the last fourteen days.

WARNINGS AND PRECAUTIONS

This product contains Chlorpheniramine Maleate.

When used for treatment of cough and cold:

- a) Not to be used in children less than 2 years of age
- b) To be used with caution and doctor's/ pharmacist's advice in children 2 to 6 years of age.

Chlorpheniramine, in common with other drugs having anticholinergic effects, should be used with caution in epilepsy; raised intra-ocular pressure including glaucoma; prostatic hypertrophy; severe hypertension or cardiovascular disease; bronchitis, bronchiectasis or asthma; hepatic impairment; renal impairment. Children and the elderly are more likely to experience the neurological anticholinergic effects and paradoxical excitation (eg. Increased energy, restlessness, nervousness).

The anticholinergic properties of chlorpheniramine may cause drowsiness, dizziness, blurred vision and psychomotor impairment in some patients which may seriously affect ability to drive and use machinery.

The effects of alcohol may be increased and therefore concurrent use should be avoided.

Should not be used with other antihistamine containing products, including antihistamine containing cough and cold medicines.

Harmful for those suffering from alcoholism. To be taken into account in pregnant and breast feeding women, children and high risk groups such as patients with liver disease or epilepsy.

Patients with rare hereditary problems of fructose intolerance, glucose-galactose malabsorption or sucrose-isomaltase insufficiency should not take this medicine.

Long term use increases the risk of dental caries and it is essential that adequate dental hygiene is maintained.

Methylparaben and Propylparaben may cause allergic reactions (possibly delayed).

Keep out of the reach and sight of children.

DRUG INTERACTION

Concurrent use of chlorpheniramine and hypnotics or anxiolytics may cause an increase in sedative effects, therefore medical advice should be sought before taking chlorpheniramine concurrently with these medicines.

Chlorpheniramine inhibits phenytoin metabolism and can lead to phenytoin toxicity.

The anticholinergic effects of chlorpheniramine are intensified by MAOIs (see Contra- indications).

Effects on ability to drive and use machines

The anticholinergic properties of chlorpheniramine may cause drowsiness, dizziness, blurred vision and psychomotor impairment, which can seriously hamper the patients' ability to drive and use machinery.

PREGNANCY AND LACTATION

Pregnancy

There are no adequate data from the use of chlorpheniramine in pregnant women. The potential risk for humans is unknown, Use during the third trimester may result in reactions in the new born or premature neonates. Not to be used during pregnancy unless considered essential by a physician.

Lactation

Chlorpheniramine maleate and other antihistamines may inhibit lactation and may be secreted in breast milk. Not to be used during lactation unless considered essential by a physician.

ADVERSE EFFECTS

Specific estimation of the frequency of adverse events for OTC products is inherently difficult (particularly numerator data). The frequency of other adverse events identified during post- marketing use is unknown.

Blood and lymphatic system disorders

Unknown: haemolytic anaemia, blood dyscrasias

Immune system disorders:

Unknown: allergic reaction, angioedema, anaphylactic reactions

Metabolism and nutritional disorders:

Unknown: anorexia

Psychiatric disorders:

Unknown: confusion*, excitation*, irritability*, nightmares*, depression

Nervous system disorders*:

Very common: sedation, somnolence

Common: disturbance in attention, abnormal coordination, dizziness, headache

Eye disorders:

Common: blurred vision

Ear and labyrinth disorders

Unknown: tinnitus

Cardiac disorders:

Unknown: palpitations, tachycardia, arrhythmias

Vascular disorders:

Unknown: Hypotension

Respiratory, thoracic and Mediastinal disorders:

Unknown: thickening of bronchial secretions

Gastrointestinal disorders:

Common: nausea, dry mouth

Unknown: vomiting, abdominal pain, diarrhoea, dyspepsia

Hepatobiliary disorders:

Unknown: hepatitis including jaundice

Skin and subcutaneous disorders:

Unknown: exfoliative dermatitis, rash, urticaria, photosensitivity,

Musculoskeletal and connective tissue disorders:

Unknown: muscular twitching, muscle weakness.

Renal and Urinary disorders:

Unknown: Urinary retention

General disorders and administration site conditions:

Common: fatigue

Unknown: chest tightness

*Children and the elderly are more susceptible to neurological anticholinergic effects and paradoxical excitation (eg increased energy, restlessness, nervousness)

OVERDOSE AND TREATMENT**Symptoms and signs**

The estimated lethal dose of chlorpheniramine is 25 to 50mg/kg body weight. Symptoms and signs include sedation, paradoxical excitation of the CNS, toxic psychosis, convulsions, apnoea, anticholinergic effects, dystonic reactions and cardiovascular collapse including arrhythmias.

Treatment

Symptomatic and supportive measures should be provided with special attention to cardiac, respiratory, renal and hepatic functions and fluid and electrolyte balance. If overdosage is by the oral route, treatment with activated charcoal should be considered provided there are no contraindications for use and the overdose has been taken recently (treatment is most effective if given within an hour of ingestion.) Treat hypotension and arrhythmias vigorously. CNS convulsions may be treated with i.v. diazepam. Haemoperfusion may be used in severe cases.

STORAGE CONDITIONS

Store below 30°C. Keep container tightly closed. Protect from light.

PRESENTATION

HDPE Bottles and cap of 60ml and 120ml

MANUFACTURED BY

NORIPHARMA SDN. BHD. 200701034604 (792633-A)

Lot 5030, Jalan Teratai. 51/2 Mile Off Jalan Meru,
41050 Klang, Selangor, Malaysia

DATE OF REVISION

June 2026