

hovid-Chlorpheniramine Tablet 4 mg

VIHORxx-0 (FEO)

DESCRIPTION

Round, yellow uncoated tablet, shallow convex faces; "HD" embossed and scored on the same face.

COMPOSITION

Chlorpheniramine Maleate 4 mg per tablet

ACTIONS AND PHARMACOLOGY

Chlorpheniramine acts by competing with histamine for H1-receptor sites on effector cells, thus preventing responses mediated by histamine. Its antimuscarinic actions will provide a drying effect on the nasal mucosa.

It is readily absorbed after oral administration, metabolised in the liver and excreted mainly in the urine.

INDICATIONS

For symptomatic treatment of:

Allergic or vasomotor rhinitis, hay fever, urticaria, angioneurotic oedema, sensitivity reactions and other itching skin conditions including pruritis ani, pruritus vulvae, pruritus of drug rashes, contact dermatitis and insect bites.

CONTRAINDICATIONS

Chlorpheniramine Tablets are contra-indicated in patients who are hypersensitive to antihistamines or to any of the tablet ingredients.

The anticholinergic properties of chlorpheniramine are intensified by monoamine oxidase inhibitors (MAOIs). Chlorpheniramine Tablets is therefore contra-indicated in patients who have been treated with MAOIs within the last fourteen days.

WARNINGS AND PRECAUTIONS

Chlorpheniramine, in common with other drugs having anticholinergic effects, should be used with caution in epilepsy; raised intra-ocular pressure including glaucoma; prostatic hypertrophy; severe hypertension or cardiovascular disease; bronchitis, bronchiectasis or asthma; hepatic impairment; renal impairment. Children and the elderly are more likely to experience the neurological anticholinergic effects and paradoxical excitation (eg. increased energy, restlessness, nervousness).

The anticholinergic properties of chlorpheniramine may cause drowsiness, dizziness, blurred vision and psychomotor impairment in some patients which may seriously affect ability to drive and use machinery.

The effects of alcohol may be increased and therefore concurrent use should be avoided.

Should not be used with other antihistamine containing products, including antihistamine containing cough and cold medicines.

Patients with rare hereditary problems of galactose intolerance, Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicine,

Keep out of sight and reach of children.

MAIN SIDE/ADVERSE EFFECTS

Blood and lymphatic system disorders:

Unknown: haemolytic anaemia, blood dyscrasias

Immune system disorders:

Unknown: allergic reaction, angioedema, anaphylactic reactions

Metabolism and nutritional disorders:

Unknown: anorexia

Psychiatric disorders:

Unknown: confusion*, excitation*, irritability*, nightmares*, depression

Nervous system disorders*:

Very common: sedation, somnolence

Common: disturbance in attention, abnormal coordination, dizziness headache

Eye Disorders:

Common: blurred vision

Ear and labyrinth disorders:

Unknown: tinnitus

Cardiac disorders:

Unknown: palpitations, tachycardia, arrhythmias

Vascular disorders:

Unknown: Hypotension

Respiratory, thoracic and mediastinal disorders:

Unknown: thickening of bronchial secretions

Gastrointestinal disorders:

Common: nausea, dry mouth

Unknown: vomiting, abdominal pain, diarrhoea, dyspepsia

Hepatobiliary disorders:

Unknown: hepatitis, including jaundice

Skin and subcutaneous disorders:

Unknown: exfoliative dermatitis, rash, urticaria, photosensitivity

Musculoskeletal and connective tissue disorders:

Unknown: muscle twitching, muscle weakness

Renal and urinary disorders:

Unknown: urinary retention

General disorders and administration site conditions:

Common: fatigue

Unknown: chest tightness

*Children and the elderly are more likely to experience the neurological anticholinergic effects and paradoxical excitation (eg. increased energy, restlessness, nervousness).

INTERACTION WITH OTHER MEDICINAL PRODUCTS

Concurrent use of chlorpheniramine and hypnotics or anxiolytics may cause an increase in sedative effects, therefore medical advice should be sought before taking chlorpheniramine concurrently with these medicines.

Chlorpheniramine inhibits phenytoin metabolism and can lead to phenytoin toxicity.

The anticholinergic effects of chlorpheniramine are intensified by MAOIs

PREGNANCY AND LACTATION**Pregnancy**

There are no adequate data from the use of chlorpheniramine maleate in pregnant women. The potential risk for humans is unknown. Use during the third trimester may result in reactions in the new born or premature neonates. Not to be used during pregnancy unless considered essentially by a physician.

Lactation

Chlorpheniramine maleate and other antihistamine may inhibit lactation and may be secreted in breast milk. Not to be used during lactation unless considered essential by a physician

SYMPTOMS OF OVERDOSE AND TREATMENTS:**Symptoms and signs**

The estimated lethal dose of chlorpheniramine is 25 to 50mg/kg body weight. Symptoms and signs include sedation, paradoxical excitation of the CNS, toxic psychosis, convulsions, apnoea, anticholinergic effects, dystonic reactions and cardiovascular collapse including arrhythmias.

Treatment

Symptomatic and supportive measures should be provided with special attention to cardiac, respiratory, renal and hepatic functions and fluid and electrolyte balance. If over dosage is by the oral route, treatment with activated charcoal should be considered provided there are no contraindications for use and

hovid-Chlorpheniramine Tablet 4 mg

the overdose has been taken recently (treatment is most effective if given within an hour of ingestion). Treat hypotension and arrhythmias vigorously. CNS convulsions may be treated with i.v. diazepam. Haemoperfusion may be used in severe cases.

Effects on Ability to Drive and Use Machine

The anticholinergic properties of chlorpheniramine may cause drowsiness, dizziness, blurred vision and psychomotor impairment, which can seriously hamper the patients' ability to drive and use machinery.

DOSAGE AND ADMINISTRATION**Adults**

Oral, 4 mg every four to six hours as needed, not exceeding 24 mg daily.

Children 6 to 12 years

Oral, 2 mg three or four times daily as needed, not exceeding 12 mg a day.

Note: The information given here is limited. For further information, consult your doctor or pharmacist.

Storage:

Store below 30°C. Protect from moisture.

Presentation / Packing:

Tablet 4mg x 1000's, 100's

Blister of 10 x 10's tablets

Product Registration Holder: HOVID PHARMACY Sdn. Bhd.

121, Jalan Tunku Abdul Rahman (Jalan Kuala Kangsar),
30010 Ipoh, Perak, Malaysia.

Manufactured by: HOVID Bhd.

Lot 56442, 7 1/2 Miles, Jalan Ipoh/Chemor,
31200 Chemor, Perak, Malaysia.

Information date: December 2018