

Aw. No.: 2000XXXXX

Front

Desloratadine Leaflet

Back

2000XXXXX

SYNERRV DESLORATADINE 5
Desloratadine Tablets 5mg


COMPOSITION:

Each film coated tablet contains:
Desloratadine 5mg
Colours: Indigo Carmine and Titanium Dioxide
Excipients q.s.

PRESENTATION: Light blue colored, round shaped, biconvex, film coated tablets with both sides plain.

PROPERTIES:

Desloratadine is used as antihistaminic, to be administered orally. The chemical name of Desloratadine is 8-chloro-6,11-dihydro-11-(4-piperidinylidene)-5H-benzo[5,6]cyclohepta [1,2- b]pyridine Its emperical formula is C₂₀H₁₉ClN₂ and molecular weight is 310.8. Structural formula is as shown:



CLINICAL PHARMACOLOGY :

PHARMACODYNAMICS :

Pharmacotherapeutic group: antihistamines – H1 antagonist, ATC code: R06A X27

Mechanism of action

Desloratadine is a non-sedating, long-acting histamine antagonist with selective peripheral H1- receptor antagonist activity. After oral administration, desloratadine selectively blocks peripheral histamine H1-receptors because the substance is excluded from entry to the central nervous system.

Desloratadine has demonstrated antiallergic properties. These include inhibiting the release of proinflammatory cytokines such as IL-4, IL-6, IL-8, and IL-13 from human mast cells/basophils, as well as inhibition of the expression of the adhesion molecule P-selectin on endothelial cells. The clinical relevance of these observations remains to be confirmed.

PHARMACOKINETICS AND DRUG METABOLISM:

Absorption

Desloratadine plasma concentrations can be detected within 30 minutes of administration. Desloratadine is well absorbed with maximum concentration achieved after approximately 3 hours; the terminal phase half-life is approximately 27 hours. The degree of accumulation of desloratadine was consistent with its half-life (approximately 27 hours) and a once daily dosing frequency. The bioavailability of desloratadine was dose proportional over the range of 5 mg to 20 mg.

Distribution

Desloratadine is moderately bound (83 % - 87 %) to plasma proteins. There is no evidence of clinically relevant medicine accumulation following once daily dosing of desloratadine (5 mg to 20 mg) for 14 days.

Biotransformation

The enzyme responsible for the metabolism of desloratadine has not been identified yet, and therefore, some interactions with other medicinal products can not be fully excluded. It has shown that the medicinal product does not inhibit CYP3A4, CYP2D6 and is neither a substrate nor an inhibitor of P-glycoprotein.

Elimination

There is no effect of food (high-fat, high caloric breakfast) and grapefruit juice on the disposition of desloratadine.

INDICATIONS & USAGE:

Desloratadine Tablets are indicated for the rapid relief of symptoms associated with allergic rhinitis, such as sneezing, nasal discharge and itching, congestion/stuffiness, as well as ocular itching, tearing and redness.

Desloratadine Tablets are also indicated for the relief of symptoms associated with chronic idiopathic urticaria such as the relief of itching and the size and number of hives.

DOSAGE & METHOD OF ADMINISTRATION:

Adults and Adolescents (12 years of age and older): One Desloratadine Tablets 5 mg film-coated tablet once a day regardless of mealtime. For oral use.

INSTRUCTIONS FOR USE:

Oral use
The dose can be taken with or without food.

SIDE EFFECTS:

Summary of the safety profile
The most frequent of adverse reactions reported are fatigue, dry mouth and headache

Paediatric population

In 12 through 17 years of age, the most common adverse event was headache;

Tabulated list of adverse reactions

The adverse reactions reported in excess of placebo and other undesirable effects reported during the post-marketing period are listed in the following table. Frequencies are defined as very common, common, uncommon, rare, very rare and not known (cannot be estimated from the available data).

System organ class	Adverse reactions seen with desloratadine
Metabolism and nutrition disorders	
Not known	Increased appetite
Psychiatric disorders	
Very rare	Hallucinations
Not known	Abnormal behaviour, aggression
Nervous system disorders	
Common	Headache
Very rare	Dizziness, somnolence, insomnia, psychomotor hyperactivity, seizures
Cardiac disorders	
Very rare	Tachycardia, palpitations
Not known	QT prolongation
Gastrointestinal disorders	
Common	Dry mouth
Very rare	Abdominal pain, nausea, vomiting, dyspepsia, diarrhoea
Hepatobiliary disorders	
Very rare	Elevations of liver enzymes, increased bilirubin, hepatitis
Not known	Jaundice
Skin and subcutaneous tissue disorders	
Not known	Photosensitivity
Musculoskeletal and connective tissue disorders	
Very rare	Myalgia
General disorders and administration site conditions	
Common	Fatigue
Very rare	Hypersensitivity reactions (such as anaphylaxis, angioedema, dyspnoea, pruritus, rash, and urticaria)
Not known	Asthenia
Investigations	
Not known	Weight increased

DRUG INTERACTIONS:

No clinically relevant interactions were observed with desloratadine tablets in which erythromycin or ketoconazole were co-administered.
Desloratadine tablets taken concomitantly with alcohol did not potentiate the performance impairing effects of alcohol. However, cases of alcohol intolerance and intoxication have been reported during post-marketing use. Therefore, caution is recommended if alcohol is taken concomitantly.

SPECIAL WARNINGS AND PRECAUTIONS FOR USE:

Efficacy and safety of Desloratadine Tablets in children under 12 years of age have not been established.
Desloratadine should be administered with caution in patients with a medical or family history of seizures. In particular, young children may be more susceptible to developing new seizures under desloratadine treatment. Healthcare providers may consider discontinuing desloratadine in patients who experience a seizure while on treatment.
In the case of severe renal insufficiency, desloratadine should be used with caution

PREGNANCY AND LACTATION:

Pregnancy

A large amount of data on pregnant women (more than 1,000 pregnancy outcomes) indicate no malformative nor foeto/neonatal toxicity of desloratadine. As a precautionary measure, it is preferable to avoid the use of desloratadine during pregnancy.

Breast-feeding

Desloratadine has been identified in breastfed newborns/infants of treated women. The effect of desloratadine on newborns/infants is unknown. A decision must be made whether to discontinue breast-feeding or to discontinue/abstain from desloratadine therapy taking into account the benefit of breast feeding for the child and the benefit of therapy for the woman.

Fertility

There are no data available on male and female fertility.

OVERDOSAGE:

The adverse event profile associated with overdosage, as seen during post-marketing use, is similar to that seen with therapeutic doses, but the magnitude of the effects can be higher.

Treatment

In the event of overdose, consider standard measures to remove unabsorbed active substance. Symptomatic and supportive treatment is recommended. Desloratadine is not eliminated by haemodialysis; it is not known if it is eliminated by peritoneal dialysis.

Symptoms

No clinically relevant effects were observed when 45mg of desloratadine was administered.

Paediatric population

The adverse event profile associated with overdosage, as seen during post-marketing use, is similar to that seen with therapeutic doses, but the magnitude of the effects can be higher.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINE

Desloratadine has no or negligible influence on the ability to drive and use machines. Patients should be informed that most people do not experience drowsiness. Nevertheless, as there is individual variation in response to all medicinal products, it is recommended that patients are advised not to engage in activities requiring mental alertness, such as driving a car or using machines, until they have established their own response to the medicinal product.

CONTRAINDICATION:

Hypersensitivity to the active substance, to any of the excipients, or to loratadine.

STORAGE CONDITION

Store below 30°C. Protect from moisture.

PACK SIZE:

Blister pack of 10 tablets. Such 1, 5 or 10 Blisters of 10 tablets each are packed in a carton alongwith insert.

KEEP ALL MEDICINES OUT OF REACH OF CHILDREN.

PRODUCT REGISTRATION HOLDER:

Synerrv Sdn Bhd, SO-29-2, MENARA 1,
KL ECO CITY, JALAN BANGSAR,
KG HAJI ABDULLAH KUMUK,
59200 KUALA LUMPUR WILAYAH
PERSEKUTUAN KUALA LUMPUR, MALAYSIA

Manufactured by :

 **Ind-Swift Laboratories Limited**
Off- NH-21, Village Jawaharpur,
Tehsil Derabassi, Distt. S.A.S Nagar
(Mohali) Punjab (India) Pin -140507

Date of Revision: Septemebr’2025

This artwork is Technically and Pharmaceutically approved (Customer Signature)

Item Code No.	2000XXXXX	Revision No.:		02		
		Prev. Aw. No.:		200011065		
Packaging Development	Marketing	Regulatory Affairs	Production	Quality Control	Quality Assurance	Party Approval
Specifications: Leaflet Dimension: 110 x 280mm Paper / GSM: Maplitho / 60GSM Pantone no.: 				Pharmacode		No.
				Front		4017
				Back		4018