Pharma code
Reading Direction

Pharma code
Reading Direction

SIDE 1

10 mm

10 mm

5 mm

320 mm

BACTAFUZ

Fusidic Acid Cream 2% w/w

PRODUCT NAME:

BACTAFUZ : Fusidic Acid Cream 2% w/w

NAME AND STRENGTH OF ACTIVE INGREDIENTS:

BACTAFUZ: Each gram contains: Fusidic acid Ph. Eur. 20mg, Potassium Sorbate Ph. Eur. (As preservative) 2.7mg

PRODUCT DESCRIPTION:

BACTAFUZ: White, smooth, homogenous cream.

PHARMACOLOGICAL PROPERTIES

Pharmacodynamics Properties

Pharmacotherapeutic group: Other antibiotics for topical use; ATC code: D06AX01

Fusidic acid is a potent antibacterial agent. Fusidic acid and its salts show fat and water solubility and strong surface activity and exhibit unusual ability to penetrate intact skin. Concentrations of 0.03 - 0.12 mcg Fusidic acid per ml inhibit nearly all strains of *Staphylococcus aureus*. Topical application of Fusidic acid is also effective against *Streptococci, Corynebacteria, Neisseria* and certain *Clostridia.*

Pharmacokinetics Properties

In vitro studies show that Fusidic acid can penetrate intact human skin. The degree of penetration depends on factors such as the duration of exposure to Fusidic acid and the condition of the skin. Fusidic acid is excreted mainly in the bile with little excreted in the urine.

INDICATIONS:

BACTAFUZ is indicated for the treatment of skin infections caused by *Staphylococci, Streptococci, Propionibacterium acnes, Corynebacterium minutissimum* and other organisms sensitive to fusidic acid. The most important indications being impetigo, infected wounds, folliculitis, boils, sycosis barbae, carbuncles, hidradenitis, paronychia and erythrasma

BACTAFUZ is an invisible, non staining preparation, which is cosmetically acceptable for treatment of face and scalp infections.

RECOMMENDED DOSAGE:

For external use only.

Apply to the affected area 2-3 times daily, generally for a period of 7 days, except in treatment of acne where therapy should be extended for a longer period according to the nature of the disease

CONTRAINDICATIONS:

Hypersensitivity to the active substance or to any of the excipients.

WARNING AND PRECAUTIONS:

Bacterial resistance among *Staphylococcus aureus* has been reported to occur with the use of topical Fusidic acid. As with all antibiotics, extended or recurrent use may increase the risk of developing antibiotic resistance.

Extended or recurrent use may increase the risk of developing contact sensitisation.

BACTAFUZ contains butylhydroxyanisole, cetyl alcohol and potassium sorbate. These excipients may cause local skin reactions (e.g. contact dermatitis). Butylhydroxyanisole may also cause irritation to the eyes and mucous membranes.

BACTAFUZ should therefore be used with care when applied in the proximity of the eyes.

INTERACTIONS WITH OTHER MEDICAMENTS:

No interaction studies have been performed. Interactions with systemically administered medicinal products are considered minimal as the systemic absorption of topical Fusidic acid is negligible.

PREGNANCY AND LACTATION:

Pregnancy

No effects during pregnancy are anticipated, since systemic exposure to topically-applied Fusidic acid/sodium fusidate is negligible. Topical Fusidic acid can be used during pregnancy.

Breast-feeding

No effects on the breast-fed new-born/infant are anticipated since the systemic exposure of topically-applied Fusidic acid/sodium fusidate to the breast-feeding woman is negligible. Topical Fusidic acid can be used during breast-feeding but it is recommended to avoid applying topical Fusidic acid on the breast.

Fertility

There are no clinical studies with topical Fusidic acid regarding fertility. No effects in women of childbearing potential are anticipated, since systemic exposure following topically-applied Fusidic acid/sodium fusidate is negligible.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINES:

Fusidic acid cream 2.0% w/w administered topically has no or negligible influence on the ability to drive and use machines

UNDESIRABLE EFFECTS:

The most frequently reported adverse reactions during treatment are various skin reactions such as pruritus and rash, followed by application site conditions such as pain and irritation, which all occurred in less than 1% of patients.

Hypersensitivity and angioedema have been reported.

Immune system disorders

Rare**:** Hypersensitivity

Eye disorders

Rare: Conjunctivitis

Skin and subcutaneous tissue disorders

Uncommon: Dermatitis (including dermatitis contact, eczema), Rash*, Pruritus, Erythema

**Various types of rash reactions such as erythematous, pustular, vesicular, maculo-papular and papular have been reported. Rash generalised has also occurred.*

Rare: Angioedema, Urticaria, Blister

General disorders and administration site conditions

5 mm

Font Size : 8 Point (Times New Roman)

150 mm

SIDE 2

5 mm

320 mm

Uncommon: Application site pain (including skin burning sensation), Application site irritation

Paediatric population: Frequency, type and severity of adverse reactions in children are expected to be the same as in adults.

OVERDOSE AND TREATMENT:

Overdose is unlikely to occur

Unless hypersensitivity to Fusidic acid or any of the excipients exists, accidental ingestion of Fusidic acid cream is unlikely to cause any harm. The total quantity of Fusidic acid will usually not exceed the approved total daily oral dose of Fusidic acid containing products except in children aged less than 1 year and weighing ≤ 10 kg. Although in this instance a child of this particular age group is unlikely to ingest a whole tube of Fusidic acid cream. The concentration of the excipients is too low to constitute a safety risk.

INCOMPATIBILITIES: Not applicable

SHELF LIFE: 36 months from the date of manufacture.

SPECIAL PRECAUTIONS FOR STORAGE:

Store below 30ºC, protected from light.
Do not freeze.
Keep medicine out of reach of children.

DOSAGE FORMS AND PACKAGING AVAILABLE:

BACTAFUZ: 15 g Aluminium collapsible tube with an interior coating of lacquer closed with HDPE cap, containing Fusidic acid Cream, packed in a carton along with package insert.

DATE OF REVISION: Mar' 2019

Manufactured by:
Encube Ethicals Pvt. Ltd.
Plot No. C1, Madkaim Ind. Estate, Madkaim,
Post : Mardol, Ponda, Goa - 403 404, (India.)

Mankind

Manufactured for:
MANKIND PHARMA LTD.
208, Okhla Industrial Estate, Phase-3
New Delhi-110020 (INDIA)

EMA9SPBJBJMY64400

MFB5B/00

5 mm