

MUCOXOL TABLET 30MG

Composition:

Each tablet contains Ambroxol Hydrochloride 30mg

Description: A white to off white colored round tablet with 9.1mm diameter and a score on one side.

Pharmacodynamic:

Ambroxol hydrochloride is a metabolite of bromhexine, which liquefies and thins the mucus of the passage. By this way it promotes the expectoration of blocked – up secretions and also eases cough.

Ambroxol has been shown to increase respiratory tract secretion. It enhances pulmonary surfactant production and stimulates ciliary activity. These actions result in improved mucus flow and transport (mucociliary clearance). Improvement of mucociliary clearance has been shown in clinical pharmacologic studies. Enhancement of fluid secretion and mucociliary clearance facilitates expectoration and eases cough.

Pharmacokinetic:

Absorption after oral administration is fast and complete. The active is metabolized to dibromoanthranilic acid under phase I metabolism and it is then further broken down under phase II metabolism to conjugates, predominantly glucuronides. Elimination half-life is estimated as 20-25 hours. Excretion is mainly through urine.

Indication:

Secretolytic therapy in acute and chronic bronchopulmonary diseases associated with abnormal mucus secretion and impaired mucus transport.

Recommended dosage:

Tablet

Adults: 1 tablet 3 times daily. The therapeutic effect may be enhanced by administering 2 tablets 2 times daily and should be taken after meals with liquid. Mucoxol is not recommended for children under 2 years of age without medical advice.

Route of Administration:

Oral administration

Contraindication:

Contraindicated during the first trimester of pregnancy and in patients who sensitive to Ambroxol hydrochloride or to any of the ingredients of this product.

Warning and precaution:

Patients with liver and kidney problems must be treated cautiously.

Very rare cases of chronically associated severe skin impairments such as Stevens Johnson Syndrome, Toxic Epidermal Necrolysis (TEN), Erythema Multiforme (EM) and Accute Generalized Exathematous Pustulosis (AGEP) have been reported. In most cases, these could be explained by the severity of the underlying disease or concomitant administration of another drug. In the early stages of such severe skin reactions, initially

only non-specific flu-like symptoms appear, e.g. fever, arthralgia, running nose, cough, and sore throat. If skin or mucous membrane damage occurs, seek medical advice immediately and discontinue treatment as precaution.

Drug interaction:

Concomitant administration of ambroxol together with antibiotics (amoxicillin, cefuroxime, erythromycin, doxycycline) leads to higher antibiotic concentration in the lung tissue.

Pregnancy and lactation:

There were no reports of ill-effects during pregnancy. Nevertheless, the use of ambroxol especially during the first trimester is not recommended. During lactation, ambroxol should only be used if absolutely necessary.

Side effects:

Occasional, mild side effects from Mucoxol include:

- Nausea
- Diarrhoea
- Indigestion
- A feeling of fullness in the stomach (a bloated feeling)

Serious allergic reactions have very rarely occurred with the use of Mucoxol.

They include:

- Skin rashes
- Difficulty in breathing
- Swelling of the face, lips, mouth, tongue or throat which may cause difficulty swallowing or breathing.

Immune System Disorders

Frequency not known: Anaphylactic reactions including anaphylactic shock.

Skin and Subcutaneous Skin Disorders

Frequency not known: Severe skin reactions (including Stevens Johnson syndrome, Toxic epidermal necrolysis (TEN), Erythema Multiforme (EM) and Acute Generalized Exanthematous Pustulosis (AGEP))

Overdosage:

No symptoms of overdosage have been reported. If they occur, symptomatic treatment should be provided.

Storage:

Store below 30°C.

Keep medicine out of reach of children.

Jauhkan ubat daripada kanak-kanak.

Presentation:

Blisters of 10 tablets. Box 100's, 250's and 500's.

Manufactured by:

Noripharma Sdn. Bhd.
Lot 5030, Jalan Teratai,
5 1/2 Mile off Jalan Meru,
41050 Klang, Selangor, MALAYSIA

Date of revision:

19 February 2019