

BUCOLYN 20MG/G POWDER

FOR ANIMAL USE ONLY

Product Name

BUCOLYN 20MG/G POWDER

Name and Strength of Active Substance(s)

Each gram contains 20 mg Bromhexine Hydrochloride (equivalent to 18.23 mg Bromhexine).

Product Description

A white to off white powder, soluble in water.

Pharmacodynamics properties

Bromhexine is a mucoregulator. By activating the secretion of the seromucous glands, bromhexine helps to re-establish the viscosity and elasticity of bronchial secretions in the tracheobronchial tree. In addition, its expectorant action encourages mobilisation of mucus and enables effective bronchial drainage, thereby improving the functioning and defence capability of the lung. These two simultaneous actions lead to an abundant discharge and facilitate a productive cough. It breaks down the network of acid glycoprotein fibres found in mucoid sputum, which are mainly responsible for the characteristic viscosity. Bromhexine modifies the distribution of antibiotics in the organism and increases their concentration in the serum and in the nasal secretions (e.g. spiramycin, tylosin and oxytetracycline). When administered concomitantly with the product, antimicrobial agents should, nevertheless, not be underdosed.

Pharmacokinetics properties

In pigs, bromhexine is rapidly absorbed following oral administration. In cattle, plasma concentrations increase progressively over several hours following administration. Due to the lipophilic character of bromhexine, it has a strong affinity for lipid tissues and a slow depletion profile from these tissues. Bromhexine is largely metabolised into more polar compounds.

Indication

Mucolytic treatment of congested respiratory tract in cattle (calves), pigs, poultry, dogs and cats.

Target Species

Cattle (Calves), pigs, poultry, dogs and cats.

Route of administration

To be mixed in animal drinking water or food and administer orally.

Dosage

0.45 mg of bromhexine per kg bodyweight daily, equivalent to 0.25 g of product per 10 kg bodyweight per day administered for 3 to 10 consecutive days, orally dissolved in the drinking water or dusted on food, which is corresponding to:

Species	Dosage (amount of product)
Calves	1.25 – 2.5 g
Pigs (Adult)	1.25 – 2.5 g
Pigs (Young)	0.5 – 1.25 g
Dogs	0.125 – 1 g
Cats	0.125 g
Poultry	0.5 g of product for 3 L of drinking water.

Direction of use: The treatment should last at least 3 days. Prepare fresh solution immediately before use. Any unused medicated water should be discarded after 24 hours.

Contraindications

Do not use in cases of pulmonary oedema. In case of serious lungworm infection, the drug should only be used 3 days after the commencement of the anthelmintic treatment.

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

Special warnings for each target species

None.

Special precautions for use

Special precautions to be taken by the person administering the veterinary medicinal product to animal: People with known hypersensitivity to bromhexine or any of the excipients should avoid contact with the veterinary medicinal product. Avoid contact with skin and eyes. In case of accidental eye contact, flush the affected eye with copious amounts of clean running water. Wash hands and exposed skin after administering the product.

Interactions with Other Medicaments

The product may be used in conjunction with antibiotics and/or sulphonamides and bronchodilators. Bromhexine modifies the distribution of antibiotics in the organism and increases their concentration in the respiratory system.

Use during pregnancy and lactation

Studies in laboratory animals have not produced evidence of foetotoxic effects or effects on fertility at the recommended dose. However this has not been specifically studied in the target species. Use only according to the benefit-risk assessment by the responsible veterinarian.

Adverse Effects

None known

Overdose and Treatment

None known.

Withdrawal Period(s)

Cattle (calves): Meat and offal: Zero days. Not permitted for use in cows producing milk for human consumption.

Pigs : Meat and offal: Zero days.

Poultry: Meat and offal: Zero days.

Eggs: Do not use in birds producing eggs for human consumption, during and 4 weeks before the laying phase.

Shelf-life

Shelf life as packaged for sale: 3 year

Shelf life after first opening : 24 hours

Shelf life after reconstitution : 24 hours

Allowable Maximum Residual Limit (MRL)

No MRL required [Ref. French Agency for Veterinary Medicinal Product MRL regulation 37/2010]

Storage Conditions

Store in a dry place below 30°C. Keep out of reach of children. Jauhi ubat daripada kanak-kanak.

Disposal of Containers

Any unused veterinary medicinal product or waste material derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Packaging available

Available in 100g, 200g, 500g and 1000g powder per sachet

Manufacturer and Product Registration Holder:

Life Biopharma Sdn. Bhd. (company no. 546163-D)

No. 250-251, Jalan Industri Galla 13, Kawasan Perindustrian Galla, 70200 Seremban, Negeri Sembilan, Malaysia.

Date of Revision of Package Insert

08th July 2019