

CLOFAN


Clotrimazole Pessary 500 mg

GENERAL CHARACTERISTICS:
General physico-chemical properties:
White to light yellow colour torpedo shaped pessary.
Composition:
Each pessary contains:
Clotrimazole BP 500 mg
Excipient: Hard fat

PHARMACODYNAMIC PROPERTIES:
Pharmacotherapeutic group: Anti-infective
ATC code: G01AF02

Mechanism of action
Clotrimazole acts against fungi by inhibiting ergosterol synthesis. Inhibition of ergosterol synthesis leads to structural and functional impairment of the fungal cytoplasmic membrane.
Clotrimazole has a broad antimycotic spectrum of action in vitro and in vivo, which includes dermatophytes, yeasts, moulds, etc.
Under appropriate test conditions, the MIC values for these types of fungi are in the region of less than 0.062-8.0 µg/ml substrate. The mode of action of clotrimazole is fungistatic or fungicidal depending on the concentration of clotrimazole at the site of infection. In-vitro activity is limited to proliferating fungal elements; fungal spores are only slightly sensitive.
Primarily resistant variants of sensitive fungal species are very rare; the development of secondary resistance by sensitive fungi has so far only been observed in very isolated cases under therapeutic conditions.

PHARMACOKINETIC PROPERTIES:
Pharmacokinetic investigations after vaginal application have shown that only a small amount of clotrimazole (3-10% of the dose) is absorbed. Due to the rapid hepatic metabolism of absorbed clotrimazole into pharmacologically inactive metabolites the resulting peak plasma concentrations of clotrimazole after vaginal application of a 500 mg dose were less than 10 mg/ml, reflecting that clotrimazole applied intravaginally does not lead to measurable systemic effects or side effects.
INDICATIONS:
Infections of the genital region (vaginitis) caused by fungi (usually Candida) and superinfections caused by Clotrimazole-sensitive bacteria.
DOSAGE AND ADMINISTRATION:
Clofan Pessary should be inserted into the vagina, as high as possible.
Adults: One 500 mg pessary should be inserted at night. A second treatment may be carried out if necessary.
Generally: Treatment during the menstrual period should not be performed due to the risk of the pessary being washed out by the menstrual flow. The treatment should be finished before the onset of menstruation.
Do not use tampons, intravaginal douches, spermicides or other vaginal products while using this product.
Vaginal intercourse should be avoided in case of vaginal infection and while using this product because the partner could become infected.
Children: Paediatric usage is not recommended.
Mode of administration: Vaginal
• Remove the pessary from its plastic packet using the easy open pessary (figure 1).
• Pull plunger A until it stops. Place a vaginal pessary into applicator B (figure 2).
• Carefully insert the applicator containing the pessary as deeply as possible into the vaginal. This is best achieved when lying on your back (figure 3).
• Push plunger A carefully until it stops, thereby depositing the pessary into the vagina. Remove the applicator.



CONTRAINDICATIONS:
Hypersensitivity to Clotrimazole or any ingredient in this medicine.
SPECIAL WARNINGS AND PRECAUTIONS FOR USE:
Medical advice should be sought if this is the first time the patient has experienced symptoms of candidal vaginitis. Before using Clofan Pessary, medical advice must be sought if any of the following are applicable:
• More than two infections of candidal vaginitis in the last 6 months.
• Previous history of sexually transmitted disease or exposure to partner with sexually transmitted disease.
• Pregnancy or suspected pregnancy.
• Age under 16 or over 60 years.
• Known hypersensitivity to imidazoles or other vaginal antifungal products.
Clofan Pessary should not be used if the patient has any of the following symptoms where upon medical advice should be sought:
• Irregular vaginal bleeding.

CLOFAN Pessary/MY/F

- Abnormal vaginal bleeding or a blood-stained discharge.
- Vulval or vaginal ulcers, blisters or sores.
- Lower abdominal pain or dysuria.
- Any adverse events such as redness, irritation or swelling associated with the treatment.
- Fever or chills.
- Nausea or vomiting.
- Diarrhoea.
- Foul smelling vaginal discharge.

Patients should be advised to consult their physician if the symptoms have not been relieved within one week of using Clofan Pessary. Clofan Pessary can be used again if the candidal infection returns after 7 days. However, if the candidal infection recurs more than twice within six months, patients should be advised to consult their physician.

INTERACTIONS WITH OTHER MEDICINAL PRODUCTS:
Laboratory tests have suggested that, when used together, this product may cause damage to latex contraceptives. Consequently, the effectiveness of such contraceptives may be reduced. Patients should be advised to use alternative precautions for at least five days after using this product. Concomitant medication with vaginal clotrimazole and oral tacrolimus (FK-506; immunosuppressant) might lead to increased tacrolimus plasma levels and similarly with sirolimus. Patients should thus be closely monitored for signs and symptoms of tacrolimus or sirolimus overdosage, if necessary by determination of the respective plasma levels.

FERTILITY, PREGNANCY AND LACTATION:
Fertility:
No human studies of the effects of clotrimazole on fertility have been performed; however, animal studies have not demonstrated any effects of the drug on fertility.

Pregnancy:
There are limited amount of data from the use of clotrimazole in pregnant women. Animal studies with clotrimazole have shown reproductive toxicity at high oral doses at the low systemic exposures of clotrimazole following vaginal treatment harmful effects with respect to reproductive toxicity are not predicted. Clotrimazole can be used during pregnancy, but only under the supervision of a physician or midwife.

Lactation:
Available pharmacodynamic/toxicological data in animals have shown excretion of clotrimazole/metabolites in milk after intravenous administration. A risk to the suckling child cannot be excluded. A decision must be made whether to discontinue breast-feeding or to discontinue/abstain from clotrimazole therapy taking into account the benefit of breast-feeding for the child and the benefit of therapy for the woman.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINES:
The medication has no or negligible influence on the ability to drive or use machinery.

UNDESIRABLE EFFECTS:
As the listed undesirable effects are based on spontaneous reports, assigning accurate frequency of occurrence for each is not possible.

Immune system disorders:
Allergic reaction (syncope, hypotension, dyspnea, urticaria, pruritus).

Reproductive system and breast disorders:
Genital peeling, pruritus, rash, oedema, erythema, discomfort, burning, irritation, pelvic pain, vaginal haemorrhage.

Gastrointestinal disorders:
Abdominal pain

OVERDOSAGE AND TREATMENT:
No risk of acute intoxication is seen as it is unlikely to occur following a single vaginal or dermal application of an overdose (application over a large area under conditions favorable to absorption) or inadvertent oral ingestion. There is no specific antidote. However, in the event of accidental oral ingestion, routine measures such as gastric lavage should be performed only if clinical symptoms of overdose become apparent (e.g. dizziness, nausea or vomiting). Gastric lavage should be carried out only if the airway can be protected adequately.

STORAGE CONDITION:
Store in dry place, temperature below 30°C. Do not freeze.
Keep all medicines out of reach of children.

DOSAGE FORM AND PACKING AVAILABLE:
PVC/PE strip of one pessary with 1 HDPE applicator. (Box of 1's)

NAME AND ADDRESS OF MANUFACTURER:
Kusum Healthcare Pvt. Ltd.
SP-289(A), RIICO Industrial Area,
Chopanki, Bhiwadi, Dist. Alwar,
(Rajasthan), India

PRODUCT REGISTRATION HOLDER:
Synerrv Sdn Bhd,
SO-29-2, Menara 1, KL Eco City,
Jalan Bangsar, KG Haji Abdullah Hukm,
59200 Kuala Lumpur,
Wilayah Persekutuan Kuala Lumpur, Malaysia

DATE OF REVISION OF PACKAGE INSERT: Dec 2025

##

CLOFAN Pessary/MY/B

INSERT CLOFAN PESSARY