

## PACK INSERT

**BACTOPIC®**

**MUPIROCIN OINTMENT 2% w/w**

### GENERAL CHARACTERISTICS

#### **General Physio-Chemical Properties:**

White or almost white ointment.

#### **Composition:**

Each gram of ointment contains:

Mupirocin BP..... 20 mg

**Excipients:** Polyethylene glycol 3350 & Polyethylene glycol 400

### PHARMACODYNAMIC PROPERTIES:

**Pharmacotherapeutic group:** Antibiotics and chemotherapeutics for dermatological use.

**ATC code:** D06AX09

#### **Mode of action**

Mupirocin is a novel antibiotic produced through fermentation by *Pseudomonas fluorescens*.

Mupirocin inhibits isoleucyl transfer-RNA synthetase, thereby arresting bacterial protein synthesis.

Mupirocin has bacteriostatic properties at minimum inhibitory concentrations and bactericidal properties at the higher concentrations reached when applied locally.

#### **Mechanism of Resistance**

Low-level resistance in staphylococci is thought to result from point mutations within the usual staphylococcal chromosomal gene (ileS) for the target isoleucyl tRNA synthetase enzyme. High-level resistance in staphylococci has been shown to be due to a distinct, plasmid encoded isoleucyl tRNA synthetase enzyme.

Intrinsic resistance in Gram negative organisms such as the *Enterobacteriaceae* could be due to poor penetration of the outer membrane of the Gram negative bacterial cell wall.

Due to its particular mode of action, and its unique chemical structure, Mupirocin does not show any cross-resistance with other clinically available antibiotics.

#### **Microbiological Susceptibility**

The prevalence of acquired resistance may vary geographically and with time for selected species, and local information on resistance is desirable, particularly when treating severe infections. As necessary, expert advice should be sought when the local prevalence of resistance is such that the utility of the agent in at least some types of infection is questionable.

|                                                                                 |
|---------------------------------------------------------------------------------|
| <b>Commonly susceptible species</b>                                             |
| <i>Staphylococcus aureus</i> *                                                  |
| <i>Streptococcus pyogenes</i> *                                                 |
| <i>Streptococcus</i> spp. ( $\beta$ haemolytic, other than <i>S. pyogenes</i> ) |
| <b>Species for which acquired resistance may be a problem</b>                   |
| <i>Staphylococcus</i> spp., coagulase negative                                  |
| <b>Inherently resistant organisms</b>                                           |
| <i>Corynebacterium</i> spp.                                                     |
| <i>Micrococcus</i> spp.                                                         |

\*Activity has been satisfactorily demonstrated in clinical studies

### **PHARMACOKINETIC PROPERTIES:**

After topical application of BACTOPIC (MUPIROCIN OINTMENT 2%), Mupirocin is only very minimally absorbed systemically and that which is absorbed is rapidly metabolized to the antimicrobially inactive metabolite, monic acid. Penetration of Mupirocin into the deeper epidermal and dermal layers of the skin is enhanced in traumatised skin and under occlusive dressings.

#### *Elderly patients*

No restrictions unless there is evidence of moderate or severe renal impairment. (see section special warning and precaution for use)

### **INDICATIONS:**

Bacterial skin infections, e.g. impetigo, folliculitis and furunculosis.

### **DOSAGE AND MODE OF ADMINISTRATION:**

#### *Adults and children:*

BACTOPIC (MUPIROCIN OINTMENT 2%) should be applied to the affected area 2 to 3 times a day, for up to 10 days, depending on the response.

The area may be covered with a dressing or occluded if desired.

Do not mix with other preparations as there is a risk of dilution, resulting in a reduction of the antibacterial activity and potential loss of stability of the mupirocin in the ointment

#### *Method of administration:*

Topical

A small quantity of BACTOPIC (MUPIROCIN OINTMENT 2%) should be applied to cover the affected area. The treated area may be covered by a dressing.

Any product remaining at the end of treatment should be discarded.

Do not mix with other preparations as there is a risk of dilution, resulting in a reduction of the antibacterial activity and potential loss of stability of the Mupirocin in the ointment.

### **CONTRAINDICATIONS:**

Hypersensitivity to the active substance or to any of the excipients.

This BACTOPIC (MUPIROCIN OINTMENT 2%) formulation is not suitable for ophthalmic or intranasal use.

### **SPECIAL WARNINGS AND PRECAUTIONS FOR USE:**

Should a possible sensitisation reaction or severe local irritation occur with the use of BACTOPIC (MUPIROCIN OINTMENT 2%) , treatment should be discontinued, the product should be washed off and appropriate therapy instituted.

As with other antibacterial products, prolonged use may result in overgrowth of non-susceptible organisms.

Pseudomembranous colitis has been reported with the use of antibiotics and may range in severity from mild to life-threatening. Therefore, it is important to consider its diagnosis in patients who develop diarrhoea during or after antibiotic use. Although this is less likely to occur with topically applied Mupirocin, if prolonged or significant diarrhea occurs or the patient experiences abdominal cramps, treatment should be discontinued immediately and the patient investigated further.

#### *Renal impairment*

Polyethylene glycol can be absorbed from open wounds and damaged skin and is excreted by the kidneys. In common with other polyethylene glycol based ointments, BACTOPIC (MUPIROCIN OINTMENT 2%) should not be used in conditions where absorption of large quantities of polyethylene glycol is possible, especially if there is evidence of moderate or severe renal impairment.

BACTOPIC (MUPIROCIN OINTMENT 2%) is not suitable for:

- Ophthalmic use
- Intranasal use
- Use in conjunction with cannulae
- At the site of central venous cannulation.

Avoid contact with the eyes. If contaminated, the eyes should be thoroughly irrigated with water until the ointment residues have been removed

### **INTERACTIONS WITH OTHER MEDICINAL PRODUCTS:**

No drug interactions have been identified.

### **FERTILITY, PREGNANCY AND LACTATION:**

#### **Pregnancy:**

Reproduction studies on Mupirocin in animals have revealed no evidence of harm to the foetus as there is no clinical experience on its use during pregnancy, Mupirocin should only be used in pregnancy when the potential benefits outweigh the possible risks of treatment.

**Fertility:**

There are no data on the effects of Mupirocin on human fertility. Studies in rats showed no effects on fertility.

**Lactation:**

There is no information on the excretion of Mupirocin in milk. If a cracked nipple is to be treated, it should be thoroughly washed prior to breast feeding.

**EFFECTS ON ABILITY TO DRIVE AND USE MACHINES:**

No adverse effects on the ability to drive or operate machinery have been identified.

**SIDE EFFECTS:**

Adverse reactions are listed below by system organ class and frequency. Frequencies are defined as: very common ( $\geq 1/10$ ), common ( $\geq 1/100$ ,  $< 1/10$ ), uncommon ( $\geq 1/1000$ ,  $< 1/100$ ), rare ( $\geq 1/10,000$ ,  $< 1/1000$ ), very rare ( $< 1/10,000$ ), including isolated reports.

*Immune system disorders:*

Very rare: Systemic allergic reactions including anaphylaxis, generalised rash, urticaria and angioedema have been reported with Bactopic<sup>®</sup> (Mupirocin ointment 2%).

*Skin and subcutaneous tissue disorders:*

Common: Burning localised to the area of application.

Uncommon: Itching, erythema, stinging and dryness localised to the area of application.  
Cutaneous sensitisation reactions to Mupirocin or the ointment base.

**OVERDOSE AND TREATMENT:****Symptoms:**

There is currently limited experience with over dosage of Mupirocin.

**Treatment:**

There is no specific treatment for an overdose of Mupirocin. In the event of overdose, the patient should be treated supportively with appropriate monitoring as necessary. Further management should be as clinically indicated or as recommended by the national poisons centre, where available.

**STORAGE CONDITIONS:**

Store in dry place, temperature below 30°C. Protect from light. Do not freeze.

Keep all medicines out of reach of children.

**DOSAGE FORMS AND PACKING AVAILABLE:**

Alu. tube of 5 g (Box of 1's)

Alu. tube of 15 g (Box of 1's)

**NAME AND ADDRESS OF MANUFACTURER:**

Kusum Healthcare Pvt. Ltd.

SP-289(A), RIICO Industrial Area, Chopanki,

Bhiwadi, Dist. Alwar (Rajasthan), INDIA

**PRODUCT REGISTRATION HOLDER:**

**Synerrv Sdn Bhd,**

SO-29-2, Menara 1, KL Eco City, Jalan Bangsar, KG

Haji Abdullah Hukum, 59200 Kuala Lumpur Wilayah

Persekutuan Kuala Lumpur, Malaysia

**DATE OF REVISION OF PACKAGE INSERT:**

October 2020