

pharmaniaga®

ALLERYL TABLET 4MG

DESCRIPTION

Clean, dry, round, flat, bevel-edged, scored on one side and pink tablet.

Each tablet contains Chlorpheniramine Maleate 4mg

Source of lactose: Bovine

ROUTE OF ADMINISTRATION

Oral

PHARMACODYNAMICS

Chlorpheniramine is a potent antihistamine (H1-antagonist).

Antihistamines diminish or abolish the actions of histamine in the body by competitive reversible blockade of histamine H1-receptor sites on tissues. Chlorpheniramine also has anticholinergic activity.

Antihistamines act to prevent the release of histamine, prostaglandins and leukotrienes and have been shown to prevent the migration of inflammatory mediators. The actions of chlorpheniramine include inhibition of histamine on smooth muscle, capillary permeability and hence reduction of oedema and wheal in hypersensitivity reactions such as allergy and anaphylaxis.

PHARMACOKINETICS

Chlorpheniramine is well absorbed from the gastro-intestinal tract, following oral administration. The effects develop within 30 minutes, are maximal within 1 to 2 hours and last 4 to 6 hours. The plasma half-life has been estimated to be 12 to 15 hours.

Chlorpheniramine is metabolised to the monodesmethyl and didesmethyl derivatives. About 22% of an oral dose is excreted unchanged in the urine.

INDICATIONS

This medicine is indicated for symptomatic control of all allergic conditions responsive to antihistamines, including hay fever, vasomotor rhinitis, urticaria, angioneurotic oedema, food allergy, drug and serum reactions, insect bites and itch associated with chickenpox.

Temporary relief of nasal and sinus congestion due to the common cold in combination with a decongestant (pseudoephedrine).

RECOMMENDED DOSAGE

Child 6-11 years: 2mg every 4 – 6 hours.
Maximum daily dose: 12mg per day.

Child 12-17 years: 4mg every 4-6 hours.
Maximum daily dose: 24mg per day.

Adult: 4mg every 4-6 hours.
Maximum daily dose: 24mg per day.

Elderly: 4mg every 4 – 6 hours.
Maximum daily dose: 12mg per day.

CONTRAINDICATIONS

Alleryl Tablet 4mg is contra-indicated in patients who are hypersensitive to antihistamines or to any of the tablet ingredients.

The anticholinergic properties of chlorpheniramine are intensified by monoamine oxidase inhibitors (MAOIs). Alleryl Tablet 4mg is therefore contra-indicated in patients who have been treated with MAOIs within the last fourteen days.

WARNINGS & PRECAUTIONS

Chlorpheniramine maleate in common with other drugs having anticholinergic effects, should be used with caution in epilepsy; raised intra-ocular pressure including glaucoma; prostatic hypertrophy; severe hypertension or cardiovascular disease; bronchitis, bronchiectasis or asthma; hepatic impairment; renal impairment. Children and the elderly are more likely to experience the neurological anticholinergic effects and paradoxical excitation (e.g. increased energy, restlessness, nervousness).

The anticholinergic properties of chlorpheniramine may cause drowsiness, dizziness, blurred vision and psychomotor impairment in some patients which may seriously affect ability to drive and use machinery.

The effects of alcohol may be increased and therefore concurrent use should be avoided.

Should not be used with other antihistamine containing products, including antihistamine containing cough and cold medicines.

Patients with rare hereditary problems of galactose intolerance, Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicine.

INTERACTIONS WITH OTHER MEDICAMENTS

Concurrent use of chlorpheniramine and hypnotics or anxiolytics may cause an increase in sedative effects, therefore medical advice should be sought before taking chlorpheniramine concurrently with these medicines.

Chlorpheniramine inhibits phenytoin metabolism and can lead to phenytoin toxicity.

The anticholinergic effects of chlorpheniramine are intensified by MAOIs.

PREGNANCY & LACTATION

Pregnancy:

There are no adequate data for this medicine used in pregnant women. The potential risk for humans is unknown. Use during the third trimester may result in reactions in the newborn or premature neonates.

Lactation:

This medicine and other antihistamine may inhibit lactation and may be secreted in breast milk.

Not to be used during pregnancy and lactation unless considered essential by a physician.

SIDE/ADVERSE EFFECTS

Blood and lymphatic system disorders:-

Unknown: haemolytic anaemia, blood dyscrasias.
Agranulocytosis, thrombocytopenia, pancytopenia and aplastic anaemia.

Immune system disorders:-

Unknown: allergic reaction, angioedema, anaphylactic reactions

Metabolism and nutritional disorders:-

Unknown: anorexia

Psychiatric disorders:-

Unknown: confusion*, excitation*, irritability*, nightmares*, depression.

Nervous system disorders*:-

Very common: sedation, somnolence
Common: disturbance in attention, abnormal coordination, dizziness headache

Eye Disorders:-

Common: blurred vision

Ear and labyrinth disorders:-

Unknown: tinnitus

Cardiac disorders:-

Unknown: palpitations, tachycardia, arrhythmias

Vascular disorders:-

Unknown: Hypotension

Respiratory, thoracic and mediastinal disorders:-

Unknown: thickening of bronchial secretions

Gastrointestinal disorders:-

Common: nausea, dry mouth
Unknown: vomiting, abdominal pain, diarrhoea, dyspepsia

Hepatobiliary disorders:-

Unknown: hepatitis, including jaundice

Skin and subcutaneous disorders:-

Unknown: exfoliative dermatitis, rash, urticaria, photosensitivity

Musculoskeletal and connective tissue disorders:-

Unknown: muscle twitching, muscle weakness

Renal and urinary disorders:-

Unknown: urinary retention

General disorders and administration site conditions:-

Common: fatigue
Unknown: chest tightness

*Children and the elderly are more likely to experience the neurological anticholinergic effects and paradoxical excitation (eg. increased energy, restlessness, nervousness).

SYMPTOMS & TREATMENT FOR OVERDOSAGE

Symptoms:

The estimated lethal dose of chlorpheniramine maleate is 25 to 50mg/kg body weight. Symptoms and signs include sedation, paradoxical excitation of the CNS, toxic psychosis, convulsions, apnoea, anticholinergic effects, dystonic reactions and cardiovascular collapse including arrhythmias.

Treatment:

Symptomatic and supportive measures should be provided with special attention to cardiac, respiratory, renal and hepatic functions and fluid and electrolyte balance. If over-dosage is by the oral route, treatment with activated charcoal should be considered provided there are no contraindications for use and the overdose has been taken recently (treatment is most effective if given within an hour of ingestion). Treat hypotension and arrhythmias vigorously. CNS convulsions may be treated with diazepam. Haemoperfusion may be used in severe cases.

EFFECTS ON ABILITY TO DRIVE & USE MACHINE

This medicine may cause drowsiness, dizziness, blurred vision and psychomotor impairment, which can seriously hamper the patients' ability to drive and use machinery.

PACK SIZES

Blister pack of 10 tablets:

- i) 10 strips of 10 tablets
- ii) 100 strips of 10 tablets

STORAGE CONDITIONS

Keep in a dry place below 30°C; Protect from light.

INSTRUCTIONS FOR USE

Do not exceed the recommended dose or frequency of dosing.

SHELF LIFE

Product should not be used beyond the expiry date imprinted on the product packaging.

REGISTRATION NUMBER

MAL19860094AZ

CONTROLLED MEDICINE UBAT TERKAWAL

KEEP MEDICINES OUT OF REACH OF CHILDREN JAUHI UBAT-UBATAN DARI KANAK-KANAK

For further information, please consult your doctor or your pharmacist.

Revision Number : 08

Revision Date : 18 Sep 2023

Lt-001A.01 Product Registration Holder and Manufacturer
IDAMAN PHARMA MANUFACTURING SDN BHD (661901-P)
LOT 24 & 25, JALAN PERUSAHAAN LAPAN,
BAKAR ARANG INDUSTRIAL ESTATE, 08000 SUNGAI PETANI
KEDAH DARUL AMAN, MALAYSIA.