

Important information. Please read carefully.

AntaminTM

Tablets

Composition

Each tablet contains Chlorpheniramine Maleate 4 mg

Properties

Antamin preparations contain chlorpheniramine maleate, a potent antihistamine which has antagonistic action against histamine. It is readily absorbed with quick action starting 15 to 30 minutes after ingestion.

Indications

Antamin is useful for the symptomatic treatment of allergic conditions such as seasonal rhinitis (hay fever), acute urticaria, angioedema, food allergies, insect bites, pruritus, atopic and contact dermatitis.

Dosage and Administration

To be administered orally.

Adults : 4 mg three to four times daily;
Children 6 to 12 years : 2 mg three or four times daily;
2 to 5 years : 1 to 2 mg three times daily;
1 to 2 years : 1 mg twice daily;
Below 1 year : as prescribed by the physician

Contraindications

In patients with known hypersensitivity to chlorpheniramine.

Precautions

As drowsiness may occur, patients taking this medication should be advised not to drive or operate machinery and to avoid alcoholic drinks. **Antamin** should be used with care in patients with glaucoma, prostatic hypertrophy, epilepsy and hepatic disease. Like all other medicines, the use of chlorpheniramine should be avoided during pregnancy unless absolutely necessary and under close medical supervision.

Drug Interactions

Avoid concurrent use with tranquillisers and anticholinergic drugs (e.g. benzhexol) as their effects and side effects may be potentiated. Chlorpheniramine has an antagonistic action towards betahistine.

Side Effects

Although Chlorpheniramine is less sedating compared with some other antihistamines, drowsiness may occur in susceptible patients. Other side effects that may occur include dizziness, headaches, insomnia, nervousness, anticholinergic effects such as dry mouth, throat and respiratory passages, blurred vision and gastrointestinal disturbances.

Overdosage

Hospital admission is strongly advised for patients showing symptoms of overdosage. Symptoms include both central depressant and stimulant effects such as severe drowsiness and coma followed by convulsions and postictal depression.

In small children the dominant effect is excitation; other effects include hallucinations, incoordination, convulsions, fixed dilated pupils, flushed face and fever.

Treatment is along general symptomatic and supportive lines.

Aspiration and gastric lavage for overdosage. Convulsions may be controlled by diazepam intra-venously. Mechanical ventilation and fluid replacement may also be required.

Shelf-life

The expiry date is indicated on the packaging.

Presentation

Yellow, round tablets in blister packs of 100's and 1000's.

Storage

Store below 30°C. Protect from light.

**KEEP OUT OF REACH OF CHILDREN
JAUHI DARI KANAK-KANAK**

For further information, please consult your pharmacist or physician.

Revision Date : 10-Jun-2020

Manufacturer and Product Registration Holder
Xepa-Soul Pattinson (Malaysia) Sdn Bhd
1-5 Cheng Industrial Estate, 75250 Melaka, Malaysia

