

pharmaniaga®

ALLERYL SYRUP 4MG/5ML

DESCRIPTION

Colourless to yellowish clear syrup, sweet, pleasant and slightly sourish after taste.

COMPOSITION

Each 5mL of the syrup contains:

Chlorpheniramine Maleate 4 mg

Preservative: Sodium Benzoate 5 mg

PHARMACODYNAMICS

ATC code: R06AB02

Chlorpheniramine Maleate is a potent antihistamine (H1-antagonist). Antihistamines diminish or abolish the actions of histamine in the body by competitive reversible blockade of histamine H1-receptor sites on tissues. Chlorpheniramine Maleate also has anticholinergic activity.

Antihistamines prevent the migration of inflammatory mediators and the release of histamine, prostaglandins and leukotrienes. The actions of Chlorpheniramine Maleate include inhibition of histamine on smooth muscle, capillary permeability and reduction of oedema and wheal in hypersensitivity reactions such as allergy and anaphylaxis.

PHARMACOKINETICS

Following oral administration, Chlorpheniramine Maleate is well absorbed from the gastro-intestinal tract. The effects develop within 30 minutes, are maximal within 1 to 2 hours and last 4 to 6 hours. The plasma half-life is estimated to be 12 to 15 hours. Chlorpheniramine Maleate is metabolised to monodesmethyl and didesmethyl derivatives. About 22% of an oral dose is excreted unchanged in the urine.

INDICATIONS

- 1) Symptomatic treatment of perennial and seasonal allergic rhinitis, vasomotor rhinitis, antihistamines including hay fever and allergic conjunctivitis due to inhalant allergens and foods.
- 2) Symptomatic treatment of pruritus associated with allergic reactions and of mild, uncomplicated allergic skin manifestations of urticaria and angioedema.
- 3) May be used in combination with a decongestant (Pseudoephedrine) for the temporary relief of nasal and sinus congestion associated with the common cold and allergic rhinitis.

RECOMMENDED DOSAGE

Adults and children 12 years and over: 5ml (4mg) every 4 to 6 hourly. Maximum daily dose: 30ml (24mg) in any 24 hours.

Elderly: The elderly are more likely to experience neurological anticholinergic effects. Consideration should be given to using a lower daily dose (e.g. a maximum of 12 mg in any 24 hours).

Children aged 6 - 12 years: 2.5ml (2mg) every 4 to 6 hourly. Maximum daily dose: 15ml (12mg) in any 24 hours.

Children aged 2 - 5 years: 1.25ml (1mg) every 4 to 6 hourly. Maximum daily dose: 7.5ml (6mg) in any 24 hours

Do not exceed the stated dose or frequency of dosing.

The minimum interval between the doses should be 4 hours.

Do not use continuously for more than two weeks without consulting a doctor.

ROUTE OF ADMINISTRATION

Oral

CONTRAINDICATIONS

Alleryl Syrup 4mg/5ml is contraindicated in patients with known hypersensitivity to Chlorpheniramine Maleate or to any of the syrup ingredients.

The anticholinergic properties of Chlorpheniramine Maleate are intensified by monoamine oxidase inhibitors (MAOIs). Alleryl Syrup 4mg/5ml is therefore contraindicated in patients who have been treated with MAOIs within the last fourteen days.

WARNINGS AND PRECAUTIONS

When used in the treatment of cough and cold:

- a. Do not use for children below 2 years of age.
- b. Use with caution on the advice of doctor / pharmacist for children aged 2 - 6 years.

Chlorpheniramine Maleate, in common with other drugs having anticholinergic effects, should be used with caution in epilepsy; raised intra-ocular pressure including glaucoma; prostatic hypertrophy; severe hypertension or cardiovascular disease; bronchitis, bronchiectasis or asthma; hepatic impairment; renal impairment. Children and the elderly are more likely to experience the neurological anticholinergic effects and paradoxical excitation (e.g., increased energy, restlessness, nervousness). Avoid use for elderly patients experiencing confusion.

The anticholinergic properties of Chlorpheniramine Maleate may cause drowsiness, dizziness, blurred vision and psychomotor impairment in some patients which may seriously affect ability to drive and use machinery.

The effects of alcohol may be increased and therefore concurrent use should be avoided.

Should not be used with other antihistamine containing products, including antihistamine containing cough and cold medicines.

Concurrent use with drugs which cause sedation such as anxiolytics and hypnotics may cause an increase in sedative effects, therefore medical advice should be sought before taking chlorpheniramine concurrently with these medicines.

Patients with rare hereditary problems of fructose intolerance, glucose-galactose malabsorption or sucrose-isomaltase insufficiency should not take this medicine.

Alleryl Syrup 4mg/5ml contains 3 g of sucrose per 5 ml. This should be taken into account in patients with diabetes mellitus.

Long term use increases the risk of dental caries and it is essential that adequate dental hygiene is maintained.

INTERACTIONS WITH OTHER MEDICAMENTS

Concurrent use of chlorphenamine and hypnotics or anxiolytics may cause an increase in sedative effects, concurrent use of alcohol may have a similar effect therefore medical advice should be sought before taking chlorphenamine concurrently with these medicines.

Chlorphenamine inhibits phenytoin metabolism and can lead to phenytoin toxicity.

The anticholinergic effects of chlorphenamine are intensified by MAOIs.

PREGNANCY AND LACTATION

Pregnancy

There are no adequate data from the use of chlorphenamine in pregnant women. The potential risk for humans is unknown, use during the third trimester may result in reactions in the new-born or premature neonates. Not to be used during pregnancy unless considered essential by a physician.

Lactation

Chlorphenamine maleate and other antihistamines may inhibit lactation and may be secreted in breast milk. Alleryl Syrup 4mg/5ml is not to be used during lactation unless considered essential by a physician.

ADVERSE EFFECTS AND UNDESIRABLE EFFECTS

System Organ Class	Adverse Reaction	Frequency
Nervous system disorders*	Sedation, somnolence	Very common
	Disturbance in attention, abnormal coordination, dizziness headache	Common
Eye disorders	Blurred Vision	Common
Gastrointestinal disorders	Nausea, dry mouth	Common
	Vomiting, abdominal pain, diarrhoea, dyspepsia	Unknown
Immune system disorders:	Allergic reaction, angioedema, anaphylactic reactions	Unknown
Metabolism and nutritional disorders	Anorexia	Unknown
Blood and lymphatic system disorders	Haemolytic anaemia, blood dyscrasias	Unknown
Musculoskeletal and connective tissue disorders	Muscle twitching, muscle weakness	Unknown
Psychiatric disorders	Confusion*, excitation*, irritability*, nightmares*, depression	Unknown
Renal and urinary disorders	Urinary retention	Unknown
Skin and subcutaneous disorders	Exfoliative dermatitis, rash, urticaria, photosensitivity	Unknown
Respiratory, thoracic and mediastinal disorders	Thickening of bronchial secretions	Unknown
Vascular disorders	Hypotension	Unknown

Hepatobiliary disorders	Hepatitis, including jaundice	Unknown
Ear and labyrinth disorders	Tinnitus	Unknown
Cardiac disorders	Palpitations, tachycardia, arrhythmias	Unknown
General disorders and administration site conditions	Fatigue	Common
	Chest tightness	Unknown

**Children and the elderly are more susceptible to neurological anticholinergic effects and paradoxical excitation (e.g., increased energy, restlessness, nervousness)*

OVERDOSAGE

Symptom

The estimated lethal dose of Chlorpheniramine Maleate is 25 to 50 mg/kg body weight. Symptoms and signs include sedation, paradoxical excitation of the CNS, toxic psychosis, convulsions, apnoea, anticholinergic effects, dystonic reactions and cardiovascular collapse including arrhythmias.

Treatment

Symptomatic and supportive measures should be provided with special attention to cardiac, respiratory, renal and hepatic functions and fluid and electrolyte balance. If overdosage is by the oral route, treatment with activated charcoal should be considered, provided there are no contraindications for use and the overdose has been taken recently (treatment is most effective if given within an hour of ingestion). Treat hypotension and arrhythmias vigorously. CNS convulsions may be treated with i.v.diazepam. Haemoperfusion may be used in severe cases.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINE

The anticholinergic properties of Chlorpheniramine Maleate may cause drowsiness, dizziness, blurred vision and psychomotor impairment, which can seriously hamper the patients' ability to drive and use machinery.

PACKING / PACK SIZE

Plastic container of 60ml.

STORAGE CONDITIONS

Keep container tightly closed in a dry place, below 30°C; Protect from light.

SHELF LIFE

Product should not be used beyond the expiry date imprinted on the product packaging.

REGISTRATION NUMBER

MAL19860099AZ.

CONTROLLED MEDICINE UBAT TERKAWAL

KEEP MEDICINES OUT OF REACH OF CHILDREN JAUHI UBAT-UBATAN DARI KANAK-KANAK

For further information, please consult your doctor or your pharmacist.

Revision Number : 01

Revision Date : 19 Dec 2025

LT-305.01 PRODUCT REGISTRATION HOLDER AND MANUFACTURER
INSPIRAPHARMA SDN. BHD. (200401023395)
LOT 24 & 25, JALAN 8,
KAWASAN PERUSAHAAN BAKAR ARANG,
08000 SUNGAI PETANI,
KEDAH DARUL AMAN, MALAYSIA.