

BEAVATE CREAM 0.1%



DESCRIPTION

A white coloured, homogenous cream containing
Betamethasone 17 -valerate eq to Betamethasone 0.1% w/w
& Preservatives: Methylparaben 0.32% w/w, Propylparaben 0.048% w/w.

INDICATIONS

Beavate cream is indicated for the treatment of mild to moderate atopic dermatitis, contact dermatitis, mild mummular dermatitis, seborrheic, facial and dematitis of the intertriginous areas; other mild to moderate inflammatory dermatoses and other forms of dermatitis; intertrigo, facial and intertriginous linchen planus, discoid lupus erythematosus, polymorphous light eruption; anogenital pruritus, pruritus senilis, psoriasis, and for treatment during the inflammatory phase of xerosis.

PHARMACODYNAMICS

Corticosteroids diffuse through cell membranes and complexes with specific cytoplasmic cell receptors. These complexes then enters the cell nucleus, binds to DNA, and stimulate transcription of mRNA to produce various inhibitory enzymes responsible for the anti-inflammatory effects of the corticosteroids. These inflammatory effects include inhibition of early process such as oedema, fibri deposition, capillary dilatation, movement of phagocytes into the area and phagocytic activities. Later process such as capillary production, collagen deposition, and keloid formation are also inhibited by corticosteroids. The overall action of corticosteroids is catabolic.

PHARMACOKINETICS

Beavate cream is absorbed systemically through the stratum corneum. The penetration process is enhanced by increasing skin hydration and/or temperature. Occlusion also enhances penetration. Increased percutaneous absorption also occurs when the mucosa or skin is abraded or inflamed; when body temperature is elevated; with prolonged use and/or with extensive use. Repeated application results in a cumulative depot effect in the skin, which may lead to prolonged duration of action, increased side effects, and increased systemic absorption.

DOSAGE

Adult

Apply Beavate cream thinly to the affected area once to three times a day.

Paediatric

Apply Beavate cream thinly to the affected area once a day.

SIDE EFFECTS

Beavate cream may cause side-effects that may warrant medical attention such as allergic contact dermatitis, folliculitis, furunculosis, pustules, pyoderma, vesiculation, hyperesthesia, numbness in fingers, purpura, skin atrophy, secondary skin infection, stripping of epidermal layer and telangiectasia. Less frequent side-effects are acneiform eruptions, posterior subcapsular cataracts, Cushing's syndrome, perioral dermatitis, ecchymosis, oedema, gastric ulcer, secondary glaucoma, hirsutism, hypertension, hypokalaemia syndrome, hypopigmentation, miliaria rubra, protein depletion, skin laceration, skin maceration, striae, subcutaneous tissue atrophy and unusual loss of hair.

CONTRAINDICATIONS

Beavate cream is contraindicated in untreated pathogenic skin lesion, rosacea (acne rosacea), perioral dermatitis, acne vulgaris, allergic to corticosteroids, pre-existing skin atrophy.

PRECAUTIONS

Patients who are allergic to corticosteroids, pregnant, lactating, young children, and the elderly should exercise caution when using Beavate cream.

OVERDOSAGE

Discontinue occlusive dressing (if used) and institute appropriate antimicrobial therapy until the infection is properly controlled should local infection occurs. Discontinuance of Beavate cream may be necessary. If irritation or sensitization occurs at the site of application, discontinue Beavate cream and initiate proper symptomatic treatment. For chronic topical overdosage, gradual withdrawal may be necessary since there is no specific antidote available. For acute oral overdosage, dilution with fluid may all be necessary as there is no specific antidote and serious adverse effects are most unlikely.

Up-to-date information on treatment of overdose can be obtained from The National Poison Centre, Universiti Sains Malaysia.

PRESENTATION

15g, 30g aluminium collapsible tube, 30g x50's
500g polypropylene (PP) jar.
Not all pack sizes are available locally.

STORAGE CONDITIONS AND USER INSTRUCTIONS

Keep container tightly closed.
Store in a dry place below 30°C.
Keep out of reach of children.
Jauhi daripada kanak-kanak.
Shelf life : Please refer to outer package.
Keep away from eyes.
Occlusive dressing in the affected area only if prescribed by your general practitioner.
Gradual withdrawal may be necessary to avoid rebound flare-up in certain conditions, e.g., psoriasis.
For external use only.

Product Registration Holder:

DUOPHARMA MANUFACTURING (BANGI) SDN. BHD.
Lot No. 2, 4, 6, 8 & 10, Jalan P/7, Section 13,
Bangi Industrial Estate, 43650 Bandar Baru Bangi,
Selangor, Malaysia.

Manufacturer:

DUOPHARMA MANUFACTURING (BANGI) SDN. BHD.
Lot No. 2 & 4, Jalan P/7, Section 13,
Bangi Industrial Estate, 43650 Bandar Baru Bangi,
Selangor, Malaysia.



150001XXXX XX
Revision Date: Jan 2021

BEAVATE CREAM