

HYSONE CREAM 1% (w/w)

DESCRIPTION

A white coloured, homogeneous cream containing Hydrocortisone Acetate 1%.
Preservatives: Methylparaben 0.32% w/w, Propylparaben 0.048% w/w.

INDICATION

For relief of symptoms of acute and chronic skin dermatoses, e.g. Eczema, contact dermatitis and psoriasis.

PHARMACODYNAMICS

Corticosteroids diffuse through cell membranes and complexes with specific cytoplasmic cell receptors. These complexes then enters the cell nucleus, binds to DNA, and stimulate transcription of mRNA to produce various inhibitory enzymes responsible for the anti-inflammatory effects of the corticosteroids. These inflammatory effects includes inhibition of early process such as oedema, fibrin deposition, capillary dilatation, movement of phagocytes into the area and phagocytic activities. Later process such as capillary production, collagen deposition, and keloid formation are also inhibited by corticosteroids. The overall action of corticosteroids is catabolic.

PHARMACOKINETICS

Hysone Cream 1% (w/w) is absorbed systemically through the stratum corneum. The penetration process is enhanced by increasing skin hydration and/or temperature. Occlusion also enhances penetration. Increased percutaneous absorption also occurs when the mucosa or skin is abraded or inflamed; when body temperature is elevated; with prolonged use and/or with extensive use. Repeated application results in a cumulative depot effect in the skin, which may lead to prolonged duration of action, increased side effects, and increased systemic absorption.

DOSAGE

A small quantity should be applied to the affected area 2 or 3 times daily until improvement occurs. It may then be possible to maintain improvement by applying once a day, or even less often. Occlusion of the area under treatment with plastic film may be used to enhance the effect in more resistant cases.

Route of Administration: For External use only

SIDE EFFECT

Prolonged and intensive treatment, particularly with occlusive dressing, may cause local atrophic changes in the skin, such as striae and may lead to secondary infections. Other minor side effects are itching, dryness, irritation eruptions and burning sensations.

CONTRAINDICATIONS

Hysone Cream 1% (w/w) is contraindicated in untreated pathogenic skin lesion, rosacea (acne rosacea), perioral dermatitis, acne vulgaris, allergic to corticosteroids, pre-existing skin atrophy.



PRECAUTIONS

Long-term or prophylactic use, rebound following sudden withdrawal; fluorinated corticosteroids are more likely to cause systemic effects or damage delicate tissues.

OVERDOSAGE

Discontinue occlusive dressing (if used) and institute appropriate antimicrobial therapy until infection is properly controlled should local infection occurs. Discontinuance of Hysone Cream 1% (w/w) may be necessary. If irritation or sensitization occurs at the site of application, discontinue Hysone Cream 1% (w/w) and initiate proper symptomatic treatment. For chronic topical overdosage, gradual withdrawal may be necessary since there is no specific antidote available. For acute overdosage, dilution with fluid may all be necessary as there is no specific antidote and serious adverse effects are most unlikely.

Up-to-date information on treatment of overdose can be obtained from The National Poison Centre. Universiti Sains Malaysia.

PRESENTATION

15g, 15g x 50's and 450gm

STORAGE CONDITIONS AND USER INSTRUCTIONS

Keep container tightly closed.

Store in a dry place below 30°C.

Protect from light.

Keep out of reach of children.

Jauhi daripada kanak-kanak.

Shelf-life: Please refer to outer package.

Product Registration Holder:

DUOPHARMA MANUFACTURING (BANGI) SDN. BHD.

Lot No. 2, 4, 6, 8 & 10, Jalan P/7, Section 13,
Bangi Industrial Estate, 43650 Bandar Baru Bangi,
Selangor, Malaysia.

Manufacturer:

DUOPHARMA MANUFACTURING (BANGI) SDN. BHD.

Lot No. 2 & 4, Jalan P/7, Section 13,
Bangi Industrial Estate, 43650 Bandar Baru Bangi,
Selangor, Malaysia.



150001XXXX XX
Revision Date: August 2022

HYSONE CREAM 1% (w/w)