

———— IDAMAN PHARMA MANUFACTURING ————

IDAMAN PHARMA DIAZEPAM TABLET 5MG

PRODUCT DESCRIPTION

Clean, dry, uniform, round, flat, bevel edged tablets and scored on one side. Colour: Yellow.

Each tablet contains Diazepam 5mg

Source of lactose: Bovine

ROUTE OF ADMINISTRATION: Oral

PHARMACODYNAMICS

Mechanism of Action: In general, diazepam act as depressants of the central nervous system (CNS), producing all levels of CNS depression from mild sedation to hypnosis to coma depending on dose.

The precise sites and mechanisms of action have not been completely established. Although various mechanisms of action have been proposed, it is believed that benzodiazepines advance or facilitate the inhibitory neurotransmitter action of gammaaminobutyric acid (GABA), which mediates both pro-and post-synaptic inhibition in all regions of the CNS, following interaction between diazepam and a specific neuronal membrane receptor.

Antianxiety Agent: Sedative-hypnotic Believed to stimulate GAMA receptors in the ascending reticular activating system. Since GAMA is inhibitory, receptor stimulation increases inhibitions and blocks both cortical and limbic arousal following stimulation of the brain stem formation.

Anticonvulsant: Appear to act, at least partially, by enhancing presynaptic inhibition. Suppress the spread of seizure activity produced by epileptogenic foci in the cortex, thalamus and limbic structures but do not abolish the abnormal discharge of the focus.

Skeletal Muscle Relaxant: The exact mechanism of action of diazepam has not been completely established but it appears to produce skeletal muscle relaxation primarily by inhibiting spinal polysynaptic afferent pathways; however, it may also inhibit monosynaptic afferent pathways. Diazepam may inhibit monosynaptic and polysynaptic reflexes by acting as an inhibitory neuronal transmitter or by blocking excitatory synaptic transmission. It may also directly depress motor nerve and muscle function.

Antitremor Agent: Mechanism of action is not known.

PHARMACOKINETICS

Diazepam is a benzodiazepine tranquiliser with anticonvulsant, sedative, muscle relaxant, anti-tremor and amnesia properties. Diazepam is readily and completely absorbed from the gastrointestinal tract, peak plasma concentrations occurring within about 30 to 90 minutes of oral administration. The rate of absorption is age-related and tend to be delayed in the elderly. Diazepam crosses the blood-brain barrier and is highly lipid soluble. Initial effects of diazepam decrease rapidly as it is redistributed into fat depots and tissues. Diazepam has a biphasis half-life with an initial rapid distribution phase followed by a prolonged terminal elimination phase of 1 or 2 days. Diazepam is extensively metabolized in the liver, in addition to desmethyldiazepam, its active metabolites include methyloxazepam and temazepam. It is excreted in the urine, mainly in the form of its metabolites. There is no biliary excretion. Diazepam is very extensively

bound to plasma protein. The plasma half-life of diazepam is prolonged in neonates and in the elderly and in patients with kidney or liver diseases. In addition to crossing the blood-brain barrier, diazepam and its metabolites also crosses the placental barrier and are excreted in breast milk.

INDICATIONS

- 1) Treatment of anxiety.
- 2) Treatment of acute alcohol withdrawal symptoms.
- 3) Indicated for preoperative procedures to relieve anxiety and tension.
- 4) Indicated as an adjunct prior to endoscopic procedures.
- 5) Indicated as short-term (7-14 days) adjunctive therapy in convulsive disorder.
- 6) Indicated as an adjunct for the treatment of skeletal muscle spasm.
- 7) Treatment of familial, senile or essential action tremors.

RECOMMENDED DOSAGE

Usual Adult Dose:

Antianxiety Agent - Orally, 2 to 10mg, two to four times a day.

Sedative-Hypnotic-Acute Alcohol Withdrawal:

Orally, 10mg, three or four times during the first 24 hours, reduces to 5mg, three or four times a day as needed.

Anticonvulsant: Orally, 2 to 10mg, three or four times a day.

Skeletal Muscle Relaxant: Orally, 2 to 10mg, three or four times a day.

Note: Geriatric or debilitated patients - Orally, 2 to 2.5mg one or two times a day, increase gradually as needed and tolerated.

Usual Pediatric Dose:

Children Up To 6 Months of Age: Use is not recommended.

Children 6 Months Of Age And Over: Orally, 1 mg to 2.5 mg, three or four times a day initially. Because of varied responses to CNS-acting drugs, initiate therapy with lowest dose and increase gradually as needed and tolerated.

CONTRAINDICATIONS

- 1) In patients with acute-close angle glaucoma.
- 2) In patients with psychotic disorder.
- 3) In patients with myasthenia gravis.
- 4) In patients with severe chronic obstructive disease.

WARNING & PRECAUTIONS

- 1) Use diazepam with caution in paediatrics; in elderly and debilitated patients; in patients with arteriosclerosis, renal or respiratory dysfunction.
- 2) Withdrawal of diazepam in patients who have been receiving high dosage should be gradual.
- 3) This preparation may be habit forming on prolonged use and may cause drowsiness, avoid alcoholic beverages.
- 4) Anaphylaxis (severe allergic reaction) and angioedema (severe facial swelling) which can occur as early as the first time the product is taken.
- 5) Complex sleep - related behaviours which may include sleep driving, making phone calls, preparing and eating food (while asleep) may occur.

Risk from Concomitant Use with Opioids

Profound sedation, respiratory depression, coma, and death may result from the concomitant use of Idaman Pharma Diazepam Tablet 5mg with opioids. Observational studies have demonstrated that concomitant use of opioids and benzodiazepines increases the risk of drug-related mortality compared to use of opioids alone. Because of

these risks, reserve concomitant prescribing of these drugs for use in patients for whom alternative treatment options are inadequate.

If the decision is made to newly prescribe a benzodiazepine and an opioid together, prescribe the lowest effective dosages and minimum durations of concomitant use.

If the decision is made to prescribe a benzodiazepine in a patient already receiving an opioid, prescribe a lower initial dose of the benzodiazepine than indicated in the absence of an opioid, and titrate based on clinical response.

If the decision is made to prescribe an opioid in a patient already taking a benzodiazepine, prescribe a lower initial dose of the opioid, and titrate based on clinical response.

Follow patients closely for signs and symptoms of respiratory depression and sedation. Advise both patients and caregivers about the risks of respiratory depression and sedation when Idaman Pharma Diazepam Tablet 5mg is used with opioids. Screen patients for risk of substance use disorders, including opioid abuse and misuse, and warn them of the risk for overdose and death associated with the use of opioids.

INTERACTIONS WITH OTHER MEDICAMENTS

Be alert for the possible drug interactions between diazepam and antacids, carbamazepine, cimetidine, CNS depressants, Isoniazid, levodopa, rifampicin, scopolamine, MAO inhibitors, tricyclic antidepressants, fentanyl derivative, phenytoin and valproic acid. Incompatible with alcohol.

Opioids

Due to additive pharmacologic effect, the concomitant use of opioids with benzodiazepines increases the risk of respiratory depression, profound sedation, coma and death.

The concomitant use of opioids and benzodiazepines increases the risk of respiratory depression because of actions at different receptor sites in the central nervous system that control respiration. Opioids interact primarily at μ -receptors, and benzodiazepines interact at GABA_A sites. When opioids and benzodiazepines are combined, the potential for benzodiazepines to significantly worsen opioid-related respiratory depression exists.

Reserve concomitant prescribing of these drugs for use in patients for whom alternative treatment options are inadequate.

Limit dosage and duration of concomitant use of benzodiazepines and opioids, and follow patients closely for respiratory depression and sedation.

PREGNANCY AND LACTATION

Pregnancy

Diazepam crosses the placenta. Diazepam have been reported to increase the risk of congenital malformations when used during the first trimester. Chronic usage during pregnancy may cause physical dependence with resulting withdrawal symptoms in the neonate. Usage during last weeks of pregnancy (especially with hypnotic doses) may result in neonatal CNS depression. Thus, use of diazepam during pregnancy should be carefully considered and should be used only if the expected benefit outweighs any potential risk.

Lactation

Diazepam and its metabolites are secreted in breast milk. Since neonates metabolise diazepam more slowly than adult, accumulation of diazepam and / or its metabolites may occur. Use by nursing mothers may cause sedation and possible feeding difficulties and weight loss in the infant. Hence risk-benefit must be considered.

SIDE/ADVERSE EFFECTS

Side effects are usually mild and infrequent, drowsiness, light-headedness and ataxia, are the most common and are often dose-dependent. Elderly and debilitated patients are especially susceptible to these reactions. Diazepam may induce amnesia. Other effects include hypotension, some respiratory depression, nausea and constipation, changes in salivation, blurred vision and diplopia, dysarthria, skin rashes, urinary retention, incontinence, mental depression, tremor, headache, confusion, slurred speech, vertigo and changes in libido. Prolonged use of diazepam may lead to the development of dependence and will produce withdrawal reaction when withdrawn abruptly.

CAUTION: This preparation may be habit forming on prolonged use.

SYMPTOMS & TREATMENT OVERDOSE

Symptoms:

Include confusion (continuing), severe drowsiness, shakiness, continuing slurred speech, staggering, unusual slow heartbeat, shortness of breath, or trouble breathing and severe weakness.

Treatment:

There are no specific treatments, if recent severe overdosage the stomach should be emptied by aspiration and lavage. The rest of treatment is symptomatic.

EFFECTS ON ABILITY TO DRIVE & USE MACHINE

Sedation, amnesia and impaired muscular function may adversely affect the ability to drive or use machines. If insufficient sleep occurs, the likelihood of impaired alertness may be increased. Impaired function and sedation may occur the following morning and for several days after.

Patients should be warned that effects on the central nervous system may persist into the day after administration even after a single dose.

This medicine can impair cognitive function and can affect a patient's ability to drive safely. When prescribing this medicine, patients should be told:

- The medicine is likely to affect your ability to drive
- Do not drive until you know how the medicine affects you
- It is an offence to drive while under the influence of this medicine

PACK SIZES

- Plastic Containers of 30's
- Blister Packs of 10's (100 Strips X 10's).

STORAGE CONDITION

Keep container tightly closed in a dry place, below 30°C; Protect from light.

INSTRUCTIONS FOR USE

May be taken with or without food.

SHELF LIFE

3 years from the date of manufacture.

REGISTRATION NUMBER
MAL19860211AZ

CONTROLLED MEDICINE
UBAT TERKAWAL

KEEP MEDICINES OUT OF REACH OF CHILDREN
JAUHI UBAT-UBATAN DARI KANAK-KANAK

For further information, please consult your doctor or your pharmacist.

Revision Number : 03
Revision Date : 19 Sep 2019

Lt-023.03 PRODUCT REGISTRATION HOLDER AND MANUFACTURER
IDAMAN PHARMA MANUFACTURING SDN BHD (661901-P)
LOT 24 & 25, JALAN PERUSAHAAN LAPAN, BAKAR ARANG INDUSTRIAL
ESTATE, 08000 SUNGAI PETANI, KEDAH DARUL AMAN, MALAYSIA.