

NEO-CORTISONE CREAM

VINEOxx-02 (MY)

DESCRIPTION

White opaque cream

COMPOSITION

Hydrocortisone 1.0% w/w
Neomycin (as sulphate) 0.5% w/w

ACTIONS AND PHARMACOLOGY

- **Pharmacodynamics:** Hydrocortisone, a less potent corticosteroid, diffuses across cell membranes, binds with DNA and stimulates transcription of messenger RNA (mRNA) resulting in protein synthesis of various enzymes. Neomycin, a bactericidal antibiotic, is actively transported across bacterial cell membrane, binds to bacterial ribosomes and interferes with initiation complex between mRNA and the ribosome resulting in inhibition of protein synthesis.
- **Pharmacokinetics:** Hydrocortisone is well absorbed through the skin after topical application. It rapidly distributed to all body tissues. It is metabolized mainly in the liver but also in other tissues, and are excreted in the urine.
Topical neomycin is readily absorbed through large denuded, burned, or granulating areas. The absorbed neomycin is rapidly excreted by the kidneys in active form.

INDICATIONS

For relief or treatment of infections accompanied by inflammation, in the following conditions:

- Minor skin irritation, itching and rashes due to
 - Eczema
 - Insect bites
 - Poison ivy, oak or sumac dermatitis
 - Detergents
 - Cosmetics and jewellery
- Anogenital pruritus

CONTRAINDICATIONS

- Avoid in patients known to be hypersensitive to aminoglycosides.
- Prolonged use is not recommended as it may lead to skin sensitisation.
- Avoid contact with eyes.
- Do not use on deep or puncture wounds, serious burns or raw areas unless directed by physician.
- Concurrent topical and systemic administration of neomycin or related drugs is not recommended since hypersensitive reactions may occur more frequently under these circumstances.

PRECAUTIONS

- Caution in pregnant patients and in patients with tuberculosis of the skin.
- Paediatric therapy continuing for longer than 2 weeks should be evaluated carefully by the physician.
- Caution in patients with extensive skin damage or perforated tympanic membranes as deafness may result.
- Do not bandage or otherwise wrap the treated skin area unless directed to do so by physician.
- Safety for use in lactation has not been established.

DRUG INTERACTION / INCOMPATIBILITIES

Concurrent use of Neomycin and the following drugs should be avoided:

- Aminoglycosides
- Vancomycin
- Cephalothin
- Polymyxins

PREGNANCY AND LACTATION

- **Pregnancy:** Problems in humans have not been documented; however, risk-benefit must be considered.
- **Lactation:** Safety for use in lactation has not been established.

MAIN SIDE / ADVERSE EFFECTS

Hypersensitive reactions; prolonged use include moon face, glaucoma, thinning of skin with easy bruising, increase in hair growth, loss of hair, reddish purple lines on arms, legs trunk or groin.

Note: It has fewer side effects on the skin than the more potent topical corticosteroids.

OVERDOSAGE

- **Clinical features:** Atrophic changes in the skin, burning, itching, irritation, dryness, folliculitis, hypertrichosis, acne and pigmentation.
- Treat overdosage by symptomatic measures, where possible the dosage being reduced or the drug slowly withdrawn.

DOSAGE AND ADMINISTRATION

Apply sparingly to the affected area.

Adults : Three or four times a day.
Children : One or two times a day.

Note: The information given here is limited. For further information consult your doctor or pharmacist.

Storage : Store below 30°C.
Protect from freezing.

Presentation/Packing : Cream 500 g, 50 g, 15 g.

Product Registration Holder / Manufactured by: HOVID Berhad
121, Jalan Tunku Abdul Rahman (Jalan Kuala Kangsar),
30010 Ipoh, Perak, Malaysia.

Date of Revision: June 2021