

NEO-BETASONE CREAM

VINEO06-var2 (MY)

DESCRIPTION

White opaque cream.

COMPOSITION

Betamethasone (as 17-valerate) 0.1% w/w
Neomycin Sulphate 0.5% w/w

ACTIONS AND PHARMACOLOGY

• Pharmacodynamics

- **Betamethasone - 17 - valerate:**

Betamethasone - 17- valerate is a corticosteroid. Its action is similar to that of cortisone. Betamethasone diffuse across cell membranes and complex with specific cytoplasmic receptors. These complexes then enter the cell nucleus, bind to DNA (chromatin) and stimulate transcription of messenger RNA (mRNA) and subsequent protein synthesis of various enzymes thought to be ultimately responsible for anti-inflammatory effects of topical application of betamethasone.

- **Neomycin:**

Neomycin, an aminoglycoside effective against most aerobic gram-negative bacilli, is actively transported across the bacterial cell membrane. It then binds to a specific receptor protein on the 30s subunit of bacterial ribosomes. This interferes with an initiation complex between mRNA (messenger RNA) and the 30s subunit, inhibiting protein synthesis. DNA may be misread, thus producing non-functional proteins. Polyribosomes are split apart and are unable to synthesize protein. Propagation of bacteria controlled.

• Pharmacokinetics

- **Absorption:**

After topical application, small amounts of the vale rate may be absorbed systematically during prolonged usage. Topical neomycin is readily absorbed from large denuded, burned or granulating areas.

- **Half-life:**

Biological half-life of neomycin is about 2 hours.

- **Distribution:**

Neomycin may enter the cerebrospinal fluid and may accumulate in renal function impairment.

- **Protein binding:**

About 60 to 65% of Betamethasone- 17-Valerate is bound to plasma proteins.

- **Excretion:**

Absorbed neomycin and betamethasone is excreted through the urine.

INDICATIONS

For treatment of infective acute or chronic inflammatory dermatoses such as:

- Localized neurodermatitis
- Psoriasis
- Atopic or seborrhoeic dermatitis
- Inflammatory phase of xerosis
- Anogenital pruritus
- Discoid lupus erythematosus
- Lichen planus
- Granuloma annulare
- Psoriatic plaques
- Allergic contact dermatitis or irritant dermatitis

CONTRAINDICATIONS

- Avoid in patients known to be hypersensitive to aminoglycosides.
- Prolonged use is not recommended as it may lead to skin sensitisation.
- Avoid contact with eyes.
- Do not use on deep or puncture wounds, serious burns, or raw areas unless directed by physician.
- Concurrent topical and systemic administration of neomycin or related drugs is not recommended since hypersensitivity reactions may occur more frequently under these circumstances.

PRECAUTIONS

- Caution in pregnant patients and in patients with tuberculosis of the skin.
- Paediatric therapy continuing for longer than 2 weeks should be evaluated carefully by the physician.
- Caution in patients with extensive skin damage or perforated tympanic membranes as deafness may result.
- Do not bandage or otherwise wrap the treated skin area unless directed to do so by physician.
- Safety for use in lactation has not been established.

DRUG INTERACTION / INCOMPATIBILITIES

- Avoid concurrent use of Neomycin and Aminoglycosides, other; Vancomycin, Cephalothin, Polymyxins.
- Betamethasone Valerate is inactivated by Coal Tar and Salicylic Acid.

PREGNANCY AND LACTATION

Safety for use in lactation has not been established.

MAIN SIDE/ADVERSE EFFECTS

Hypersensitivity reactions; prolonged use include moon face, glaucoma, thinning of skin with easy bruising, increase in hair growth, loss of hair, reddish purple lines on arms, legs, trunk or groin.

OVERDOSAGE

- **Clinical features:** Atrophic changes in the skin, burning, itching, irritation, dryness, folliculitis, hypertrichosis, acne and pigmentation.
- Treat overdosage by symptomatic measures, where possible the dosage being reduced or the drug slowly withdrawn.

DOSAGE AND ADMINISTRATION

Apply sparingly to the affected area.

Adults : One to three times a day.

Children : Once a day.

Note: The information given here is limited. For further information, consult your doctor or pharmacist.

Storage : Store below 30°C.
Protect from light.

Presentation/Packing : Cream 15 g

Product Registration Holder / Manufactured by: HOVID Bhd.
121, Jalan Tunku Abdul Rahman (Jalan Kuala Kangsar),
30010 Ipoh, Malaysia.

Revision date : October 2020