

CHLORPYRIMINE



(Chlorpyrimine Injection 10 mg/ml (10 ml))

Name and Strength of Active Substance(s)

Chlorpheniramine maleate 10 mg/ml (10 ml)

Product Description

Chlorpyrimine injection is a clear mobile liquid.

Pharmacodynamics

Antihistamine competitively antagonized histamine at H₁ receptor site but do not bind with histamine to inactivate it. They also have anticholinergic, antipruritic and sedative effects. Although symptoms of the common cold might be modified with antihistamine use, the drugs do not prevent or cure colds nor do they shorten the course of disease.

Pharmacokinetics

Following IV administration in humans, chlorpheniramine undergoes rapid and extensive IV administration reportedly average 2.5-3.2 L/kg in adults and 3.8 L/kg in children. Chlorpheniramine is distributed into saliva, and the drug and/or its metabolites appear to be distributed in small amounts into bile.

Following IV and administration maleate plasma concentrations of the drug have generally been reported to decline in biphasic manner; with a very rapid distribution phase. In adults with normal renal and hepatic function, the terminal elimination half-life of chlorpheniramine reportedly ranges from 12-43 hours; in children with normal renal and hepatic function, the terminal elimination half-life reportedly averages 9.6-13.1 hours (range: 5.2-23.1). In some patients with chronic renal failure undergoing hemodialysis the elimination half-life of chlorpheniramine reportedly ranged from 280-230 hours.

Chlorpheniramine and its metabolites are apparently excreted almost completely in urine. Urinary excretion of chlorpheniramine and its N-dealkylated metabolites varies with urinary pH and urine flow, following a single IV dose of chlorpheniramine maleate in healthy individuals with normal renal and hepatic function in one study about 20% of the dose was excreted in urine within 48 hours.

Indication

For the relief of some of the manifestation of allergic states such as hay fever, urticaria, allergic eczema, vasomotor rhinitis, serum sickness, atopic contact dermatitis.

Recommended Dosage

Dosage of chlorpheniramine should be individualized according to the patient's response and tolerance.

For the relief of allergic reactions to blood or plasma in adults and children 12 years of age and older a single chlorpheniramine maleate dose of 10-20 mg may be administered subcutaneously, IV, or IM. For the adjunctive therapy of anaphylaxis in adults and children 12 years of age and older a single dose of 10-20 mg may be administered IV. The maximum parenteral dosage of chlorpyramine maleate in adults and children may be given 0.0875 mg/kg or 2.5 mg/m² subcutaneously 4 times daily.

Route of Administration

Administered by intramuscular, subcutaneous or intravenous injection (drip slowly).

Contraindication

Hypersensitivity to antihistamines.

Warning and Precautions

CNS effect: CNS depression is common usual dosage of antihistamines. Sedation, ranging from mild drowsiness to deep sleep, occurs most frequently. Individuals who perform potentially hazardous tasks requiring mental alertness or physical coordination should be warned about possible drowsiness, dizziness.

Pediatric precautions: Like other antihistamine, chlorpheniramine should not be used in premature neonates. Safety and efficacy of chlorpheniramine maleate injection in children younger than 12 years of age have not been established.

Interactions with other Medicaments

Not applicable.

Statement on usage during pregnancy and lactation

Pregnancy

Not applicable.

Lactation

Not applicable.

Adverse Effects/ Undesirable Effects

Frequent drowsiness, dry mouth, dizziness and irritability occur with intermittent therapy. However, most patients develop tolerance to these side effects during continuous therapy, particularly if dosage is increased slowly.

Overdose and Treatment

Symptom: Effects may vary from mild CNS depression and cardiovascular collapse to stimulation, especially in children and geriatric patients.

Treatment: Treatment includes usual measures.

Storage Condition

Store below 30°C

Dosage forms and packaging available

10 mg/1 ml (10 ml) x 10 vials.

Name and address of manufacturer

ATLANTIC LABORATORIES CORPORATION LTD.

111 Moo 7, Bang Phli Noi, Bang Bo, Samut Prakan 10560, Thailand.

Name and address of product registration holder

ATLANTIC LABORATORIES (M) SDN. BHD.

S-G-11, Gaya Resort Homes, Persiaran Kemuning Damai, Seksyen 32, 40460 Shah Alam, Selangor, Malaysia.

Date of revision of PI

20/11/2025

Atlantic Laboratories Corporation Ltd.
Samut Prakan, Thailand.

201125 MAL(R)