

Pharmaniaga Frusemide Tablet

DESCRIPTION

A white to off white, round, biconvex tablet with scoreline on one side and a hexagon shaped marking on the other.

COMPOSITION

Each tablet contains Frusemide 40 mg.

ACTIONS

Frusemide increased diuresis generally even in cases in which this effect cannot be achieved in other ways. Increased urinary excretion usually begins within the first hour after ingestion of the tablets and lasts for an average of 6 to 8 hours. Duration and intensity of diuresis may vary in different patients. In the early phases of treatment a more pronounced effect is to be expected.

It is easily absorbed from the gastro-intestinal tract and have a rapid action of short duration, lasting for about 6 hours. It is strongly bound to plasma proteins and mainly excreted in the urine, almost entirely unchanged.

INDICATIONS

Oedema due to heart disease if satisfactory diuresis cannot be obtained by adequate treatment with cardiac drugs, ascites due to diseases of the liver and kidneys, complications of pregnancy such as oedema and eclampsia; oedema due to burns; stasis due to varicosis. Moreover, frusemide is suitable for treatment of hypertension of mild to moderate degree, alone or in combination with other drugs.

ADVERSE REACTIONS

General manifestations of intolerance (gastrointestinal symptoms) and allergic reactions such as skin rashes occur rarely under frusemide treatment. Side-effects resulting from the increased urinary excretion are more likely to occur, especially after prolonged high-dosage treatment.

If the patient salt intake is too low, electrolyte depletion may be reflected in leg cramps, anorexia, weakness, dizziness, drowsiness, and other disturbances.

CONTRAINDICATIONS AND PRECAUTIONS

Frusemide should not be used in cases of renal failure with reduced or absent urinary excretion (oliguria, anuria) resulting from poisoning with nephrotoxic (mercury bichloride) or hepatotoxic agents (carbon tetrachloride, poisonous amanita). This also applies to severely impaired renal function with simultaneous liver failure.

Severe hypokalaemia must be compensated before frusemide treatment. Potassium loss maybe due to unbalanced diet, to vomiting or diarrhoea (frequent use of laxatives), or to disease of the liver or adrenal cortex, and, in diabetics, to hyperglycaemia.

In diabetics, a change of diet or medication may become necessary during frusemide treatment.

During pregnancy, especially at first three months, frusemide should only be used if absolutely necessary (e.g. in the presence of eclampsia).

It should be noted that frusemide passes into breast milk and, like all diuretics, may inhibit lactation.

Special Notes

Patients with manifest or latent diabetes mellitus should be checked regularly during frusemide treatment. The same applies to patients with gout. Therefore, patients who suffer from diabetes or gout or have a case of diabetes or gout in their family, should inform their physician about this before frusemide treatment is started. Also the existence of urinary tract obstruction (e.g. in case of prostatic hypertrophy) should be mentioned.

DOSAGE AND ADMINISTRATION

Frusemide tablets are swallowed whole and in the dosage prescribed by the physician. Generally, treatment is initiated in adults with a daily dose of ½ - 1 tablet, taken preferably in the morning. Further treatment depends on the extent of the diuresis. Special dosage guidelines apply to children, each case being considered individually. For this reason, only the physician can decide on the dose suitable for each case. Generally, daily doses of 1 - 3 mg/kg body weight, up to maximum of 40 mg/day are recommended for infants and children. More exact directions are given in the following table.

Frusemide dosage for children:		
Age of child	Body weight in kg	Daily dosage (1 tablet = 40 mg)
1 month	3.5	7.0 mg
2 months	4.0	8.0 mg
3 months	4.8	10.0 mg
4 months	5.5	11.0 mg
5 months	5.8	12.0 mg
6 months	6.5	13.0 mg
1 year	9.0 - 9.5	18.0 mg
2 years	10.0 - 10.5	20.0 mg
3 years	12.5 - 13.0	25.0 mg
4 years	14.5 - 15.0	29.0 mg
5 years	15.5 - 16.5	30.0 mg
6 years	16.5 - 17.0	33.0 mg
10 years	27.0 - 29.0	40.0 mg
12 years	31.0 - 32.0	40.0 mg

Rigorous salt restriction is not necessary during frusemide treatment. In long-term therapy, a high potassium diet is recommended (lean meat, potatoes, tomatoes, spinach, cauliflower and dried fruit).

OVERDOSAGE

Overdosage will lead to fluid and salt depletion subsequent to initial excessive diuresis. Medical help must be called in without delay so as to ensure immediate measures for an adequate restoration of water and electrolyte balances.

ROUTE OF ADMINISTRATION

Oral.

STORAGE CONDITIONS

Store below 30°C.
Keep container tightly closed.
Protect from light.

SHELF LIFE

Product should not be used beyond the expiry date imprinted on the product packaging.

PRESENTATION

In blisters of 500 tablets.

Date of revision: 11th July 2024

PRODUCT REGISTRATION HOLDER/

MANUFACTURER:

PHARMANIAGA MANUFACTURING BERHAD
(198001006232)

No. 11A, Jalan P/1, Kawasan Perusahaan Bangi,
43650 Bandar Baru Bangi, Selangor Darul Ehsan,
Malaysia.

PRP 0034.11 110724