



H-CORT CREAM & OINTMENT 1% W/W

NAME AND STRENGTH OF ACTIVE SUBSTANCE: Hydrocortisone 1% w/w.

PRODUCT DESCRIPTION:

Cream: A white cream.

Other ingredients: White soft paraffin, Liquid paraffin, Chlorocresol, Cetostearyl alcohol, Propylene glycol, Macrogol cetostearyl ether (Cetomacrogol 1000) and Purified water.

Ointment: A white ointment.

Other ingredients: White soft paraffin and Liquid paraffin.

PHARMACODYNAMICS : Hydrocortisone is a topically active corticosteroid which has anti-inflammatory, antipruritic and vasoconstrictive actions.

PHARMACOKINETICS : Corticosteroids are extensively bound to plasma protein.

Only unbound corticosteroids has pharmacological effects or is metabolised. They are metabolised mainly in the liver and also in the kidney, and are excreted in the urine.

INDICATIONS : Suppression of skin diseases in which inflammation is a prominent feature, e.g. eczema, infantile eczema, atopic dermatitis, dermatitis herpetiformis, contact dermatitis, neurodermatitis, psoriasis and intertrigo.

CONTRAINDICATION : Hypersensitivity to hydrocortisone, other corticosteroids, or any component in the base.

DRUG INTERACTION : Not known.

ADVERSE EFFECT : Reported local adverse reactions are burning sensation, itching, irritation, dryness, folliculitis, hypertrichosis, acneiform eruptions, hypopigmentation, maceration of the skin, secondary infection, skin atrophy, striae and milaria. Systemic adverse reactions, such as vision blurred, have also been reported with the use of topical corticosteroids.

PRECAUTIONS/WARNINGS: Discontinue use if irritation, sensitivity or other reaction develop and institute appropriate treatment. Corticosteroid applied to the skin can be absorbed in sufficient amount to produce systemic effects such as hypothalamic-pituitary-adrenal axis suppression, manifestation of Cushing's syndrome, hyperglycaemia and glucosuria. Thus suitable precautions should be taken during prolonged treatment, when treating extensive body surface areas, when using the occlusive technique and when treating children (because of a large skin surface area to body weight ratio).

Not suitable for ophthalmic use.

Visual disturbance may be reported with systemic and topical corticosteroids use. If a patient presents with symptoms such as blurred vision or other visual disturbances, the patient should be considered for referral to an ophthalmologist for evaluation of possible causes which may include cataract, glaucoma, or rare diseases such as central serous chorioretinopathy (CSCR) which may have been reported after use of systemic and topical corticosteroids.

PREGNANCY AND LACTATION : Safety of use of topical corticosteroids during pregnancy and lactation has not been established. Thus it should be used during pregnancy only if the potential benefit justifies the potential risk to the foetus. It should not be used extensively on pregnant patients in large amounts or for prolonged periods of time. Caution should be exercised when administering it to nursing mothers.

RECOMMENDED DOSAGE : Apply thinly to the affected area one to four times daily, depending on the seriousness of the condition. Occlusive dressings may be used for the management of psoriasis or recalcitrant conditions.

ROUTE OF ADMINISTRATION : Topical.

OVERDOSE AND TREATMENT : If hypothalamic-pituitary-adrenal axis suppression is found, then the drug should be withdrawn, frequency of application reduced or weaker steroid used. Supplemental systemic corticosteroids may be required if signs and symptoms of steroid withdrawal occur.

PACKING : Collapsible aluminium tube of 15g. Plastic jars of 450g (For Export Only). Not all pack size may be available locally.

STORAGE : Keep containers well closed. Protect from strong light. Store below 30°C.

For external use only. Keep out of reach of children.

Recommended shelf-life: 3 years.

MANUFACTURER : HOE Pharmaceuticals Sdn. Bhd.
Lot 10, Jalan Sultan Mohamed 6, Bandar Sultan Suleiman,
42000 Port Klang, Selangor, MALAYSIA.

Revision date : March 2021