

PACKAGE INSERT

1 cm
Barcode

hovid

DEXCOPHAN

VIDEX11-var (MY)

DESCRIPTION

Dexcophan Tablet

Round, white film-coated tablet, bevel-edged, flat-faces; "HD" and break-bar embossed on the same face.

Dexcophan Paediatric Linctus

Red coloured liquid with sweet blackcurrant taste.

COMPOSITION

Dexcophan Tablet

Each film-coated tablet contains: Dextromethorphan HBr 15 mg

Dexcophan Paediatric Linctus

Each 5 ml contains: Dextromethorphan HBr 5.5 mg

PHARMACODYNAMICS

Dextromethorphan Hydrobromide has a depressant action on the cough center similar to that produced by codeine. However unlike codeine, it has no analgesic or expectorant effect and no other depressant action on the central nervous system.

Dextromethorphan acts centrally to elevate the threshold for coughing and is therefore used as an antitussive.

PHARMACOKINETICS

Absorption: Dextromethorphan is fully absorbed from the gastrointestinal tract.

Blood Concentration: After an oral dose of 30mg of dextromethorphan HBr, 10, 25 and 10% of the dose was detected in the blood at 1, 2.5 and 6 hours respectively after administration. Nevertheless plasma levels of unmetabolised dextromethorphan following the administration of therapeutic oral doses are very low due to extensive metabolism in the liver (first pass effect) and distribution which remove the drug rapidly from the blood.

Metabolism: After oral administration, dextromethorphan is absorbed from the intestinal tract and passes via the portal vein to the liver before entering the general circulation. In the liver it is metabolised (first pass effect) to an extent which is large but which varies widely between individuals. Three metabolites have been identified in man. All three metabolites have approximately the same antitussive potency as dextromethorphan in guinea pigs. There is a wide variation between individuals in the large first pass effect. Therefore, the proportions of dextromethorphan that are O-demethylated and N-demethylated vary widely between individuals. After O- or N-demethylation, the metabolites undergo glucuronic acid or sulphate conjugation.

Excretion: Up to 56% of a dose is excreted in the urine, about 8% being excreted unchanged in 6 hours. However there is a wide variation in the reported excretion of dextromethorphan and its metabolites. This reflects the large intersubject differences in the first pass effect.

INDICATIONS

Dextromethorphan is a cough suppressant used in the treatment of unproductive cough.

CONTRAINDICATIONS

Contraindicated in patients hypersensitive to propylene glycol.

WARNINGS AND PRECAUTIONS

- Caution in patients with asthma and hepatic function impairment.
- Safety for use in pregnancy and lactation has not been established.

- Not to be used in children less than 2 years of age.
- To be used with caution and doctor's / pharmacist's advice in children 2 to 6 years of age.

PREGNANCY AND LACTATION

Safety for use in pregnancy and lactation has not been established.

DRUG INTERACTIONS

Concurrent use with monoamine oxidase (MAO) inhibitors is not recommended since it may result in CNS depression.

MAIN SIDE/ADVERSE EFFECTS

Drowsiness, dizziness, gastrointestinal disturbances, mental confusion.

OVERDOSE

Clinical features: Nausea, vomiting, abdominal discomfort, hallucinations, hypotension, ataxia, respiratory depression.

Treat overdose by emesis or gastric lavage, if appropriate. Administer naloxone 0.4 - 1.2 mg IV or IM in an adult or 5 - 10 µg per kg body weight IV or IM in children, repeated as necessary.

DOSAGE AND ADMINISTRATION

Oral.

Adults

Tablet

15 to 30 mg every four hours as needed; up to 120 mg per day

Children

Tablet

7 to 12 years of age: 7.5 mg to 15 mg every four hours as needed; up to 60 mg per day.

Linctus

- 2 to 6 years of age: 5 ml every four hours as needed; up to 27 ml per day.
- 7 to 12 years of age: 10 ml every four hours as needed; up to 54 ml per day.

Note: The information given here is limited. For further information, consult your doctor or pharmacist.

Dexcophan Tablet

Storage: Store below 25°C. Protect from light.

Presentation/Packing:

Film-coated tablet 15 mg x 1000's,
Blisters of 10 x 10's, 100 x 10's

Product Registration Holder: HOVID Bhd.

121, Jalan Tunku Abdul Rahman, 30010 Ipoh, Malaysia.

Manufactured by: HOVID Bhd.

Lot 56442, 7 1/2 Miles, Jalan Ipoh / Chemor,
31200 Chemor, Malaysia.

Dexcophan Paediatric Linctus

Storage: Store below 30°C. Protect from light and freezing.

Presentation/Packing:

Paediatric Linctus 5.5 mg/5 ml x 60 ml, 100 ml.

Product Registration Holder /

Manufactured by: HOVID Bhd.

121, Jalan Tunku Abdul Rahman
(Jalan Kuala Kangsar),
30010 Ipoh, Perak, Malaysia.

Revision date: March 2021

hovid