

POCIN H EAR DROPS

DESCRIPTION

A white to off-white, homogenous suspension.
Each ml contains: Polymyxin B Sulphate 10,000 iu, Neomycin Sulphate 3400 iu and Hydrocortisone 10mg

INDICATIONS

Pocin H ear drops are indicated in the treatment of external ear canal infections and mastoidectomy cavity infections caused by susceptible organisms.

PHARMACODYNAMICS

Neomycin Sulphate

Neomycin is an aminoglycoside which is actively transported across the bacterial cell membrane where it binds to a specific receptor protein on the 30 S subunit of bacterial ribosomes. At this site, it interferes with an initiation complex between mRNA and the 30 S subunit, inhibiting protein synthesis. DNA may be misread, thus producing nonfunctional proteins, polyribosomes are split apart and are unable to synthesize protein. Neomycin is bactericidal.

Polymyxin B Sulphate

Polymyxin B is bactericidal and active against *Pseudomonas aeruginosa* and other gram-negative bacteria. It is a surface-active basic polypeptide that binds to anionic phospholipid sites in bacterial cytoplasmic membranes, disrupts membrane permeability to allow leakage of intracellular contents. Its action is antagonised by calcium and magnesium.

Hydrocortisone

Hydrocortisone diffuses across cell membranes and complex with specific cytoplasmic receptors. These complexes enter the cell nucleus, bind to DNA and stimulate transcription of mRNA and subsequent protein synthesis of enzymes responsible for the anti-inflammatory effects of hydrocortisone. In high concentrations, hydrocortisone may exert direct membrane effects. Hydrocortisone decreases cellular and fibrinous exudation and tissue infiltration, inhibits fibroblastic and collagen forming activity, retards epithelial, regeneration, diminishes postinflammatory neovascularisation, and reduces toward normal levels the excessive permeability of inflamed capillaries.

PHARMACOKINETICS

Neomycin, polymyxin B and hydrocortisone may be absorbed following topical application to the ear if the eardrum is perforated or tissue damage is present.

DOSAGE

Adult

External ear canal infection

4 drops to the ear canal 3 or 4 times a day.

Mastoidectomy cavity infections

4 to 10 drops to the mastoidectomy cavity or ear canal every 6 to 8 hours.

Paediatric

External ear canal infection

3 drops to the ear canal 3 or 4 times a day.

Mastoidectomy cavity infections

4 or 5 drops to the mastoidectomy cavity every 6 to 8 hours.

Route of Administration: Ear

SIDE EFFECTS

Hypersensitivity reactions have been reported following application of Pocin H ear drops.

CONTRAINDICATIONS

Pocin H ear drops should not be used in patients with a sensitivity to aminoglycoside, polymyxins or sulphates. The use of Pocin H ear drops are also contraindicated in bullous myringitis, herpes simplex, herpes zoster oticus, tubercular or fungal infections of the ear, vaccinia, varicella, or other viral disease of the ear, chronic otitis media or perforated eardrum.

PRECAUTIONS

Patients sensitive to one aminoglycoside or polymyxin may be sensitive to other aminoglycosides or polymyxins also.

Risk of sensitization in relation to thiomersal and other preservatives.

INTERACTIONS WITH OTHER MEDICAMENTS

Co-treatment with CYP3A inhibitors, including cobicistat-containing products, is expected to increase the risk of systemic side-effects. The combination should be avoided unless the benefit outweighs the increased risk of systemic corticosteroid side-effects, in which case patients should be monitored for systemic corticosteroid side-effects.

OVERDOSAGE

There is no relevant information available for neomycin, polymyxin B and hydrocortisone.

TOXICOLOGY

Pregnancy/Reproduction

Topical corticosteroids have been shown to be teratogenic in rabbits when applied at concentration of 0.5% on days 6 to 18 of gestation and in mice when applied at a concentration of 15% on days 10 to 13 of gestation.

(FDA Pregnancy Category C)

No breast-feeding, paediatric and geriatric specific problems have been reported to date.

PRESENTATION

5ml x 12 bottles

STORAGE CONDITIONS AND USER INSTRUCTIONS

Store in a dry place below 25°C.

For external use only.

Discard contents 4 weeks after opening.

Keep out of reach of children.

Jauhi daripada kanak-kanak.

Shelf life : Please refer to outer package.

Product Registration Holder:

Duopharma Manufacturing (Bangi) Sdn. Bhd.

Lot No. 2, 4, 6, 8, & 10, Jalan P/7, Section 13, Bangi Industrial Estate, 43650 Bandar Baru Bangi, Selangor, Malaysia.

Manufacturer:

Duopharma Manufacturing (Bangi) Sdn. Bhd.

Lot No. 2 & 4, Jalan P/7, Section 13, Bangi Industrial Estate, 43650 Bandar Baru Bangi, Selangor, Malaysia.



1500008664 N.1

Revision Date: December 2023

POCIN H EAR DROPS

170mm(w) x 185mm (h)

Black 100%