

PAC2505-0002
Label - Xepagan Elixir 60ml (MY)
Material code: KLF0052L1
Size: 195(L) x 170(H)mm
Color: Black
Revision updated: 005



Xepa-Soul Pattinson (Malaysia) Sdn. Bhd. 1-5 Cheng Industrial Estate 75250 Melaka, Malaysia

Important information. Please read carefully.

XEPAGAN / XEPAGAN-5

Elixir

Composition

Xepagan Elixir : Each 5ml contains Promethazine Hydrochloride 10mg. Contains Sodium Benzoate 0.1% w/v as preservative. Alcohol content: 0.19 % v/v.
Xepagan-5 Elixir : Each 5ml contains Promethazine Hydrochloride 5mg. Contains Sodium Benzoate 0.1% w/v as preservative. Alcohol content: 0.86 % v/v.

Pharmacodynamics

Xepagan contains a long acting antihistamine which is used for its anti-allergic, anti-emetic, sedative and hypnotic actions.

Pharmacokinetics

Promethazine is well absorbed after oral doses and peak plasma concentration occur 2 to 3 hours after dose. Metabolised in the liver, then excreted mainly as metabolites in the urine. Promethazine undergoes extensive metabolism, mainly to promethazine sulfoxide, and also to N-desmethylpromethazine. It is excreted slowly via the urine and bile.

Indications

Xepagan is indicated for symptomatic relief of allergy such as hay fever, urticaria, treatment of nausea and vomiting and motion sickness; and as a sedative for mild insomnia.

Dosage and Administration

For oral administration only.

		Recommended Dose	
		Xepagan	Xepagan-5
Allergic conditions	Adults	5ml twice daily. This may be increased to 10ml three times daily if necessary.	10ml twice daily. This may be increased to 20ml three times daily if necessary.
	Children 6-12 years	Single dose at night 5ml-12.5ml or 2.5ml-5ml two times daily (max 12.5ml daily)	Single dose at night 10ml-25ml or 5ml-10ml two times daily (max 25ml daily)
	Children 2-5 years	Single dose at night 2.5ml-7.5ml or 2.5ml two times daily (max 7.5ml daily)	Single dose at night 5ml-15ml or 5ml two times daily (max 15ml daily)
Nausea and Vomiting	Adults	12.5ml every 4 to 6 hours (max 50ml daily)	25ml every 4 to 6 hours (max 100ml daily)
	Children 6-12 years	5ml every 4 to 6 hours (max 12.5ml daily)	10ml every 4 to 6 hours (max 25ml daily)
	Children 2-5 years	2.5ml every 4 to 6 hours (max 7.5ml daily)	5ml every 4 to 6 hours (max 15ml daily)
Travel Sickness (the night before travel and repeated after 6-8 hours if required)	Adults	Single dose 12.5ml	Single dose 25ml
	Children 6-12 years	Single dose 5ml	Single dose 10ml
	Children 2-5 years	Single dose 2.5ml	Single dose 5ml

Contraindications

Xepagan should not be used in patients with jaundice induced by other phenothiazine derivatives or with known hypersensitivity to phenothiazines as cross-sensitivity may occur.

Warnings

Xepagan is strictly not to be used for infants six months of age and younger.It is also not recommended for children below 2 years old because younger children are more prone to serious side-effects of promethazine.This product contains promethazine hydrochloride.It should not be used in pediatric patients less than 2 years of age because of the potential for fatal respiratory depression. **Xepagan** is also to be used with caution in children 2 years and older. This preparation contains Sodium metabisulphite that may cause serious allergic type reactions in certain susceptible patients. Do not use if known to be hypersensitive to bisulphites.

Precautions

Xepagan should be used with caution in patients with cardiovascular disease and liver disease. As the preparations may cause drowsiness, adult patients should be advised against driving or operating machinery and to avoid consuming alcoholic drinks while on this medication.

Promethazine may cause the skin to be more sensitive to the sun especially if the patients have eczema or a tendency to rheumatism.The skin should be protected from exposure to bright sunlight.

To be used with cautions, and as advised by the physician/pharmacist for children age 2 to 6 years.

Pregnancy and Lactation

As with all medicines, use in pregnancy and lactation is best avoided unless necessary, where close medical supervision is required.

Drug Interactions

CNS Depressants- Xepagan may increase, prolong, or intensify the sedative action of other central-nervous-system depressants, such as alcohol, sedatives/hypnotics (including barbiturates), narcotics, narcotic analgesics, general anesthetics, tricydic antidepressants, and tranquilizers. Such agents should be avoided or administered in reduced dosage to patients receiving promethazine HCl.

Epinephrine- Xepagan is potential to reverse epinephrine's vasopressor effect, Epinephrine should NOT be used to treat hypotension associated with **Xepagan**.

Anticholinergics- Concomitant use of other agents with anticholinergic properties should be undertaken with caution.

Monoamine Oxidase Inhibitors (MAOI)- Concomitant use of MAOI and phenothiazines have been reported to increase incidence of extrapyramidal effects.

Side Effects

For the doses recommended, side effects are usually mild and transient. The more common side effects include drowsiness, tinnitus, lassitude, fatigue, dryness of mouth, dysuria, loss of appetite, nausea or vomiting, diarrhoea or constipation, restlessness, dizziness, blurred vision and gastrointestinal disturbances. Other side effects which have been reported include occasional hypotension, leucopenia. Less common but more serious side effects include fever, breathing difficulty, irregular heart beat, jaundice, tremor or convulsions.

Overdosage

In cases of overdosage and poisoning, hospital admission is strongly advised. Symptoms include convulsions and/or tremors, hallucinations, ataxia, incoordination, difficulty in breathing, drowsiness and coma. Fixed dilated pupils with a flushed face and fever are common signs of overdosage. As there is no specific antidote available, aspiration and gastric lavage may be necessary followed by symptomatic and supportive treatment.

Shelf-life : The expiry date is indicated on the packaging.

Presentation

Yellowish brown colour elixir with orange flavour in packs of 60ml.

Storage : Store below 30°C. Protect from light.

KEEP OUT OF REACH OF CHILDREN
JAUIH DARI KANAK-KANAK

For further information, please consult your pharmacist or physician.

Revision Date : 15-May-2025

Manufacturer and Product Registration Holder
Xepa-Soul Pattinson (Malaysia) Sdn. Bhd.
1-5, Cheng Industrial Estate,
75250 Melaka, Malaysia.

KLF0052L1

005



Xepa-Soul Pattinson (Malaysia) Sdn. Bhd. 1-5 Cheng Industrial Estate 75250 Melaka, Malaysia

PATIENT INFORMATION LEAFLET

XEPAGAN / XEPAGAN-5

Elixir

Please read this leaflet carefully before you start to take your medicine. This leaflet is a summary of important information about your medicine. If you have any questions or are not sure about anything, ask your doctor or pharmacist.

What is in Xepagan?

The active ingredient of Xepagan is Promethazine Hydrochloride.

Xepagan Elixir contains Promethazine Hydrochloride 10mg in each 5ml of syrup. Also contains Sodium Benzoate 0.1% w/v as preservative. Alcohol content is 0.19% v/v.

Xepagan-5 Elixir contains Promethazine Hydrochloride 5mg in each 5ml of syrup. Also contains Sodium Benzoate 0.1% w/v as preservative. Alcohol content is 0.86% v/v.

Xepagan preparations are yellowish brown colour elixir with orange flavour.This syrup liquid is available in pack sizes of 60ml.

Xepagan belongs to a group of medicines called phenothiazines. It is an antihistamine that has anti-allergic, anti-emetic and sedative actions.

What is Xepagan used for?

Xepagan is used for relief of allergy such as hay fever, rashes, hives, treatment of nausea and vomiting, prevention of motion sickness, and as a short term sedative.

How much and how often should you use Xepagan?

		Recommended Dose	
		Xepagan Elixir	Xepagan-5 Elixir
Allergic conditions	Adults	5ml twice daily. This may be increased to 10ml three times daily if necessary.	10ml twice daily. This may be increased to 20ml three times daily if necessary.
	Children 6-12 years	Single dose at night 5ml-12.5ml or 2.5ml-5ml two times daily (max 12.5ml daily)	Single dose at night 10ml-25ml or 5ml-10ml two times daily (max 25ml daily)
	Children 2-5 years	Single dose at night 2.5ml-7.5ml or 2.5ml two times daily (max 7.5ml daily)	Single dose at night 5ml-15ml or 5ml two times daily (max 15ml daily)
Nausea and Vomiting	Adults	12.5ml every 4 to 6 hours (max 50ml daily)	25ml every 4 to 6 hours (max 100ml daily)
	Children 6-12 years	5ml every 4 to 6 hours (max 12.5ml daily)	10ml every 4 to 6 hours (max 25ml daily)
	Children 2-5 years	2.5ml every 4 to 6 hours (max 7.5ml daily)	5ml every 4 to 6 hours (max 15ml daily)
Travel Sickness (the night before travel and repeated after 6-8 hours if required)	Adults	Single dose 12.5ml	Single dose 25ml
	Children 6-12 years	Single dose 5ml	Single dose 10ml
	Children 2-5 years	Single dose 2.5ml	Single dose 5ml

When should you not take this medicine?

Do not take this medicine if you are allergic to promethazine or any of the ingredients in the product. Xepagan should be used with caution in patients with medical conditions such as cardiovascular disease and liver disease.If you are taking other medicines or suffering from other medical conditions, consult a doctor or pharmacist before taking this medicine.

As with all medicines, use in pregnancy is best avoided unless necessary, where dose medical supervision is required. Do not take this medicine during pregnancy or breast feeding, before consulting your doctor or pharmacist.

Special Warnings

Xepagan is strictly not to be used for infants six months of age and younger.

It is also not recommended for children below 2 years old because younger children are more prone to serious side-effects of promethazine.

Xepagan is also to be used with caution in children 2 years and older.

As the preparations may cause drowsiness, adult patients should be advised against driving or operating machinery and to avoid consuming alcoholic drinks while on this medication.

Promethazine may cause your skin to be more sensitive to the sun especially if you have eczema or a tendency to rheumatism.You should protect your skin from exposure to bright sunlight.

To be used with cautions, and as advised by the physician/pharmacist for children age 2 to 6 years.

Does Xepagan have side effects?

A medicine may cause side effects in certain individuals.If you experience side effects and are concerned, you should seek advice from your doctor or pharmacist.

For the doses recommended, side effects are usually mild and transient.

The more common side effects include drowsiness, tinnitus, lassitude, fatigue, dryness of mouth, dysuria, loss of appetite, nausea or vomiting, diarrhoea or constipation, restlessness, dizziness, blurred vision and gastrointestinal disturbances.

Other side effects which have been reported include occasional hypotension, leucopenia. Less common but more serious side effects include fever, breathing difficulty, irregular heart beat, jaundice, tremor or convulsions.

What other medicine or food should be avoided whilst taking Xepagan?

Avoid using alcohol, sleeping pills, medicines for reducing anxiety and tension or any other medicines that suppress breathing.

What should you do if you miss a dose?

If you are taking this medicine regularly and miss a dose, take it as soon as possible. However, if it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take a double dose to make up for the dose that you have missed.

How should you keep Xepagan?

Always ensure the container is well closed, kept below 30°C and protected from light.

Keep Xepagan in a safe place and out of reach of children.

What are the symptoms of overdose and what to do when you have taken more than the recommended dosage?

Symptoms of overdose include over-excitement, shaky and unsteady movements, convulsions and/or tremors, hallucinations, difficulty in breathing, drowsiness, unconscious, fixed dilated pupils with a flushed face and fever.

If you experience any of the symptoms, or if you have taken more than the recommended dose, seek immediate medical attention.

When should you consult your doctor?

You need to consult a doctor when:

- You experience common side effects that bother you and serious side effects that need urgent medical attention.
- You have taken this medicine for 7-10 days in a row and your condition is not improving.
- You have taken an overdose of the medicine.

KLF0052L1

005

