
PIRI EXPECTORANT

Each 5ml contains:

Chlorpheniramine Maleate BP.....	2.5 mg
Ammonium Chloride BP.....	125mg
Trisodium Citrate BP.....	55mg
Ethanol.....	0.48% v/v
Preservatives:	
Methyl Paraben.....	0.1% w/v
Propyl Paraben.....	0.01% w/v

Description:

Product Description

Red liquid with raspberry flavor

Pharmacodynamics

It is widely distributed throughout the body tissues and appears to be subject to enterohepatic circulation. About 70% is bound to plasma proteins. It is extensively metabolized to form polar and non-polar metabolites. About 35% a dose is excreted as unchanged drug but it is increased by an increase in urinary pH and urinary flow, less than 1% of a dose is excreted in the faeces.

Pharmacokinetics

It is readily absorbed after oral administration. Peak concentrations appear within 2 to 3 hours. Plasma half-life has been reported as 12 to 15 hours.

Indications

For symptomatic treatment of perennial and seasonal allergic rhinitis, vasomotor rhinitis, and allergic conjunctivitis due to inhalant allergens and foods. Symptomatic treatment of mild, uncomplicated allergic skin manifestation of urticaria and angioedema.

Recommended Dose

Adults: 5 - 10ml, 3 - 4 times daily.

Children: 2 - 5 years: 2.5 – 5 ml 3 times daily.

Children 6 - 12 years: 5 – 10 ml, 3 – 4 times daily.

Route of Administration

Oral.

Contraindications

Chlorpheniramine is contraindicated in patients who are hypersensitive to antihistamines or any of the excipients listed.

Chlorpheniramine is contraindicated in patients who had treatment with Monoamine Oxidase Inhibitors (MAOI's) within the last 14 days as the anticholinergic properties of chlorpheniramine are intensified by MAOI's.

Warning and Precautions

When used for treatment of cough and cold;

- (a) Not to be used in children less than 2 years of age.
- (b) To be used with caution and doctors/ pharmacist's advice in children 2 to 6 years of age.

Chlorpheniramine has an anticholinergic effects and should be used with caution in patients with epilepsy, raised intra-ocular pressure including glaucoma, prostatic hypertrophy, severe hypertension, cardiovascular disease, bronchitis, bronchiectasis, asthma, hepatic impairment. Children and the elderly patients are more likely to experience the neurological anticholinergic effects and paradoxical excitation (e.g. increased energy, restlessness, nervousness).

The effects of alcohol may be increased and therefore should be avoided. Chlorpheniramine should not be used with other antihistamine containing products such as antihistamine containing cough and cold medicines.

Patients with rare hereditary problems of fructose intolerance should not take this medicine.

Interactions with Other Medicaments

Alcoholic drinks and certain other central nervous system depressants such as anxiolytics or hypnotics can potentiate the sedative effects of chlorpheniramine.

Phenytoin metabolism is inhibited by chlorpheniramine and this can cause phenytoin toxicity.

The anticholinergic effects of chlorpheniramine are intensified by the use of other anticholinergic drugs such as atropine, tricyclic antidepressants and MAOI's.

Pregnancy & Lactation

Pregnancy

There are no adequate data for the use of chlorpheniramine in pregnant woman, the potential risk in humans is unknown. Use during the third trimester may result in reactions in the newborn or premature neonates, therefore it should not to be used during pregnancy unless considered essential by a physician.

Lactation

Chlorpheniramine maleate and other antihistamines may inhibit lactation and may be secreted into the breast milk. Should not be used during lactation unless considered essential by a physician.

Side Effects

The most common effect is sedation, varying from slight drowsiness to deep sleep and including lassitude, dizziness and incoordination, disturbances in concentrating, dizziness and headache. Sedative effects may diminish after a few days.

Other side effects include gastro-intestinal disturbances such as nausea, vomiting, diarrhoea or constipation, anorexia or increased appetite, insomnia, irritability, nightmare, hallucinations, euphoria and epigastric pain. It also produce anti-muscarinic effects including blurred vision, difficulty in micturition, dysuria, dryness of the mouth and tightness of the chest.

Symptoms and Treatment of Overdose

Symptoms of overdose include: anticholinergic effects (clumsiness or unsteadiness; severe dryness of mouth, nose or throat; flushing or redness of face; shortness of breath or troubled breathing); CNS depression (severe drowsiness); CNS stimulation (hallucination, seizures, trouble in sleeping); severe hypotension (feeling faint); cardiac arrhythmias (fast or irregular heart beat).

If vomiting has not occurred spontaneously, emesis should be induced. Syrup of ipecac is preferred. Precautions against aspiration must be taken especially in children and infants. If vomiting is unsuccessful, gastric lavage is indicated within 3 hours after ingestion and even later if large amounts of milk or cream were given. Isotonic and 0.45% Sodium Chloride Solution is the lavage solution of choice.

In unconscious patient, the airway should be secured with a cuffed endotracheal tube before attempting to evacuate the gastric contents.

Hypotension is an early sign of impending cardiovascular collapse and should be treated vigorously. Treatment with IV infusion of a vasopressor (eg: Dopamine titrated to maintain blood pressure is also necessary. Do not treat CNS depression with analeptics which might precipitate convulsions. Convulsions should be controlled by careful use of a short-acting barbiturates.

Effects and Ability to Drive and Use Machine

As with all antihistamines, dizziness, drowsiness, blurred vision and psychomotor impairment may occur. Extreme caution should be advised when driving or operating machinery.

Storage Condition

Keep container tightly closed. Protect from light. Store below 30°C.

Presentation:

120ml bottle.

Registration Number

MAL19871712AZ

Manufactured by & Product Registration Holder

KCK Pharmaceutical Industries Sdn. Bhd.

Plot 61, Bayan Lepas Industrial Park Phase 4,
Lintang Bayan Lepas 1, 11900 Bayan Lepas,
Pulau Pinang, Malaysia

Date of Revision:

28 April 2023