

Important Information. Please read carefully.



Zaricort[®]

Cream

COMPOSITION

Contains Miconazole Nitrate 2% w/w and Hydrocortisone 1% w/w in a non greasy, non-staining cream basis. Benzoic Acid 0.15% w/w as preservative.

PHARMACODYNAMICS

Miconazole is an imidazole antifungal agent that interferes with ergosterol synthesis and therefore alters the permeability of the cell membrane of sensitive fungi. Miconazole is active against the common dermatophytes *Trichophyton rubrum*, *T. mentagrophytes* and *Epidermophyton*; the yeast *Candida albicans*; and the organism responsible for tinea versicolor, *Malassezia furfur*.

Hydrocortisone is a mild corticosteroid with glucocorticoid activity.

PHARMACOKINETICS

There is little absorption through skin or mucous membranes when miconazole nitrate is applied topically. Miconazole is metabolized in the liver to inactive metabolites. The half-life of miconazole is about 24 hours.

Hydrocortisone is absorbed through the skin, particularly in denuded areas. Hydrocortisone is metabolized in the liver and most body tissues to degraded forms and are excreted in the urine. Hydrocortisone readily crosses the placenta.

INDICATIONS

Zaricort Cream is used for the topical treatment of fungal infections where inflammation of the skin is also present such as those of tinea pedis (athlete's foot), tinea versicolor (white spot), tinea corporis (ringworm) and candidiasis.

DOSAGE AND ADMINISTRATION

For external use only.

Clean the skin before applying this medicine. Apply a thin layer to completely cover the affected areas once or twice daily and rub well to ensure good absorption. Continue to use this medicine for at least one week after disappearance of symptoms to completely destroy the fungus which may still be present.

For nail infections, apply daily and cover with bandage.

CONTRAINDICATIONS

Do not use for other skin problems without first checking with your doctor. Topical hydrocortisone should not be used in patients with tuberculosis and herpes simplex infections and also in patients vaccinated with live vaccine.

WARNINGS AND PRECAUTIONS

Inform your doctor if you have any previous unusual or allergic reactions to any topical preparation. Use with care on children especially when applied to large areas or with occlusive dressing.

Do not bandage or otherwise cover the area of skin being treated unless directed by the doctor. Such an occlusive dressing (airtight covering) may increase both the amount of steroid absorbed through the skin as well as increase the risk of side effect. Avoid contact with your eyes. It is advisable to consult a doctor if no improvement is observed in 4 weeks in the case of 'athlete's foot' or 2 weeks in 'jock itch'.

In patients on warfarin, caution should be exercised and the anticoagulant effect should be monitored (see Interactions).

Visual disturbance may be reported with systemic and topical corticosteroid use. If a patient presents with symptoms such as blurred vision or other visual disturbances, the patient should be considered for referral to an ophthalmologist for evaluation of possible causes which may include cataract, glaucoma or rare diseases such as central serous chorioretinopathy (CSCR) which have been reported after use of systemic and topical corticosteroids.

Use in infants

In infants, long term continuous topical corticosteroid therapy should be avoided.

PREGNANCY AND LACTATION

Should be used cautiously during pregnancy and lactation period. Avoid treatment of large surfaces and application under occlusive dressing.

INTERACTIONS

Miconazole administered systemically is known to inhibit CYP2C9 enzyme system. Due to the limited systemic availability after topical application, clinically relevant interactions occur very rarely. In patients on warfarin which is subjected to metabolism by CYP2C9, caution should be exercised and the anticoagulant effect should be monitored. (See Warnings and Precautions).

SIDE EFFECTS

Check with your doctor if your skin problems become worse or if there is redness, swelling, pain, itch, stinging or irritation at the site of application.

The following side effects may occur if you use this medicine for a long time: acne or oily skin; filling or rounding out of the face; reddish stretch marks; thinning of skin with easy bruising; unusual weakness; swelling of the feet or legs; wounds that will not heal; persistent sore throat or fever; vision blurred.

OVERDOSAGE AND TREATMENT

Overdose of corticosteroid may cause adrenal atrophy and show signs of Cushing's syndrome. Treat symptomatically and withdraw treatment.

SHELF-LIFE

The expiry date is indicated on the packaging.

PRESENTATION

White, water miscible cream in tubes of 15 g.

STORAGE

Store at 2°C - 8°C.

KEEP OUT OF REACH OF CHILDREN

JAUHI DARI KANAK-KANAK

For further information, please consult your pharmacist or physician.

Revision Date: 01-Mar-2024

Manufacturer and Product Registration Holder

Xepa-Soul Pattinson (Malaysia) Sdn Bhd

1-5, Cheng Industrial Estate, 75250 Melaka, Malaysia.