

PRODUCT LITERATURE

COREDIN EXPECTORANT LINCTUS

Each 5 ml contains:

Ammonium Chloride	135 mg
Diphenhydramine HCl	14 mg

Preservative

Methyl Paraben	0.1 %w/v
Propyl Paraben	0.01 %w/v

Description

Brown- coloured clear syrup with sweet taste and fruity odour.

Pharmacodynamics Properties

Ammonium chloride has an irritant effect on the gastric mucous membrane. Diphenhydramine has pronounced sedative properties. It also has anti-emetic, anticholinergic and local anaesthetic properties.

Pharmacokinetics Properties

Ammonium chloride is effectively absorbed from the gastro-intestinal tract. The ammonium ion is converted into urea in the liver. The anion liberated into the blood stream and extra-cellular fluid causes a metabolic acidosis and decreases the pH of the urine. This is followed by transient diuresis.

Antihistamines are readily absorbed from the gastro intestinal tract, metabolised in the liver and excreted usually mainly as metabolites in the urine.

Indication

For the treatment of cough and upper respiratory congestion due to cold and allergy.

Recommended Dose

To be taken 3 or 4 times a day

Adult : 1-2 teaspoonful (5ml - 10ml)

Children (6-12 years): 1/2 - 1 teaspoonful (2.5 ml - 5 ml)

Route of Administration

Oral

Contraindications

- Ammonium chloride is contraindicated in the presence of impaired hepatic or renal function.
- Monoamine oxidase inhibitors (MAOIs) or within 14 days of stopping treatment
- Hypersensitivity to the active substances(s) or any of the excipients
- Not to be used in children under the age of 6 years

Warnings and Precautions

When used for treatment of cough and cold:

- (i) Not to be used in children less than 2 years of age
- (ii) To be used with caution and doctor's / pharmacist's advice in children 2 to 6 years of age.
- (iii) May cause drowsiness in susceptible individual.

- Do not exceed stated dose
- Use with caution in prostatic hypertrophy, urinary retention, susceptibility to angle closure, hepatic disease.

- The product may cause drowsiness. This product should not be used to sedate a child.
- Ask a doctor before use if you suffer from a chronic or persistent cough, if you have asthma, suffering from an acute asthma attack or where cough is accompanied by excessive secretions.
- Patients with rare hereditary problems of fructose intolerance, glucose galactose malabsorption or sucrase-isomaltase insufficiency should not take this medicine.

Interactions with Other Medicaments

- CNS depressants: May enhance the sedative effects of CNS depressants including barbiturates, hypnotics, opioid analgesics, anxiolytic sedatives, antipsychotics and alcohol.
- Antimuscarinic drugs: May have an additive muscarinic action with other drugs, such as atropine and some antidepressants.
- MAOIs: Not to be used in patients taking MAOIs or within 14 days of stopping treatment as there is a risk of serotonin syndrome.

Pregnancy and lactation

- This medicine should be taken under medical supervision.

Side Effects

Common side effects:

- *CNS effects:* Drowsiness (usually diminishes within a few days), paradoxical stimulation, headache, psychomotor impairment.
- *Antimuscarinic effects:* Urinary retention, dry mouth, blurred vision, gastrointestinal disturbances, thickened respiratory tract secretions.

Rare side-effects:

Hypotension, extrapyramidal effects, dizziness, confusion, depression, sleep disturbances, tremor, convulsions, palpitation, arrhythmia, hypersensitivity reactions, blood disorders and liver dysfunction.

ADR: Urticaria, Cough

Symptoms and Treatment of Overdose

Ammonium chloride: large doses may cause nausea, vomiting, thirst, headache, hyperventilation and progressive drowsiness and lead to profound acidosis and hypokalaemia.

Diphenhydramine: Antihistamines may cause gastrointestinal disturbances, headache, blurred vision, tinnitus, elation or depression, irritability, nightmares, anorexia, difficulty in micturition, dryness of the mouth, tightness of the chest and tingling, heaviness and weakness of the hands.

Packing

Packed in plastic container of 90 ml and 100 ml.

Pharmaceutical Precautions

Store below 30°C in a dry place, protected from direct light.

Keep away from reach of children.

Jauhkan daripada kanak-kanak

SHAKE WELL BEFORE USE.

Expiry Date

2 years from date of manufacture

Name and Address of Distributor:

ZONTRON PHARMACEUTICALS SDN.BHD. (445695-T)

Lot 10 & 11, PERDA Industrial Park,
Lorong IKS Simpang Ampat B,
14100 Simpang Ampat, S.P.S.,
Pulau Pinang, Malaysia

Name and Address of Product Registration Holder / Manufacturer:

TERAPUTICS SDN. BHD. (590500-W)

Lot 10 & 11, PERDA Industrial Park,
Lorong IKS Simpang Ampat B,
14100 Simpang Ampat, S.P.S.,
Pulau Pinang, Malaysia

Malaysian Drug Registration No.

MAL19880310AZ

Date of Revision: 20/06/2024